



<http://ir4.rutgers.edu/Ornamental/ornamentalSummaryReports.cfm>

IR-4 Ornamental Horticulture Program Dimethenamid-p Crop Safety

**Authors: Ely Vea and Cristi L. Palmer
Date: 12/20/2016**

Acknowledgements

**Susan Bierbrunner
Kathleen Hester
Diane Infante
Lori Harrison
Karen Sims**

This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2015-34383-23710 with substantial cooperation and support from the State Agricultural Experiment Stations and USDA-ARS.

Table of Contents

Table of Contents	2
Table of Tables	3
Abstract	4
Introduction.....	5
Materials and Methods.....	5
Results and Summary	5
Phytotoxicity	5
Label Suggestions	34
Appendix 1: Contributing Researchers.....	35

Table of Tables

Table 1.	List of Tower EC treated crops with no or minimal transitory injury.	6
Table 2.	List of Tower EC treated crops with no or minimal transitory injury seen at the 1X rate, but the 2X or 4X rate did cause significant phytotoxicity	7
Table 3.	List of Tower EC treated crops exhibiting significant injury.	7
Table 4.	List of Tower EC treated crops where more information is needed.	8
Table 5.	Detailed Summary of Crop Safety Testing with Dimethenamid-p.....	9

Abstract

From 2007 to 2016, IR-4 completed 504 trials on Tower EC (dimethenamid-p). The data contained in this report was generated to register uses of dimethenamid-p on and around ornamental horticulture plants with over-the-top applications. The dimethenamid-p rates in the testing program were 0.97, 1.94 and 3.88 pounds active ingredient per acre (lb ai per A) as the 1X, 2X and 4X rates. Tower EC had been applied to 146 plant genera or species. Of these, 62 plant species exhibited no or minimal transient injury after application at all three rates. Twenty crops exhibited no phytotoxicity at 0.97 lb ai per acre but did have some injury at 1.94 and 3.88 lb ai per acre. Nine crops – *Aquilegia sp.*, *Catharanthus roseus*, *Cladrastis sp.*, *Echeveria sp.*, *Echinacea sp.*, *Epilobium canum*, *Muhlenbergia dubia*, *Teucrium chamaedrys* and *Viburnum opulus* – exhibited significant phytotoxicity at even the lowest rate.

Introduction

Control of broadleaved weeds and sedges in the production of woody and herbaceous perennials can be problematic because nurseries grow many different types of plants and not all genera or species are listed on labels. These weeds can also be difficult to control in landscape settings for the same reason. Four herbicides, Freehand G (BAS 659 G; dimethenamid-p + pendimethalin), F6875 0.3G (sulfentrazone + proflamone), Mesotrione SC, and Tower EC (BAS 656 EC; dimethenamid-p), were chosen for 2007 research activities into level of crop safety on over 38 different plant species. During 2008, 2009 and 2010, Freehand (dimethenamid-p + pendimethalin), Broadstar 0.25G VC1604 (flumioxazin), EXC3898 G (mesotrione), Tower EC and V-10142 G (imazasulfuron) were tested. This summary covers the results for Tower EC from 2007 to 2016.

Materials and Methods

In the 2007 protocol, two applications of Tower EC were made approximately 8 weeks apart. In the protocols between 2008 and 2016 protocols, two applications of Tower EC were made approximately 6 weeks apart. The application rates were 0.97, 1.94 and 3.88 lb ai per acre, plus a water treated control. A minimum of four plants (replicate treatments) were required with many researchers exceeding this minimum. Phytotoxicity was recorded on a scale of 0 to 10 (0 = No phytotoxicity; 10 = Complete kill) at 1, 2, 4, 8, and 12 weeks after initial application. Some researchers also included readings 3 to 4 days after the initial and second applications. The following protocols were used: 07-009, 08-010, 09-011, 10-001, 11-004, 12-014, 13-014, 14-009, 15-009 and 16-010. Please visit <http://ir4.rutgers.edu/ornamental/OrnamentalDrafts.cfm> to view and download these protocols.

Tower EC was supplied to researchers (See list of researchers in Appendix 1) by BASF Corporation.

Results and Summary

Phytotoxicity

Based on the type and nature of injury seen with Tower EC applications in the conducted research, tested plant species were placed into four categories: 1) no significant phytotoxicity or growth differences from the untreated check or any injury was transitory, 2) no or minimal transitory injury seen at the 1X rate, but the 2X and/or 4X rates did cause significant phytotoxicity, 3) significant injury sufficient to recommend growers not utilize this product, and 4) more data is needed to make informed recommendations.

Tower EC exhibited no or minimal negative impact on 62 plant species with over the top applications (Table 1). Some minimal injury may be acceptable for growers if applications are made several weeks to months in advance of crop sale particularly for woody ornamental crops. In the research presented here, 20 plant species exhibited significant injury at higher rates even though little injury was observed at 1X (Table 2). Nine crops tested from 2007 to 2016 exhibited damage sufficient to recommend growers not utilize Tower EC as an over-the-top treatment for pre-emergent weed control (Table 3). For 52 genera/species, more information is needed either

because only 1 or 2 trials were conducted or because consistent results were not achieved among the research sites (Table 4).

Please see Table 5 for a list of individual trial summaries on Tower EC.

Table 1. List of Tower EC treated crops with no or minimal transitory injury.

<i>Abelia grandiflora</i> (see Neal) ¹	<i>Leucanthemum maximum</i>
<i>Abies fraseri</i> ¹	<i>Leucothoe</i> sp. ¹
<i>Acer palmatum</i> ¹	<i>Loropetalum</i> sp. (See Neal) ¹
<i>Acer rubrum</i> ¹	<i>Magnolia</i> sp. ¹
<i>Agapanthus africanus</i> ¹	<i>Malus domestica</i> ¹
<i>Agave</i> sp. ¹	<i>Nandina domestica</i> ¹
<i>Berberis</i> sp. (See Lieth, Mathers) ¹	<i>Ophiopogon japonicas</i> ¹
<i>Betula nigra</i> ¹	<i>Paeonia lactiflora</i>
<i>Buxus</i> sp. (See Mathers) ¹	<i>Picea abies</i> ¹
<i>Callistemon</i> sp. ¹	<i>Picea glauca</i> ¹
<i>Camelia</i> sp. ¹	<i>Picea pungens</i> ¹
<i>Carex</i> sp. ¹	<i>Pieris japonica</i> ¹
<i>Caryopteris x clandonensis</i> (See Beste/Frank) ¹	<i>Pinus</i> sp. ¹
<i>Ceanothus</i> sp. ¹	<i>Potentilla fruticosa</i> ¹
<i>Clematis</i> sp. ¹	<i>Pseudotsuga menziesii</i> ¹
<i>Cotoneaster</i> sp. ¹	<i>Quercus acutissima</i> ¹
<i>Delosperma</i> sp. ¹	<i>Raphiolepis indica</i> (See Lieth) ¹
<i>Dendranthema x morifolium</i> (See Klett 2013) ^{1,2}	<i>Rosa</i> sp. ¹
<i>Euonymus</i> sp.	<i>Sabal minor</i>
<i>Gaura lindheimeri</i>	<i>Spiraea</i> sp. ¹
<i>Gleditsia</i> sp. ¹	<i>Taxus</i> sp. ¹
<i>Hemerocallis</i> sp. ¹	<i>Ternstroemia</i> sp.
<i>Hosta</i> sp. ¹	<i>Thuja occidentalis</i> (See Lieth, Mathers) ¹
<i>Hydrangea macrophylla</i> ¹	<i>Thuja orientalis</i> ¹
<i>Hydrangea paniculata</i> ¹	<i>Thuja plicata</i> ¹
<i>Hydrangea quercifolia</i> ¹	<i>Trachelospermum jasminoides</i> ¹
<i>Ilex cornuta</i> ¹	<i>Tsuga heterophylla</i> (See Beste/Frank) ¹
<i>Juglans nigra</i> ¹	<i>Viburnum plicatum</i> (See Beste/Frank) ¹
<i>Juniperus</i> sp. ¹	<i>Viburnum tinus</i> (See Neal) ¹
<i>Lagerstroemia indica</i> ¹	<i>Wisteria</i> spp. ¹
<i>Lantana</i> sp. ¹	
<i>Lavandula</i> sp. ¹	

¹ Registered already on EPA label

² More information on cultivar and or species differences is warranted.

Table 2. List of Tower EC treated crops with no or minimal transitory injury seen at the 1X rate, but the 2X or 4X rate did cause significant phytotoxicity

<i>Buddleia davidii</i> ^{1,2}	<i>Iberis sp.</i>
<i>Clethra alnifolia</i>	<i>Ligustrum sp.</i> ¹
<i>Cornus florida</i> ¹	<i>Nepeta cataria</i>
<i>Cornus kousa</i> ¹	<i>Nepeta x faasseni</i>
<i>Cornus sericea</i> ¹	<i>Quercus sp.</i> ^{1,2}
<i>Cotoneaster dammeri</i>	<i>Rhododendron sp.</i> ^{1,2}
<i>Cryptomeria japonica</i> ¹	<i>Salvia nemorosa/sylvestris</i> ¹
<i>Dryopteris sp.</i> ^{1,2}	<i>Scaevola sp.</i> ²
<i>Forsythia sp.</i> ^{1, 3}	<i>Syringae sp.</i> ¹
<i>Hibiscus sp.</i> ^{1,2}	<i>Viburnum nudum</i> ¹

Table 3. List of Tower EC treated crops exhibiting significant injury.

<i>Aquilegia sp.</i> ⁴	<i>Epilobium canum</i> ¹
<i>Catharanthus roseus</i>	<i>Mulenbergia dubia</i> ¹
<i>Cladrastis sp.</i> ³	<i>Teucrium chamaedrys</i>
<i>Echeveria sp.</i>	<i>Viburnum opulus</i> ¹
<i>Echinacea sp.</i> ⁴	

Table 4. List of Tower EC treated crops where more information is needed.

<i>Acer negundo</i> ¹	<i>Fothergila gardenii</i>	<i>Philadelphus viginalis</i>
<i>Acer saccharinum</i> ¹	<i>Fraxinus americana</i> ¹	<i>Photinia fraserii</i>
<i>Albizia julibrissin</i>	<i>Gardenia sp.</i>	<i>Photinia villosa</i>
<i>Aloe sp.</i>	<i>Heteromeles arbutifolia</i>	<i>Quercus alba</i> ¹
<i>Arctostaphylos uva ursi</i>	<i>Heuchera sanguinea</i>	<i>Quercus pinnata</i> ¹
<i>Asclepias sp.</i>	<i>Ilex x meserveae</i> ¹	<i>Quercus rubra</i> ¹
<i>Berlandiera lyrata</i>	<i>Ilex verticillata</i> ¹	<i>Ribes viburnifolium</i>
<i>Betula alba</i> ¹	<i>Iris sp.</i> ¹	<i>Rudbeckia sp.</i>
<i>Caladium sp.</i>	<i>Itea virginica</i>	<i>Sedum sp.</i> ⁴
<i>Chasmanthium latifolium</i>	<i>Kerria japonica</i>	<i>Stewartii pseudocamelia</i>
<i>Cornus amomum</i> ¹	<i>Lonicera sp.</i>	<i>Syringa sp.</i> ^{1,2}
<i>Crassula ovata</i>	<i>Mammillaria sp.</i>	<i>Tulipa sp.</i>
<i>Crassula sp.</i>	<i>Muhlenbergia capillaris</i>	<i>Viburnum odoratissimum</i> ¹
<i>Crataegus coccinoid</i>	<i>Myrica pensylvanica</i>	<i>Viburnum x pragense</i> ¹
<i>Cupressus arizonica</i>	<i>Narcissus sp.</i> ¹	<i>Washingtonia robusta</i>
<i>Echinacea sp.</i> ²	<i>Osmanthus heterophyllus</i> ¹	<i>Zelkova serrata</i>
<i>Echinocactus grusonii</i>	<i>Osmundo regalis</i>	
<i>Fagus sp.</i>	<i>Penstemon sp.</i>	

¹ Registered already on EPA master label

² More information on cultivar differences might be warranted.

³ Perhaps a single application only could be a label restriction.

⁴ Listed in the label under sensitive ornamental plants

Table 5. Detailed Summary of Crop Safety Testing with Dimethenamid-p

Notes: Table entries are sorted by crop Latin name. Only those trials with research reports received by 12/12/16 are listed below.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27074	Abelia (Abelia sp.) A. gaucheri	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
27074	Abelia (Abelia sp.) A. grandiflora	Field Container	Neal	NC	2009	Over the top	No significant injury at 0.97, moderate stunting at 1.94 and 3.88 lb ai per acre.
27074	Abelia (Abelia sp.) A. grandiflora	Field Container	Neal	NC	2009	Over the top	No significant injury at 0.97, slight to moderate stunting at 1.94 and 3.88 lb ai per acre.
27074	Abelia (Abelia sp.) A. grandiflora 'Edward Goucher'	Field Container	Trader	MS	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27074	Abelia (Abelia sp.) 'Edward Goucher'	Field Container	Lieth	CA	2008	Over the top	Slight, acceptable injury and no significant growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26537	Fir, Fraser (Abies fraseri)	Field In-Ground	Ahrens/Mervosh	CT	2007	Over the top	No injury at 0.97 and 1.94 lb ai per acre; moderate injury only after 1st application, with complete recovery, at 3.88 lb ai per acre
26537	Fir, Fraser (Abies fraseri)	Field In-Ground	Beste/Frank (ARS)	MD	2007	Over the top	Results not useful because of severe injury caused by very high temperature and drought conditions
26320	Fir, Fraser (Abies fraseri)	Field Container	Boydston	WA	2007	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26320	Fir, Fraser (Abies fraseri)	Field Container	Freiberger	NJ	2007	Over the top	Slight injury at 0.97, 1.94 and 3.99 lb ai per acre after 1st, moderate after 2nd application
26320	Fir, Fraser (Abies fraseri) 'Roan Mountain'	Field Container	Marshall	MI	2007	Over the top	One application. No injury at 0.97, 1.94 and 3.88 lb ai per acre
27808	Box elder (Acer negundo L. ssp. Negundo)	Field Container	Freiberger	NJ	2008	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre.
27808	Box elder (Acer negundo L. ssp. Negundo)	Field Container	Freiberger	NJ	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27808	Box elder (Acer negundo L. ssp. Negundo)	Field Container	Jones	OH	2013	Over the top	No injury or significant growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27082	Maple, Japanese (Acer palmatum) 'Atropurpureum'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; reduced marketability for all treatments because of environmental stress
27082	Maple, Japanese (Acer palmatum) 'Atropurpureum'	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable.
26541	Maple, Red (Acer rubrum)	Field In-Ground	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
26194	Maple, Red (Acer rubrum)	Field Container	Freiberger	NJ	2007	Over the top	No injury at 0.97, 1.94 and 3.99 lb ai per acre after 1st, slight after 2nd application
26194	Maple, Red (Acer rubrum)	Field Container	Senesac	NY	2007	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26194	Maple, Red (<i>Acer rubrum</i>) 'Summer'	Field Container	Gilliam	AL	2008	Over the top	No injury after the first application using 0.97, 1.84 and 3.88 lb ai per acre, but no to moderate injury after the second application increasing with rate; there was also a significant reduction in growth at the highest rate.
27091	Maple (<i>Acer</i> sp.) <i>A. palmatum</i>	Field Container	Ahrens/Mervosh	CT	2009	Over the top	No crop injury or reduction in growth with two applications at 0.97, 1.94, 3.88 lb ai per acre.
27091	Maple (<i>Acer</i> sp.) <i>A. rubrum</i> 'Sun Valley'	Field Container	Mathers (OSU)	OH	2008	Over the top	Significant injury at 0.97, 1.94 and 3.88 lb ai per acre after 1st, none after 2nd application
27091	Maple (<i>Acer</i> sp.) <i>A. saccharinum</i>	Field Container	Derr	VA	2009	Over the top	No significant injury at 0.97 and 1.94, significant at 3.88 lb ai per acre; no growth reduction.
27091	Maple (<i>Acer</i> sp.) Maple Japanese	Field Container	Harvey	WA	2009	Over the top	No injury at 0.97 and 1.94, slight at 3.88 lb ai per acre.
31084	Bearberry (<i>Arctostaphylos</i> sp.) <i>A. uva ursi</i> 'Massachusetts'	Field Container	Senesac	NY	2016	Over the top	Minor to moderate injury with complete recovery with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28201	Ohio Buckeye (<i>Aesculus glabra</i>)	Field Container	Siefer	OH	2015	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28468	Lily-Of-The-Nile (<i>Agapanthus</i> sp.) <i>A. africanus</i> 'Big Blue'	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
28468	Lily-Of-The-Nile (<i>Agapanthus</i> sp.) <i>A. africanus</i> 'Peter Pan'	Field Container	Uber	CA	2008	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
28468	Lily-Of-The-Nile (<i>Agapanthus</i> sp.) 'Dwarf Blue'	Field Container	Wilén	CA	2012	Over the top	Slight, acceptable injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
31325	Agave (<i>Agave</i> sp.) <i>A. ellemetiana</i> 'Santina'	Field Container	Wilén	CA	2013	Over the top	No to minor injury with 21, 42 and 84 fl oz per acre after 1st applic., unacceptable after 2nd application.
31325	Agave (<i>Agave</i> sp.) <i>A. parrasana</i>	Field Container	Villavicencio	CA	2013	Over the top	No injury with 21, minor with 42 and 84, fl oz per acre applied twice; no growth reduction.
31325	Agave (<i>Agave</i> sp.) 'Blue Flame'	Field Container	Villavicencio	CA	2012	Over the top	No injury with 21 and 42 fl oz per acre applied twice, slight discoloration and spotting only with 84 fl oz; no growth reduction.
29172	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Czarnota	GA	2013	Over the top	Results inconclusive due to high injury at 21 fl oz per acre, and moderate injury, but not significant from untreated, with 42 and 84 fl oz.
29172	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
31324	Barbados Aloe (<i>Aloe</i> sp.) <i>A. brevifolia</i>	Field Container	Villavicencio	CA	2013	Over the top	Severe injury (leaf splitting) with 21,42 and 84 fl oz per acre applied twice.
31324	Barbados Aloe (<i>Aloe</i> sp.) 'Blue Elf'	Field Container	Villavicencio	CA	2012	Over the top	No injury with 21 fl oz per acre applied twice, slight necrosis with 24 and 84 fl oz; no growth reduction.
31324	Barbados Aloe (<i>Aloe</i> sp.) 'Little Gator'	Field Container	Wilén	CA	2014	Over the top	Slight injury with 21, 42 and 84 fl oz per acre; no significant growth reduction.
28222	Canadian Serviceberry (<i>Amelanchier canadensis</i>)	Field In-Ground	Cochran	IA	2014	Over the top	Moderate injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice; slight growth reduction with 4X.
28222	Canadian Serviceberry (<i>Amelanchier canadensis</i>)	Field In-Ground	Cochran	IA	2015	Over the top	Minor injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice; no growth reduction.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
28222	Canadian Serviceberry (<i>Amelanchier canadensis</i>)	Field In-Ground	Persad	OH	2014		No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
29533	Columbine (<i>Aquilegia</i> sp.)	Field Container	Boydston	WA	2013	Over the top	Very slight necrosis with full recovery at 0.98, 1.97, and 3.94 lb ai per acre applied twice; no growth reduction; all plants saleable.
29533	Columbine (<i>Aquilegia</i> sp.) A. <i>canadensis</i> 'Little Lanterns'	Field Container	Senesac	NY	2014	Over the top	Severe injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
29533	Columbine (<i>Aquilegia</i> sp.) A. <i>chrysantha</i>	Field Container	Klett	CO	2013	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
29533	Columbine (<i>Aquilegia</i> sp.) A. <i>vulgaris</i> 'Origami Mix'	Field Container	Boydston	WA	2014	Over the top	No significant injury with 0.98 and 1.97, moderate to severe injury (stunting and leaf burn) with 3.94 lb ai per acre applied twice; 1X- and 2X-treated plants marketable.
29975	Butterfly Flower (<i>Asclepias</i> sp.) A. <i>incarnata</i>	Field Container	Senesac	NY	2014	Over the top	Slight injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
29975	Butterfly Flower (<i>Asclepias</i> sp.) A. sp.	Field Container	Derr	VA	2010	Over the top	Little to no crop injury with 1.0 lb ai per acre. 62 DAT and 14DAT2. Fair to v. good control of longstalked phyllanthus, tasselflower, s. crabgrass.
29975	Butterfly Flower (<i>Asclepias</i> sp.) A. <i>tuberosa</i>	Field Container	Beste/Frank (ARS)	MD	2011	Over the top	Acceptable injury, but moderate to severe growth reduction increasing with rates (0.98, 1.97 and 3.94 lb ai per acre) applied twice.
29975	Butterfly Flower (<i>Asclepias</i> sp.) A. <i>tuberosa</i>	Field Container	Klett	CO	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice; high growth reduction 2X and 4X.
32953	Fern, Sprengeri (<i>Asparagus densiflorus</i>)	Field Container	Rivera	PR	2014	Over the top	Minor initial injury with quick recovery at 0.98, 1.97 and 3.94 lb ai per acre applied twice.
26283	Barberry (<i>Berberis</i> sp.)	Field Container	Harvey	WA	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26283	Barberry (<i>Berberis</i> sp.) B. <i>thunbergii</i>	Field Container	Uber	CA	2010	Over the top	No significant injury at 0.97, slight to moderate at 1.94 and 3.88 lb ai per acre; no significant growth reduction.
26283	Barberry (<i>Berberis</i> sp.) B. <i>thunbergii</i> 'Amber Glow'	Field Container	Uber	CA	2008	Over the top	Results inconclusive due to environmental stress; no significant injury at 0.97, 1.94 and 3.88 lb ai per acre after 1st application
26283	Barberry (<i>Berberis</i> sp.) B. <i>thunbergii</i> 'Atropurpurea'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
26283	Barberry (<i>Berberis</i> sp.) B. <i>thunbergii</i> atropurpureum 'Crimson Pygmy'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	Very slight injury at 0.97, 1.94 and 3.88 lb ai per acre; growth reduction at 2X and 4X but all plants marketable
26283	Barberry (<i>Berberis</i> sp.) B. <i>thunbergii</i> 'Crimson Pigmy'	Field Container	Lieth	CA	2008	Over the top	Unacceptable injury and growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
30152	Japanese Barberry (<i>Berberis thunbergii</i>) B. 'Crimson Pygmy'	Greenhouse	Mathers (OSU)	MI	2010	Over the top	Lincoln: Unacceptable crop injury with 32 oz/a.
31085	Chocolate Flower (<i>Berlandiera lyrata</i>)	Field Container	Mansue	NJ	2014	Over the top	Acceptable injury with 21, unacceptable injury and stunting with 42 and 84 fl oz per 100 gal applied twice.
29171	Birch, River (<i>Betula nigra</i>)	Field Container	Beste/Frank (ARS)	MD	2010	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice; no reduction in marketability.
29171	Birch, River (<i>Betula nigra</i>)	Field Container	Beste/Frank (ARS)	MD	2011	Over the top	No significant injury, growth or marketability reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
29171	Birch, River (<i>Betula nigra</i>)	Field Container	DeFrancesco	OR	2011	Over the top	No significant injury after 1st application with 21 and 42 fl oz per acre., moderate, with good recovery, with 84 fl oz; no significant injury after 2nd application; no growth reduction.
29171	Birch, River (<i>Betula nigra</i>)	Field Container	Neal	NC	2010	Over the top	No crop injury with any rate at any time during the evaluation period.
29171	Birch, River (<i>Betula nigra</i>) <i>B. alba</i>	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
26335	Butterfly Bush (<i>Buddleia davidii</i>) 'Nanho Purple'	Field Container	Wade	SC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26335	Butterfly Bush (<i>Buddleia davidii</i>) 'Royal Red'	Field Container	Fraelich	GA	2007	Over the top	Moderate injury at 0.97, severe at 1.94 and 3.88 lb ai per acre
26335	Butterfly Bush (<i>Buddleia davidii</i>) 'White Ball'	Field Container	Marshall	MI	2007	Over the top	One application. No significant injury at 0.97, 1.94 and 3.88 lb ai per acre
30151	Boxwood, Japanese (<i>Buxus microphylla</i>) <i>B. microphylla</i> 'Green velvet'	Greenhouse	Mathers (OSU)	MI	2010	Over the top	Lincoln: Unacceptable crop injury with 32 oz/a.
26305	Boxwood (<i>Buxus</i> sp.) <i>B. microphylla</i> 'Faulkner'	Field Container	Klett	CO	2007	Over the top	Two trials; no injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26305	Boxwood (<i>Buxus</i> sp.) <i>B. sinica</i> var. <i>insularis</i> 'Wintergreen'	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26305	Boxwood (<i>Buxus</i> sp.) 'Green Velvet'	Field Container	Marshall	MI	2007	Over the top	One application. No injury at 0.97, 1.94 and 3.88 lb ai per acre
29976	Elephant's-Ear, Angel-Wings (<i>Caladium</i> sp.) <i>C. 'Florida Cardinal'</i>	Field Container	Derr	VA	2010	Over the top	No crop injury with 1.0 lb ai per acre. Fair to v. good control of longstalked <i>phyllanthus</i> , tasselflower, s. crabgrass.
28727	Bottlebrush (<i>Callistemon</i> sp.)	Field Container	Beste/Frank (ARS)	MD	2010	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice; no reduction in marketability.
28727	Bottlebrush (<i>Callistemon</i> sp.)	Field Container	Denny	MS	2013	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28727	Bottlebrush (<i>Callistemon</i> sp.)	Field Container	Lieth	CA	2010	Over the top	No injury or growth reduction with 0.97, 1.94 and 3.88 lb ai per acre applied twice.
28727	Bottlebrush (<i>Callistemon</i> sp.) 'Little John'	Field Container	Wilén	CA	2012	Over the top	Minor injury with 21, 42 and 84 fl oz per acre after 1st applic; acceptable injury and growth reduction with 1X and 2X, unacceptable with 4X after 2nd applic.
28727	Bottlebrush (<i>Callistemon</i> sp.) 'Little John'	Field Container	Wilén	CA	2012	Over the top	Slight, acceptable injury and growth reduction with 21 and 42, moderate with 84, fl oz per acre applied twice.
27109	Camellia (<i>Camellia</i> sp.) <i>C. japonica</i>	Field Container	Wade	GA	2009	Over the top	No injury or significant growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27109	Camellia (<i>Camellia</i> sp.) <i>C. japonica</i> 'Colonel Fiery'	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
27109	Camellia (<i>Camellia</i> sp.) <i>C. sasanqua</i> 'White'	Field Container	Trader	MS	2008	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27109	Camellia (<i>Camellia</i> sp.) <i>C. vernalis</i> 'Yuletide'	Field Container	Gilliam	AL	2009	Over the top	No injury or growth reduction at 0.97, 1.84 and 3.88 lb ai per acre.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
28707	Sedge, Grassland (<i>Carex divulsa</i>)	Field Container	Wilén	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre; root dry weight significantly reduced at all rates.
28707	Sedge, Grassland (<i>Carex divulsa</i>)	Field Container	Wilén	CA	2010	Over the top	All plants were marketable. Minor chlorosis 4WAT2 with 0.97, 1.94, and 3.88 lb ai per acre increasing with rate.
28707	Sedge, Grassland (<i>Carex divulsa</i>) C. oshimensis 'Evergold'	Field Container	Boydston	WA	2010	Over the top	Two sequential applications of Tower 63.9% EC applied 6 weeks apart at 0.97, 1.94, and at 3.88 lb ai per acre did not injure sedge plants.
28224	Pecan (Non-Bearing) (<i>Carya illinoensis</i>)	Field In-Ground	Cochran	IA	2015	Over the top	Minor injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice; no growth reduction.
27420	Caryopteris (<i>Caryopteris x clandonensis</i>)	Field Container	Neal	NC	2008	Over the top	No injury at 0.97, 1.94, and 3.88 lb ai per acre.
27420	Caryopteris (<i>Caryopteris x clandonensis</i>) 'Longwood Blue'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury or flower number reduction, significant growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
27420	Caryopteris (<i>Caryopteris x clandonensis</i>) 'Sunshine Blue'	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
28202	Spanish Chestnut (<i>Castanea sativa</i>)	Field Container	Cochran	IA	2015	Over the top	Minor injury with 0.98 and 1.97, moderate with 3.94 lb ai per acre applied twice; no growth reduction.
28203	Rose Periwinkle (<i>Catharanthus roseus</i>)	Field Container	Mansue	NJ	2015	Over the top	Moderate injury (petals bleaching and falling) with 21, 42 and 84 fl oz per acre applied twice; no growth reduction.
28203	Rose Periwinkle (<i>Catharanthus roseus</i>) 'Pacifica XP Polka Dot'	Field Container	Senesac	NY	2014	Over the top	Severe injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28731	Ceanothus, maritime (<i>Ceanothus maritimus</i>)	Field Container	Grunwald	OR	2010	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all treated plants saleable.
28731	Ceanothus, maritime (<i>Ceanothus maritimus</i>)	Field Container	Lieth	CA	2011	Over the top	Results not conclusive because of suspected product overdose for all treatments; researcher suggested to repeat trial.
28731	Ceanothus, maritime (<i>Ceanothus maritimus</i>) 'Dark Star'	Field Container	Wilén	CA	2010	Over the top	No crop injury 28 DAT with one application 21, 42, or 84 oz/A but unacceptable injury with two applications for all rates.
29236	Ceanothus sp. (<i>Ceanothus</i> sp.) 'Concha'	Field Container	Wilén	CA	2009	Over the top	No significant injury at 0.97 and 1.94, moderate at 3.88 lb ai per acre; root and shoot dry weights significantly reduced at all rates.
28735	Ceanothus (<i>Ceanothus x pallida</i>)	Field Container	Lieth	CA	2010	Over the top	No injury with 0.97 and 1.94, very slight with 3.88 lb ai per acre, applied twice; no growth effects.
28735	Ceanothus (<i>Ceanothus x pallida</i>) C. americanus	Field Container	Czarnota	GA	2013	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
28735	Ceanothus (<i>Ceanothus x pallida</i>) 'Marie Simon'	Field Container	Grunwald	OR	2010	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all treated plants saleable.
31087	Northern Sea Oats, Wild Oats (<i>Chasmanthium latifolium</i>)	Field Container	Hanson	CA	2013	Over the top	No injury after 1st applic., slight to moderate injury (leaf necrosis) with 0.97, 1.94 and 3.88 lb ai per acre after 2nd applic.; moderate growth reduction with 2X and 4X.
31087	Northern Sea Oats, Wild Oats (<i>Chasmanthium latifolium</i>)	Field Container	Mansue	NJ	2013	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
31087	Northern Sea Oats, Wild Oats (<i>Chasmanthium latifolium</i>)	Field Container	Senesac	NY	2013	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
31087	Northern Sea Oats, Wild Oats (<i>Chasmanthium latifolium</i>)	Field Container	Wilén	CA	2014	Over the top	Moderate to high injury increasing with rates (21, 42 and 84 fl oz per acre) applied twice.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27820	Chrysanthemum, Garden (Chrysanthemum/Dendranthema sp.)	Field Container	Mansue	NJ	2014	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice.
27820	Chrysanthemum, Garden (Chrysanthemum/Dendranthema sp.) 'Sheffield Pink'	Field Container	Czarnota	GA	2007	Over the top	No significant injury at 0.97 and 1.94 lb ai per acre, significant at 3.88 lb
27820	Chrysanthemum, Garden (Chrysanthemum/Dendranthema sp.) C. x morifolium 'Bolero'	Field Container	Beste/Frank (ARS)	MD	2013	Over the top	No injury or growth reduction or reduction of flower number with 21, 42 and 84 fl oz per acre applied twice; no reduction in marketability.
27820	Chrysanthemum, Garden (Chrysanthemum/Dendranthema sp.) D. zawadskii 'Clara Curtis'	Field Container	Klett	CO	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre after 1st, slight with 4X after 2nd applic.; moderate growth reduction with 4X.
27548	Hardy Mum (Chrysanthemum/Dendranthema x morifolium)	Field Container	Derr	VA	2013	Over the top	No injury or flower rating reduction with 0.98 and 1.97 lb ai per acre applied twice; slight with 3.94 lb.
27548	Hardy Mum (Chrysanthemum/Dendranthema x morifolium) D. zawadskii	Field Container	Klett	CO	2013	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
27548	Hardy Mum (Chrysanthemum/Dendranthema x morifolium) 'Dazzler Stacy'	Field Container	Derr	VA	2007	Over the top	0, 1, and 6 % injury at 1, 2 and 4 lb ai per acre; 100 % control of rice flatsedge and fragrant flatsedge
26316	Yellowwood (Cladrastis sp.) C. kentukea	Field Container	Beste/Frank (ARS)	MD	2011	Over the top	Slight injury with good recovery at 0.98 and 1.97, moderate injury at 3.94 lb ai per acre; no significant growth reduction. Reduced market value at 4X rate.
26316	Yellowwood (Cladrastis sp.) C. kentukea	Field Container	Freiberger	NJ	2007	Over the top	No significant injury at 0.97, 1.94 and 3.99 lb ai per acre after 1st, Minor at 2X and 4X after 2nd application
26316	Yellowwood (Cladrastis sp.) C. kentukea	Field Container	Mathers (OSU)	OH	2011	Over the top	Significant crop injury (leaf burn) with 1.94 and 3.88 lb ai per acre
27115	Clematis (Clematis sp.)	Field Container	Harvey	WA	2009	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre.
27115	Clematis (Clematis sp.) C. integrifolia	Field Container	Klett	CO	2008	Over the top	Trial 1: No injury at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
27115	Clematis (Clematis sp.) C. integrifolia	Field Container	Klett	CO	2008	Over the top	Trial 2: No injury at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
27115	Clematis (Clematis sp.) 'Gillian Blades' and others	Field Container	Boydston	WA	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants saleable.
27120	Summersweet (Clethra alnifolia)	Field Container	Ahrens/Mervosh	CT	2011	Over the top	No crop injury with two sequential applications at 0.97, 1.94, or 2.65 lb ai per acre.
27120	Summersweet (Clethra alnifolia)	Field Container	Gilliam	AL	2010	Over the top	Injury unrelated to treatments (0.97, 1.94, 3.88 lb ai per acre) observed in all plants. Significant injury with 4x at 2WAT2 and stunting with 2x.
27120	Summersweet (Clethra alnifolia) C. alnifolia 'Sixteen Candles'	Field Container	Neal	NC	2010	Over the top	Significant crop injury (necrosis followed by stunting) with 2x and 4x rates (1.94 and 3.88 lb ai per acre).

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27380	Dogwood, Silky (Cornus amomum)	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
21424	Dogwood, Silky (Cornus amomum)	Field In-Ground	Garrett	MD	2001		
27380	Dogwood, Silky (Cornus amomum)	Field Container	Senesac	NY	2011	Over the top	Minor injury with two applications at 0.97, 1.94, and 3.88 lb aia but all plants marketable.
27125	Dogwood, Flowering (Cornus florida)	Field Container	Ahrens/Mervosh	CT	2008	Over the top	Slight injury at 0.98, moderate to severe at 1.97 and 3.94 lb ai per acre after 2nd application; 1st application rates were 3.8 times higher than protocol rates.
27125	Dogwood, Flowering (Cornus florida)	Field Container	Ahrens/Mervosh	CT	2009	Over the top	Two applications at 0.97, 1.94, 3.88 lb ai per acre appear to be safe although 1x caused early injury. Powdery mildew problem made it difficult to rate and suggests tolerance be investigated further.
27125	Dogwood, Flowering (Cornus florida)	Field Container	Freiberger	NJ	2009	Over the top	Slight injury at 0.97 and 1.94. moderate at 3.88 lb ai per acre.
28862	Dogwood, Kousa (Cornus kousa)	Field Container	Ahrens/Mervosh	CT	2009	Over the top	No crop injury with two application at 0.97, 1.94, 3.88 lb ai per acre.
28862	Dogwood, Kousa (Cornus kousa)	Field Container	Ahrens/Mervosh	CT	2011	Over the top	Little to no crop injury with two sequential applications at 0.97, 1.94 or 2.65 lb ai per acre.
28862	Dogwood, Kousa (Cornus kousa)	Field Container	DeFrancesco	OR	2010	Over the top	All rates (0.97, 1.94, 3.88 lb ai per acre) exhibited minor but statistically different from the untreated with one and two applications.
28862	Dogwood, Kousa (Cornus kousa) C. kousa 'Milky Way'	Field Container	Mathers (OSU)	OH	2010	Over the top	Ohio: Data inconclusive because untreated exhibited injury.
28862	Dogwood, Kousa (Cornus kousa) 'Chinensis'	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre; reduced growth and marketability with 4X.
27379	Dogwood, Red Osier (Cornus sericea)	Field Container	Freiberger	NJ	2009	Over the top	Slight injury but some plant death at 0.97, 1.94 and 3.88 lb ai per acre.
21425	Dogwood, Red Osier (Cornus sericea)	Field In-Ground	Garrett	MD	2001		
27379	Dogwood, Red Osier (Cornus sericea)	Field Container	Peachey	OR	2013	Over the top	Slight injury and growth reduction with 21, moderate with 42 and 84 fl oz per acre, applied twice.
21425	Dogwood, Red Osier (Cornus sericea)	Field In-Ground	Persad	OH	2014	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27379	Dogwood, Red Osier (Cornus sericea)	Field Container	Persad	OH	2014	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27419	Cotoneaster (Cotoneaster dammeri)	Field Container	Neal	NC	2010	Over the top	No injury at 1x (0.97 lb ai per acre), injury with 2x only immediately following retreatment, significant injury with 4x.
27419	Cotoneaster (Cotoneaster dammeri) 'Royal Beauty'	Field Container	Beste/Frank (ARS)	MD	2011	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice; no effect on marketability.
27130	Cotoneaster (Cotoneaster sp.) C. dammeri 'Lowfast'	Field Container	Regan	WA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre after 1st, slight to high injury increasing with rates after 2nd application; significant growth reduction.
27130	Cotoneaster (Cotoneaster sp.) C. glaucophyllus	Field Container	Uber	CA	2008	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre; significant growth reduction at 4X

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27130	Cotoneaster (Cotoneaster sp.) C. horizontalis 'Perpusillus'	Field Container	Lieth	CA	2008	Over the top	Phytotoxicity data inconclusive, significant growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27130	Cotoneaster (Cotoneaster sp.) C. pameyii	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
31088	Stonecrop (Crassula sp.)	Field Container	Gilliam	AL	2015	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
31088	Stonecrop (Crassula sp.) C. ovata	Field Container	Villavicencio	CA	2012	Over the top	Slight to moderate reddish leaf color with 21, 42 and 84 fl oz per acre applied twice, slight necrosis at 1X and 2X, moderate at 4X; moderate growth reduction.
27135	Hawthorn (Crataegus sp.) C. coccinoid	Field Container	Freiberger	NJ	2009	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre.
27135	Hawthorn (Crataegus sp.) C. mollis	Field Container	Jones	OH	2013	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27140	Japanese Cedar (Cryptomeria japonica) 'Black Dragon'	Field Container	Neal	NC	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27140	Japanese Cedar (Cryptomeria japonica) C. japonica	Field Container	Mickelbart	IN	2009	Over the top	No significant differences in treated (21,42,84 fl oz/A) for height, width, % growth 12 WAT. The high rate did cause significant phytotoxicity rating at 6WAT.
27140	Japanese Cedar (Cryptomeria japonica) 'Yoshino'	Field Container	Czarnota	GA	2009	Over the top	No significant injury at 0.97, moderate at 1.94 and severe at 3.88 lb ai per acre
30565	Cypress, Leyland (Cupressocyparis leylandii)	Field Container	Beste	MD	2015	Over the top	No injury with 0.98 and 1.97 lb ai per acre, minor with 3.94 lb ai after 2nd applic with complete recovery; no growth reduction.
30565	Cypress, Leyland (Cupressocyparis leylandii)	Field Container	Witcher	TN	2016	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27382	Cypress (Cupressus sp.)	Field Container	Uber	CA	2011	Over the top	No crop injury or reduction in growth with 0.97, 1.94, 3.88 lb ai per acre.
27382	Cypress (Cupressus sp.) C. arizonica 'Carolina Sapphire'	Field Container	Czarnota	GA	2009	Over the top	No significant injury at 0.98 and 1.97, slight at 3.94 lb ai per acre
30833	Delosperma sp. (Delosperma sp.) 'Cooper's Ice'	Greenhouse	Wilen	CA	2012	Over the top	No injury with 21 and 32 fl oz per acre; poor liverwort control.
31089	Delosperma sp. (Delosperma sp.) D. congestum 'White Nugget'	Field Container	Klett	CO	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre after 1st, very slight with 4X after 2nd applic.; moderate growth reduction with 2X and 4X.
31089	Delosperma sp. (Delosperma sp.) D. cooperi	Field Container	Derr	VA	2013	Over the top	Slight injury but no flower count reduction at 0.98, 1.97 and 3.94 lb ai per acre applied twice.
30833	Delosperma sp. (Delosperma sp.) D. cooperi 'Fire Spinner'	Greenhouse	Derr	VA	2012	Over the top	Minor initial injury, with complete recovery, at 21 and 32 fl oz per 100 gal w/ or w/o irrigation. Acceptable liverwort control only with high rate w/o irrig.
31089	Delosperma sp. (Delosperma sp.) D. cooperi 'Purple Mountain'	Field Container	Boydston	WA	2014	Over the top	No significant injury or growth reduction with 0.98, slight and moderate with 1.97 and 3.94 lb ai per acre applied twice; 1X- and 2X-treated plants marketable.
31089	Delosperma sp. (Delosperma sp.) D. nubigenum 'Basutoland'	Field Container	Senesac	NY	2013	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
31089	Delosperma sp. (Delosperma sp.) 'Fire Spinner'	Field Container	Derr	VA	2013	Over the top	Very minor injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
30528	Fern, Autumn (Dryopteris erythrosora)	Field Container	Senesac	NY	2013	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
30528	Fern, Autumn (Dryopteris erythrosora) 'Brilliance'	Field Container	Neal	NC	2013	Over the top	Virtually no injury with 21, moderate with 42 and 84, fl oz per acre applied twice.
30529	Fern, Shaggy Shield (Dryopteris sp.)	Field Container	Neal	NC	2013	Over the top	Slight injury with 21 and 42, moderate with 84, fl oz per acre applied twice.
30529	Fern, Shaggy Shield (Dryopteris sp.)	Field Container	Peachey	OR	2013	Over the top	No injury with 21 and 42, moderate with 84 fl oz per acre, applied twice; no growth reduction.
31207	Echeveria sp. (Echeveria sp.) 'Blue 2'	Field Container	Wilén	CA	2016	Over the top	Moderate to severe injury with 21, 42 and 84 fl oz per acre applied twice.
31207	Echeveria sp. (Echeveria sp.) 'Imbricata'	Field Container	Villavicencio	CA	2012	Over the top	Severe necrosis and moderate discoloration with 21, 42 and 84 fl oz per acre applied twice; significant growth reduction.
31207	Echeveria sp. (Echeveria sp.) E. shaviana 'Black Prince'	Field Container	Senesac	NY	2016	Over the top	Minor injury with 0.98, moderate with 1.97 and 3.94 lb ai per acre applied twice.
29534	Purple Coneflower (Echinacea sp.)	Field Container	Mansue	NJ	2013	Over the top	Trial not conducted because plants did not develop correctly.
29534	Purple Coneflower (Echinacea sp.) E. purpurea alba	Field Container	Klett	CO	2013	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
29534	Purple Coneflower (Echinacea sp.) 'Magnus'	Field Container	Boydston	WA	2013	Over the top	No injury or growth reduction with 0.98 and 1.97 lb ai per acre applied twice; moderate with 3.94 lb; 4X plants not saleable.
31208	Echinocactus grusonii (Echinocactus grusonii)	Field Container	Uber	CA	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
31208	Echinocactus grusonii (Echinocactus grusonii)	Field Container	Villavicencio	CA	2012	Over the top	Very minor spotting with 21, 42 and 84 fl oz per acre applied twice; slight chlorosis and growth reduction only at 4X.
31208	Echinocactus grusonii (Echinocactus grusonii)	Field Container	Villavicencio	CA	2013	Over the top	No injury with 21 and 42, very minor chlorosis with 84, fl oz per acre applied twice; very slight growth reduction only at 4X.
28711	California fuschia (Epilobium canum)	Field Container	Klett	CO	2015	Over the top	plants failed to thrive
28711	California fuschia (Epilobium canum)	Field Container	Wilén	CA	2009	Over the top	Moderate to high injury increasing with rates (0.97, 1.94 and 3.88 lb ai per acre); root and shoot dry weights significantly reduced at all rates.
27145	Winged Burning Bush (Euonymus alatus) E. alatus 'Compactus'	Field Container	Boydston	WA	2008	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
28205	Wintercreeper (Euonymus fortunei) 'Coloratus'	Field Container	Boydston	WA	2014	Over the top	No significant injury or growth reduction with 0.98 and 1.97, slight with 3.94 lb ai per acre applied twice; all treated plants marketable.
28205	Wintercreeper (Euonymus fortunei) 'Emerald N Gold'	Field Container	DeFrancesco	OR	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice; moderate growth reduction at all rates.
28205	Wintercreeper (Euonymus fortunei) 'Ivory Jade'	Field Container	Beste	MD	2014	Over the top	No injury or growth reduction with 0.97, 1.91 and 3.84 lb ai per acre applied twice; no reduction in marketability.
30566	Spindle Tree (Euonymus japonicus)	Field Container	Peachey	OR	2013	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice.
30566	Spindle Tree (Euonymus japonicus)	Field Container	Uber	CA	2013	Over the top	No injury with 21 and 42, minor with 84, fl oz per acre applied twice; no growth reduction.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
30566	Spindle Tree (<i>Euonymus japonicus</i>) <i>E. japonica</i> 'Elata'	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27273	Beech (<i>Fagus</i> sp.)	Field Container	Freiberger	NJ	2012	Over the top	No sufficient data collected because plants never broke dormancy and died, or died before trial conclusion.
27273	Beech (<i>Fagus</i> sp.) <i>F. grandifolia</i>	Field Container	Peachey	OR	2013	Over the top	Phytotoxicity determination inconclusive; no growth reduction with 21, 42 and 84 fl oz per acre applied twice.
27273	Beech (<i>Fagus</i> sp.) <i>F. grandifolia</i>	Field Container	Persad	OH	2014	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27273	Beech (<i>Fagus</i> sp.) <i>F. sylvatica</i>	Field Container	DeFrancesco	OR	2014	Over the top	Injury results unreliable; moderate growth reduction with 21, 42 and 84 fl oz per 100 gal.
27273	Beech (<i>Fagus</i> sp.) <i>F. sylvatica</i>	Field Container	Jones	OH	2013	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28257	Golden Bells (<i>Forsythia</i> sp.)	Field Container	Freiberger	NJ	2012	Over the top	No injury after the 1st application with 21, 42 and 84 fl oz per acre; slight injury (leaf burn) with 1X and 2X, moderate with 4X after 2nd application.
28257	Golden Bells (<i>Forsythia</i> sp.) <i>F. × intermedia</i> 'Lynwood Gold'	Field Container	Gilliam	AL	2013	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice.
28257	Golden Bells (<i>Forsythia</i> sp.) <i>F. × intermedia</i> 'Golden Bell'	Field Container	Mickelbart	IN	2008	Over the top	No injury at 2.65, 5.3 and 10.6 lb ai per acre with single application 3 weeks after transplanting.
28257	Golden Bells (<i>Forsythia</i> sp.) 'Lynwood Gold'	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
28206	Witch Alder (<i>Fothergila gardenii</i>) 'Mount Airy'	Field Container	Senesac	NY	2016	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
21426	Ash, Green (<i>Fraxinus pennsylvanica</i>)	Field In-Ground	Garrett	MD	2001		
26258	Ash (<i>Fraxinus</i> sp.) <i>F. americana</i>	Field In-Ground	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
31090	Jasmine, Cape, Common Gardenia (<i>Gardenia</i> sp.) <i>G. jasmonoides</i> 'Frostproof'	Field Container	Witcher	TN	2016	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
31090	Jasmine, Cape, Common Gardenia (<i>Gardenia</i> sp.) <i>G. veitchii</i>	Field Container	Uber	CA	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
29535	Gaura (<i>Gaura lindheimeri</i>) 'Belleza White Evolution'	Field Container	Klett	CO	2014	Over the top	No injury or significant growth reduction with 21, 42 and 84 fl oz per acre applied twice.
29535	Gaura (<i>Gaura lindheimeri</i>) 'Crimson Butterflies'	Field Container	Senesac	NY	2014	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
29535	Gaura (<i>Gaura lindheimeri</i>) 'Whirling Butterflies'	Field Container	Boydston	WA	2014	Over the top	No significant injury with 0.98 and 1.97, moderate injury with good recovery at 3.94 lb ai per acre applied twice; slight growth reduction at all rates; all treated plants marketable.
27151	Honey Locust (<i>Gleditsia</i> sp.) <i>G. triacanthos</i> L. 'Inermis'	Field Container	Beste/Frank (ARS)	MD	2010	Over the top	No crop injury or reduction in growth with two applications at 1.97 and 3.94 lb ai per acre.
26230	Honey Locust (<i>Gleditsia</i> sp.) <i>G. triacanthos</i>	Field In-Ground	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27151	Honey Locust (<i>Gleditsia</i> sp.) <i>Gleditsia triacanthos</i>	Field Container	Mathers (OSU)	OH	2009	Over the top	Acceptable injury at 0.97, 1.94 and 3.88 lb ai per acre after 1st, no injury after 2nd application; no growth reduction
29168	Daylily (<i>Hemerocallis</i> sp.)	Field Container	Rivera	PR	2014	Over the top	Minor initial injury with quick recovery at 0.98, 1.97 and 3.94 lb ai per acre applied twice.
29168	Daylily (<i>Hemerocallis</i> sp.)	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
29168	Daylily (<i>Hemerocallis</i> sp.) 'Green Flutter'	Field Container	Senesac	NY	2010	Over the top	Slight crop injury with two applications at 0.97, 1.94, 3.88 recovering during latter half of evaluation period.
29168	Daylily (<i>Hemerocallis</i> sp.) 'H. Happy Returns'	Field Container	Derr	VA	2010	Over the top	No crop injury with 1.0, 2.0, 4.0 lb ai per acre 19 and 33 DAT and 4DAT2. Excellent control of spotted spurge with 2-4.0 lb ai per acre 19 and 33DAT.
29168	Daylily (<i>Hemerocallis</i> sp.) H. sp. 'Stella d'Oro'	Field Container	Mathers (OSU)	OH	2010	Over the top	Ohio: Little to no crop injury to 'Stella d'Oro' at 0.97, 1.94, and 3.88 lb ai per acre.
29168	Daylily (<i>Hemerocallis</i> sp.) H. sp. 'Strawberry Candy'	Field Container	Mathers (OSU)	MI	2010	Over the top	Lincoln: Minor stunting at higher rates (1.94, 3.88 lb ai per acre).
29168	Daylily (<i>Hemerocallis</i> sp.) H. 'Stella d'Oro'	Field Container	Boydston	WA	2011	Over the top	No crop injury or reduction in growth with two applications at 0.97, 1.94, 3.88 lb ai per acre.
30838	Daylily (<i>Hemerocallis</i> sp.) 'Happy Returns' & 'Stella de Oro'	Greenhouse	Ahrens/Mervosh	CT	2012	Over the top	Virtually no injury with 21, 42 and 84 fl oz per acre applied once.
29168	Daylily (<i>Hemerocallis</i> sp.) 'Stella de Oro'	Field Container	Boydston	WA	2010	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all treated plants saleable.
28743	Toyon, Christmas Berry (<i>Heteromeles arbutifolia</i>)	Field Container	Wilen	CA	2009	Over the top	Moderate injury at 0.97, 1.94 and 3.88 lb ai per acre; root and shoot dry weights not significantly reduced.
30671	Coral Bells, Alumroot (<i>Heuchera sanguinea</i>) 'Bressingham'	Field Container	Senesac	NY	2014	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
30671	Coral Bells, Alumroot (<i>Heuchera sanguinea</i>) 'Firefly'	Field Container	Boydston	WA	2014	Over the top	Slight injury with good recovery with 0.98 and 1.97, moderate injury with 3.94 lb ai per acre applied twice; 1X- and 2X-treated plants marketable.
30671	Coral Bells, Alumroot (<i>Heuchera sanguinea</i>) 'Pink Lipstick'	Field Container	Klett	CO	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre after 1st, very slight with 4X after 2nd applic.; moderate growth reduction with 4X.
30671	Coral Bells, Alumroot (<i>Heuchera sanguinea</i>) 'Ruby Bells'	Field Container	Beste	MD	2014	Over the top	No significant injury or growth reduction with 0.98, 1.96 and 3.89 lb ai per acre applied twice; no reduction in marketability.
26298	Rosemallow (<i>Hibiscus</i> sp.)	Field Container	Harvey	WA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
26298	Rosemallow (<i>Hibiscus</i> sp.)	Field Container	Rivera	PR	2014	Over the top	Minor initial injury with quick recovery at 0.98, 1.97 and 3.94 lb ai per acre applied twice.
26298	Rosemallow (<i>Hibiscus</i> sp.) H. moscheuto	Field Container	Senesac	NY	2008	Over the top	Slight injury at 0.98, moderate at 1.98 and 3.94 lb ai per acre
26298	Rosemallow (<i>Hibiscus</i> sp.) H. rosa-sinensis 'White Wing'	Field Container	Uber	CA	2009	Over the top	Slight injury at 0.97 and 1.94, unacceptable at 3.88 lb ai per acre.
26298	Rosemallow (<i>Hibiscus</i> sp.) H. syriacus 'Blue Satin'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury at 0.97 and 1.94, very slight at 3.88 lb ai per acre; no growth or flower number reduction, all plants marketable
26298	Rosemallow (<i>Hibiscus</i> sp.) H. syriacus 'Red Heart'	Field Container	Freiberger	NJ	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26298	Rosemallow (Hibiscus sp.) 'Rose Queen'	Field Container	Wade	SC	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre; moderate stunting at 2X and 4X to make plants unsaleable
30842	Hosta (Hosta sp.) 'Blue Hawaii'	Greenhouse	Derr	VA	2013	Over the top	Low injury with 21 and 32 fl oz per acre w/ or w/o irrigation. High liverwort control at both rates w/o irrig.
30672	Hosta (Hosta sp.) 'Fortunei Aureomarginata'	Field Container	Boydston	WA	2013	Over the top	No injury or growth reduction with 0.98, 1.97, and 3.94 lb ai per acre applied twice; all plants saleable.
30672	Hosta (Hosta sp.) H. fortunei 'Francee'	Field Container	Fraelich	GA	2013	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice; all plants marketable.
30672	Hosta (Hosta sp.) H. fortunei 'Francee'	Field Container	Mansue	NJ	2013	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice.
30842	Hosta (Hosta sp.) H. fortunei 'Gold Standard'	Greenhouse	Ahrens/Mervosh	CT	2012	Over the top	Slight injury with 21, 42 and 84 fl oz per acre applied once.
30672	Hosta (Hosta sp.) H. fortunei 'Gold Standard'	Field Container	Senesac	NY	2014	Over the top	Minor injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
30672	Hosta (Hosta sp.) 'Wide Brim'	Field Container	Reding	OH	2014	Over the top	No significant injury with 0.98 and 1.47, slight with 3.94, lb ai per acre applied twice; no significant growth reduction; all plants marketable.
26250	Hydrangea (Hydrangea sp.)	Field Container	Ahrens/Mervosh	CT	2011	Over the top	Little to no crop injury or reduction in growth with two sequential applications at 0.97, 1.94 or 3.88 lb ai per acre.
26250	Hydrangea (Hydrangea sp.)	Field Container	Mathers (OSU)	OH	2007		Moderate to significant yellowing and stunting increasing with rate.
26250	Hydrangea (Hydrangea sp.)	Field Container	Rivera	PR	2014	Over the top	Minor injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice; no growth reduction.
26250	Hydrangea (Hydrangea sp.) 'Angel Robe'	Field Container	Lieth	CA	2007	Over the top	Unacceptable injury at 0.97, 1.94 and 3.88 lb ai per acre.
30150	Hydrangea (Hydrangea sp.) H. arborescens 'Invincibelle amourenses'	Greenhouse	Mathers (OSU)	MI	2010	Over the top	Slight crop injury decreasing with time from 32 oz/A.
26250	Hydrangea (Hydrangea sp.) H. macrophylla 'Nikko Blue'	Field Container	Czarnota	GA	2007	Over the top	No significant injury at 0.97 and 1.94 lb ai per acre, significant at 3.88 lb
30150	Hydrangea (Hydrangea sp.) H. macrophylla 'Endless Summer'	Greenhouse	Ahrens/Mervosh	CT	2012	Over the top	Slight injury with 21 and 42, moderate with 84 fl oz per acre applied once.
26250	Hydrangea (Hydrangea sp.) H. macrophylla 'Endless Summer'	Field Container	Wade	SC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26250	Hydrangea (Hydrangea sp.) H. macrophylla 'Nikko Blue'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	No injury at 0.97 and 1.94, slight injury with complete recovery at 3.88 lb ai per acre; all plants marketable
26250	Hydrangea (Hydrangea sp.) H. macrophylla 'Nikko Blue'	Field Container	Fraelich	GA	2007	Over the top	Slight to moderate injury (chlorosis) at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
26250	Hydrangea (Hydrangea sp.) H. paniculata 'PeeGee'	Field Container	Mickelbart	IN	2008	Over the top	No injury at 2.65, 5.3 and 10.6 lb ai per acre with single application 3 weeks after transplanting.
26250	Hydrangea (Hydrangea sp.) H. quercifolia 'Alice'	Field Container	Senesac	NY	2007	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre
31091	Candytuft (Iberis sp.)	Field Container	Uber	CA	2013	Over the top	Moderate to severe injury increasing with rates (21, 42 and 84 fl oz per acre) applied twice.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
31091	Candytuft (Iberis sp.) 'Candytuft'	Field Container	Boydston	WA	2013	Over the top	No injury or growth reduction with 0.98, 1.97, and 3.94 lb ai per acre applied twice; all plants saleable.
31091	Candytuft (Iberis sp.) Iberis sempervirens	Field Container	Klett	CO	2013	Over the top	Slight injury and growth reduction with 21, severe with 42 and 84 fl oz per acre applied twice.
28204	Inkberry (Ilex glabra) 'Shamrock'	Field Container	Neal	NC	2015	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
28204	Inkberry (Ilex glabra) 'Shamrock'	Field Container	Persad	OH	2014	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
26331	Holly (Ilex sp.) 'Conaf'	Field Container	Czarnota	GA	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26331	Holly (Ilex sp.) I. cornuta 'Bufordii Nana'	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26331	Holly (Ilex sp.) I. cornuta 'Carissa'	Field Container	Fraelich	GA	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26331	Holly (Ilex sp.) I. cornuta 'Needlepoint'	Field Container	Neal	NC	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
26331	Holly (Ilex sp.) I. crenata 'Compacta'	Field Container	Wade	SC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26331	Holly (Ilex sp.) I. verticillata 'Winter Gold'	Field Container	Senesac	NY	2007	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre
30153	Holly, Blue (Ilex x meserveae) I. 'China Girl'	Greenhouse	Mathers (OSU)	MI	2010	Over the top	Moderate crop injury with 32 oz per acre.
31898	Flag (Iris sp.)	Field In-Ground	Senesac	NY	2014	Over the top	Trial not completed due to crop failure; will be repeated in 2015.
31898	Flag (Iris sp.) 'Blue Diamond'	Field In-Ground	Miller	WA	2015	Over the top	Moderate injury (less flowers, lower bulb weight) with 21, 42 and 84 fl oz per acre applied twice.
31898	Flag (Iris sp.) I. reticulata	Field In-Ground	Senesac	NY	2015	Over the top	No injury with 0.97 and 1.94, slight with 3.88 lb ai per acre applied twice.
28207	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Gilliam	AL	2014	Over the top	No foliar injury, but slight to severe growth reduction, with 21, 42 and 84 fl oz per acre applied twice; only 1X plants marketable.
28207	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Marble	FL	2016	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice; slight to moderate growth reduction; all plants marketable.
28207	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Persad	OH	2014	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
19628	Walnut, Black (Non-Bearing) (Juglans nigra)	Field In-Ground	Beste/Frank (ARS)	MD	2008	Over the top	Virtually no injury at 0.97, 1.94 and 3.88 lb ai per acre; growth reduction only at 4X; all plants marketable
27377	Walnut, Black (Non-Bearing) (Juglans nigra)	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	Data not reliable. No significant injury at 0.97 and 1.94, moderate at 3.88 lb ai per acre.
19628	Walnut, Black (Non-Bearing) (Juglans nigra)	Field In-Ground	Beste/Frank (ARS)	MD	2010	Over the top	No significant injury with 0.98, moderate initial injury with good recovery at 1.97 and 3.94 lb ai per acre applied twice; significant height reduction increasing with rates; plants marketable.
19628	Walnut, Black (Non-Bearing) (Juglans nigra)	Field In-Ground	Beste/Frank (ARS)	MD	2011	Over the top	No significant injury or growth reduction with 0.99, 1.97 and 3.94 lb ai per acre applied twice; plants marketable.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27377	Walnut, Black (Non-Bearing) (<i>Juglans nigra</i>)	Field Container	Freiberger	NJ	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
19628	Walnut, Black (Non-Bearing) (<i>Juglans nigra</i>)	Field In-Ground	Garrett	MD	2001		
27377	Walnut, Black (Non-Bearing) (<i>Juglans nigra</i>)	Field Container	Senesac	NY	2011	Over the top	No crop injury with two applications at 0.97, 1.94, and 3.88 lb aia.
26268	Juniper (<i>Juniperus</i> sp.)	Field Container	Harvey	WA	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26268	Juniper (<i>Juniperus</i> sp.)	Field Container	Mathers (OSU)	OH	2007	Over the top	No injury observed.
26268	Juniper (<i>Juniperus</i> sp.) <i>J. chinensis</i> 'Sea Green'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
26268	Juniper (<i>Juniperus</i> sp.) <i>J. chinensis</i> 'Sea Green'	Field Container	Mickelbart	IN	2008	Over the top	No injury at 2.65, 5.3 and 10.6 lb ai per acre with single application 3 weeks after transplanting.
26268	Juniper (<i>Juniperus</i> sp.) <i>J. communis</i> 'Gold Totem Pole'	Field Container	Senesac	NY	2007	Over the top	Virtually no injury at 0.97, 1.94 and 3.88 lb ai per acre
26268	Juniper (<i>Juniperus</i> sp.) <i>J. davunica</i> 'Parsonii'	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26268	Juniper (<i>Juniperus</i> sp.) <i>J. horizontalis</i> 'Youngstown'	Field Container	Regan	OR	2007	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26268	Juniper (<i>Juniperus</i> sp.) <i>J. squamata</i> 'Blue Star'	Field Container	Ahrens/Mervosh	CT	2007	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre
31092	Japanese Kerria, Japanese Rose (<i>Kerria japonica</i>) <i>K. japonica</i> 'Pleniflora'	Field Container	Senesac	NY	2013	Over the top	Minor injury with complete recovery with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
26339	Crape Myrtle (<i>Lagerstroemia indica</i>)	Field Container	Fraelich	GA	2007	Over the top	Slight injury at 0.97, moderate at 1.94 and 3.88 lb ai per acre
26339	Crape Myrtle (<i>Lagerstroemia indica</i>) 'Catawea'	Field Container	Wade	SC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26339	Crape Myrtle (<i>Lagerstroemia indica</i>) 'Dynamite'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	Slight injury after 1st application, with quick recovery at 0.97, 1.94 and 3.88 lb ai per acre; significant growth reduction at 4X but all plants marketable
26339	Crape Myrtle (<i>Lagerstroemia indica</i>) <i>L. x 'Muskogee'</i>	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
27312	Shrub Verbena (<i>Lantana</i> sp.) <i>L. camara</i> 'Dwarf Radiation'	Field Container	Lieth	CA	2009	Over the top	No significant crop injury with two applications at 0.97, 1.94, 3.88 lb aia during 10 week trial.
27312	Shrub Verbena (<i>Lantana</i> sp.) <i>L. montevidensis</i> 'Lynwood Gold'	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
27312	Shrub Verbena (<i>Lantana</i> sp.) 'New Gold'	Field Container	Czarnota	GA	2007	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per
27312	Shrub Verbena (<i>Lantana</i> sp.) 'New Gold'	Field Container	Gilliam	AL	2009	Over the top	No injury or growth reduction at 0.97, 1.84 and 3.88 lb ai per acre.
27169	Lavender (<i>Lavandula</i> sp.) <i>L. angustifolia</i> 'Hidcote'	Field Container	Boydston	WA	2009	Over the top	No injury at 0.97 and 1.94, moderate at 3.88 lb ai per acre; all treated plants saleable.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27169	Lavender (<i>Lavandula</i> sp.) <i>L. angustifolia</i> 'Munstead'	Field Container	Neal	NC	2009	Over the top	No significant injury at 0.97 and 1.94, moderate stunting at 3.88 lb ai per acre.
27169	Lavender (<i>Lavandula</i> sp.) <i>L. angustifolia</i> 'Munstead'	Field Container	Trader	MS	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27169	Lavender (<i>Lavandula</i> sp.) <i>L. intermedia</i> 'Provence'	Field Container	Boydston	WA	2008	Over the top	No injury at 0.97 and 1.94, slight at 3.88 lb ai per acre; no growth reduction; all plants marketable
27169	Lavender (<i>Lavandula</i> sp.) 'Munstead'	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
31093	Shasta Daisy (<i>Leucanthemum maximum</i>) 'Alaska'	Field Container	Derr	VA	2013	Over the top	Virtually no injury with 0.98 and 1.97, minor with 3.94, lb ai per acre applied twice.
31093	Shasta Daisy (<i>Leucanthemum maximum</i>) <i>L. x superbum</i> 'Becky'	Field Container	DeFrancesco	OR	2014	Over the top	No to slight injury with 21, 42 and 84 fl oz per acre applied twice; no growth reduction.
31093	Shasta Daisy (<i>Leucanthemum maximum</i>) <i>L. x superbum</i> 'Becky'	Field Container	Senesac	NY	2013	Over the top	Minor injury with complete recovery at 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27174	Fetterbush, Drooping <i>Leucothoe</i> (<i>Leucothoe</i> sp.) <i>L. (or Agarista) populifolia</i>	Field Container	Derr	VA	2011	Over the top	Little to no injury with 1, 2, 4 lb ai per acre. Excellent control of s. crabgrass, but no control of redroot pigweed, annual sedge, or doveweed with 1 lb ai per acre.
27174	Fetterbush, Drooping <i>Leucothoe</i> (<i>Leucothoe</i> sp.) <i>L. fontanesiana</i>	Field Container	Mickelbart	IN	2009	Over the top	There were no treatment effects on growth ratings with 21, 42, 84 fl. Oz./A. Phytotoxicity was observed after first and second application but by 6WAT only the high rate was significantly different from the untreated.
27174	Fetterbush, Drooping <i>Leucothoe</i> (<i>Leucothoe</i> sp.) <i>L. fontanesiana</i> 'Rainbow'	Field Container	Grunwald	OR	2010	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all treated plants saleable.
27383	Privet (<i>Ligustrum</i> sp.) <i>L. lucidum</i>	Field Container	Uber	CA	2008	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre; significant growth reduction at 4X
27383	Privet (<i>Ligustrum</i> sp.) <i>L. vulgare</i>	Field Container	Freiberger	NJ	2008	Over the top	Slight injury at 0.97 and 1.94, moderate at 3.88 lb ai per acre
27383	Privet (<i>Ligustrum</i> sp.) <i>L. x vicaryi</i>	Field Container	Senesac	NY	2009	Over the top	No injury at 0.97, very slight at 1.9 and 3.9 lb ai per acre after 2nd application.
32213	Honeysuckle (<i>Lonicera</i> sp.) <i>L. nitida</i> 'Lemon Beauty'	Field Container	DeFrancesco	OR	2014	Foliar	No injury with 21 and 42 fl oz per 100 gal applied twice, slight with 84 fl oz at end of trial; moderate to severe growth reduction.
32213	Honeysuckle (<i>Lonicera</i> sp.) <i>L. sempervirens</i> 'Alabama'	Field Container	Senesac	NY	2016	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
26368	<i>Loropetalum</i> (<i>Loropetalum</i> sp.)	Field Container	Derr	VA	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre.
26368	<i>Loropetalum</i> (<i>Loropetalum</i> sp.) <i>L. chinense rubrum</i>	Field Container	Czarnota	GA	2007	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre
26368	<i>Loropetalum</i> (<i>Loropetalum</i> sp.) <i>L. chinense rubrum</i>	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26368	<i>Loropetalum</i> (<i>Loropetalum</i> sp.) <i>L. chinensis</i> 'Ruby'	Field Container	Neal	NC	2009	Over the top	No significant injury at 0.97, slight to moderate stunting at 1.94 and 3.88 lb ai per acre.
26312	<i>Magnolia</i> (<i>Magnolia</i> sp.) <i>M. grandiflora</i> 'Alta'	Field Container	Wade	SC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26312	Magnolia (Magnolia sp.) M. stellata 'Centennial'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
26312	Magnolia (Magnolia sp.) M. virginiana	Field Container	Fraelich	GA	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
27185	Apple & Crabapple (Non-Bearing) (Malus sp.) M. domestica	Field Container	DeFrancesco	OR	2011	Over the top	No injury after 1st application with 21, very slight, with complete recovery, with 42 and 84 fl oz per acre; no injury after 2nd application; no growth reduction.
27185	Apple & Crabapple (Non-Bearing) (Malus sp.) 'M.domestica'	Field Container	Mathers (OSU)	OH	2009	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre.
27185	Apple & Crabapple (Non-Bearing) (Malus sp.) 'Prairie Fire'	Field Container	Jones	OH	2013	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27185	Apple & Crabapple (Non-Bearing) (Malus sp.) 'Spring Snow'	Field Container	Grunwald	OR	2010	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all treated plants saleable
31086	Mammillaria sp. (Mammillaria sp.) Mammillaria melanocentra	Field Container	Uber	CA	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
28212	Muhly, hairyawn (Muhlenbergia capillaris)	Field Container	Gilliam	AL	2014	Over the top	No to slight injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice; all plants marketable.
28212	Muhly, hairyawn (Muhlenbergia capillaris)	Field Container	Klett	CO	2015	Over the top	Tower not included in this trial.
28212	Muhly, hairyawn (Muhlenbergia capillaris)	Field Container	Senesac	NY	2014	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28715	Mexican deergrass (Muhlenbergia dubia)	Field Container	Klett	CO	2013	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
28715	Mexican deergrass (Muhlenbergia dubia)	Field Container	Senesac	NY	2014	Over the top	Very slight injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28715	Mexican deergrass (Muhlenbergia dubia)	Field Container	Wilén	CA	2009	Over the top	No significant injury at 0.97, moderate at 1.94 and 3.88 lb ai per acre; significant growth reduction at 4X.
28200	Bayberry (Myrica pensylvanica)	Field Container	Freiberger	NJ	2012	Over the top	No sufficient data collected because plants never broke dormancy and died, or died before trial conclusion.
28200	Bayberry (Myrica pensylvanica)	Field Container	Senesac	NY	2016	Over the top	No injury with 0.97, 1.94 and 3.88 lb ai per acre applied twice.
26324	Heavenly Bamboo (Nandina domestica)	Field Container	Gilliam	AL	2009	Over the top	No injury or growth reduction at 0.97, 1.84 and 3.88 lb ai per acre.
26324	Heavenly Bamboo (Nandina domestica) 'Fire Power'	Field Container	Gilliam	AL	2008	Over the top	No injury at 0.97, 1.84 and 3.88 lb ai per acre with two applications.
26324	Heavenly Bamboo (Nandina domestica) 'Firepower'	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97 and 1.94, slight at 3.88 lb ai per acre.
26324	Heavenly Bamboo (Nandina domestica) N. domestica	Field Container	Lieth	CA	2009	Over the top	Significant crop injury with 0.97, 1.94, and 3.88 lb ai compared to untreated increasing with time and second application (20-40% by 60DAT). Dramatic reduction in height compared to untreated.
31899	Daffodil (Narcissus sp.)	Field In-Ground	Senesac	NY	2014	Over the top	Trial not completed due to crop failure; will be repeated in 2015.
31899	Daffodil (Narcissus sp.) 'Flower Carpet'	Field In-Ground	Miller	WA	2015	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
31899	Daffodil (Narcissus sp.) 'Golden Echo'	Field In-Ground	Senesac	NY	2015	Over the top	Slight injury with complete recovery at 0.97 and 1.94, moderate to slight with 3.88 lb ai per acre applied twice.
27193	Catnip (Nepeta cataria) N. cataria	Field Container	Lieth	CA	2009	Over the top	Significant crop injury (10-25%) with 0.97, 1.94, and 3.88 lb aia compared to untreated 2WAT. (Crop injury ratings after 2WAT inconclusive due to damage on all plants.) Plant volume decreased by half with 1.94 and 3.88 lb aia rates 60DAT.
27193	Catnip (Nepeta cataria) 'Psfike'	Field Container	Klett	CO	2008	Over the top	Trial 1: Some injury (chlorosis) at 0.97, 1.94 and 3.88 lb ai per acre; growth reduction at 4X
27193	Catnip (Nepeta cataria) 'Psfike'	Field Container	Klett	CO	2008	Over the top	Trial 2: No injury at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
27198	Catmint (Nepeta x faasseni)	Field Container	Lieth	CA	2008	Over the top	Unacceptable injury at 0.97, 1.94 and 3.88 lb ai per acre
27198	Catmint (Nepeta x faasseni) 'Walker's Low'	Field Container	Boydston	WA	2012	Over the top	No injury or growth reduction with 0.97, 1.94 and 3.88 lb ai per acre applied twice; all plants saleable.
27198	Catmint (Nepeta x faasseni) 'Walker's Low'	Field Container	Senesac	NY	2008	Over the top	Slight injury at 0.98 and 1.98, moderate at 3.94 lb ai per acre
27198	Catmint (Nepeta x faasseni) 'Walker's Low'	Field Container	Wilén	CA	2012	Over the top	Minor, acceptable injury with 21, 42 and 84 fl oz per acre after 1st applic; unacceptable injury and growth reduction after 2nd applic.
29245	Mondo Grass, Lilyturf, Ker-Gawl (Ophiopogon sp.) O. japonica	Field Container	Uber	CA	2012	Over the top	No injury with 0.97, 1.94 and 3.88 lb ai per acre applied twice; very slight growth reduction at 2X and 4X.
29245	Mondo Grass, Lilyturf, Ker-Gawl (Ophiopogon sp.) O. japonicas	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
29245	Mondo Grass, Lilyturf, Ker-Gawl (Ophiopogon sp.) O. japonicas	Field Container	Czarnota	GA	2012	Over the top	No injury with 21 and 42 fl oz per acre, slight with complete recovery with 84 fl oz, applied twice; no height reduction.
29245	Mondo Grass, Lilyturf, Ker-Gawl (Ophiopogon sp.) O. japonicas	Field Container	Gilliam	AL	2013	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice.
30846	Mondo Grass, Lilyturf, Ker-Gawl (Ophiopogon sp.) O. japonicus 'Nana'	Greenhouse	Wilén	CA	2013	Over the top	Excellent crop safety with 21 and 32 fl oz per acre; excellent liverwort control by 44 DAT.
29253	Holly Olive; False (Osmanthus heterophyllus) Holly Olive; false	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
29253	Holly Olive; False (Osmanthus heterophyllus) O. fragrans	Field Container	Gilliam	AL	2013	Over the top	Moderate injury after 1st application with 21, 42 and 84 fl oz per acre, with good recovery to slight injury after 2nd applic; no growth reduction; treated plants marketable.
31877	Fern, Royal (Osmunda regalis)	Greenhouse	Derr	VA	2013	Over the top	Low injury with 21 and 32 fl oz per acre w/ or w/o irrigation. High liverwort control at both rates w/o irrig.
31956	Fern, Royal (Osmunda regalis)	Field Container	Neal	NC	2013	Over the top	Slight injury with 21, moderate with 42 and 84, fl oz per acre applied twice.
29084	Peony (Paeonia sp.) P. lactiflora 'Karl Rosenfield'	Field Container	Boydston	WA	2011	Over the top	No crop injury or reduction in growth with two applications at 0.97, 1.94, and 3.88 lb ai per acre.
29084	Peony (Paeonia sp.) 'President Roosevelt'	Field In-Ground	Miller	WA	2015	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
29084	Peony (Paeonia sp.) P. lactiflora 'Edulis Superba'	Field Container	Senesac	NY	2011	Over the top	No injury with 0.97, 1.94, and 3.88 lb ai per acre applied twice.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
31094	Beard-Tongue (Penstemon sp.) P. barbatus 'Navigator Mix'	Field Container	Reding	OH	2014	Over the top	No significant injury or growth reduction with 0.98 and 1.47, moderate with good recovery with 3.94, lb ai per acre applied twice; all plants marketable.
31094	Beard-Tongue (Penstemon sp.) P. digitalis 'Huskies Red'	Field Container	Boydston	WA	2014	Over the top	Slight to moderate injury (leaf burn) with good recovery at 0.98, 1.97 and 3.94 lb ai per acre applied twice; no significant growth reduction; 1X- and 2X-, and most 4X-treated plants marketable.
31094	Beard-Tongue (Penstemon sp.) P. pinifolius 'Mersea Yellow'	Field Container	DeFrancesco	OR	2015	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
28739	Mock Orange (Philadelphus sp.)	Field Container	Freiberger	NJ	2012	Over the top	Inconclusive results because plants died off early in both treated and control plants.
28739	Mock Orange (Philadelphus sp.) P. viginialis 'Snow Dwarf'	Field Container	Czarnota	GA	2009	Over the top	No significant injury at 0.97 and 1.94, slight at 3.88 lb ai per acre.
28475	Photinia (Photinia sp.)	Field Container	Czarnota	GA	2013	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
28475	Photinia (Photinia sp.) P. fraseri	Field Container	Uber	CA	2008	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre after 1st, moderate at 2X and 4X after 2nd application; significant growth reduction at 4X
28475	Photinia (Photinia sp.) P. fraserii	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
28475	Photinia (Photinia sp.) P. villosa	Field Container	Beste	MD	2014	Over the top	Acceptable injury with 1, 2 and 4 lb ai per acre applied twice, but moderate growth reduction with 2X and 4X; marketability reduction at 2X and 4X.
19626	Spruce, Norway (Picea abies)	Field In-Ground	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
19626	Spruce, Norway (Picea abies)	Field In-Ground	Beste/Frank (ARS)	MD	2008	Over the top	Injury moderate though not significant at trial end at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction; reduced marketability at 2X and 4X
29511	Spruce, Colorado (Picea pungens)	Field In-Ground	Beste/Frank (ARS)	MD	2011	Over the top	No significant injury, growth or marketability reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
26291	Spruce (Picea sp.) P. abies	Field Container	Mathers (OSU)	OH	2008	Over the top	No injury to very slight chlorosis increasing with rate (0.97, 1.94, 3.88 lb ai per acre).
26291	Spruce (Picea sp.) P. abies	Field Container	Regan	WA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre; slight growth reduction at 4X.
26291	Spruce (Picea sp.) P. glauca	Field Container	Freiberger	NJ	2007	Over the top	No injury at 0.97, 1.94 and 3.99 lb ai per acre
26291	Spruce (Picea sp.) P. glauca	Field Container	Freiberger	NJ	2012	Over the top	No sufficient data collected because plants never broke dormancy and died, or died before trial conclusion.
26291	Spruce (Picea sp.) P. glauca	Field Container	Mickelbart	MI	2009	Over the top	Phytotoxicity was zero at all observation times with no treatment effects on growth involving 21, 42, 84 fl oz./A 12 WAT.
26291	Spruce (Picea sp.) P. pungens	Field Container	Senesac	NY	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26291	Spruce (Picea sp.) P. pungens	Field Container	Senesac	NY	2008	Over the top	No significant injury at 0.98, slight at 1.98, moderate at 3.94 lb ai per acre

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26291	Spruce (Picea sp.) P. pungens glauca	Field Container	Klett	CO	2008	Over the top	Trial 1: No injury at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
26291	Spruce (Picea sp.) P. pungens glauca	Field Container	Klett	CO	2008	Over the top	Trial 2: No injury at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
27209	Andromeda (Pieris sp.) P. japonica 'Dorothy Wycoff'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury at 0.97 and 1.94, slight at 3.88 lb ai per acre; no growth reduction, all plants marketable
27209	Andromeda (Pieris sp.) P. japonica 'Mountain Fire'	Field Container	Trader	MS	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27209	Andromeda (Pieris sp.) P. japonica 'Dorothy Wycoff'	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable.
27214	Pine (Pinus sp.)	Field Container	Freiberger	NJ	2012	Over the top	No sufficient data collected because plants never broke dormancy and died, or died before trial conclusion.
27214	Pine (Pinus sp.) P. eldarica	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27214	Pine (Pinus sp.) P. halapensis	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27214	Pine (Pinus sp.) P. mugo	Field Container	Harvey	WA	2008	Over the top	No injury at 0.97, minor injury at 1.94 3.88 lb ai per acre after 2nd application that was gone 2 weeks later
27214	Pine (Pinus sp.) P. strobus	Field Container	Trader	MS	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27214	Pine (Pinus sp.) P. taeda	Field Container	Fraelich	GA	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable.
27214	Pine (Pinus sp.) Pine Mungo	Field Container	Harvey	WA	2009	Over the top	No injury at 0.97 and 1.94, slight at 3.88 lb ai per acre.
27378	Pine, Loblolly (Pinus taeda)	Field Container	Fraelich	GA	2015	Over the top	No injury or difference in plant growth and marketability with 21, 42 and 84 fl oz per acre applied twice.
27378	Pine, Loblolly (Pinus taeda)	Field Container	Neal	NC	2015	Over the top	No injury with 21 and 42 fl oz per acre applied twice, very slight with 84 fl oz.
27219	Cinquefoil (Potentilla sp.) P. fruticosa 'Abbotswood'	Field Container	Lieth	CA	2009	Over the top	Slight(15-20%) but statistically significant crop injury with 0.97, 1.94, and 3.88 lb aia compared to untreated 6WAT. Lack of significant plant growth treatment effect suggest Tower is considered safe for application to Potentilla.
27219	Cinquefoil (Potentilla sp.) P. fruticosa 'Goldfinger'	Field Container	Uber	CA	2008	Over the top	No significant injury at 0.97 and 1.94, significant at 3.88 lb ai per acre; no significant growth reduction
27219	Cinquefoil (Potentilla sp.) P. fruticosa 'Monsidh'	Field Container	Klett	CO	2008	Over the top	Trial 1: No injury at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
27219	Cinquefoil (Potentilla sp.) P. fruticosa 'Monsidh'	Field Container	Klett	CO	2008	Over the top	Trial 2: No injury at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
27219	Cinquefoil (Potentilla sp.) P. neumaniana 'Verna'	Field Container	Boydston	WA	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants saleable.
26222	Fir, Douglas (Pseudotsuga menziesii)	Field Container	Boydston	WA	2007	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26222	Fir, Douglas (Pseudotsuga menziesii)	Field Container	Harvey	WA	2007	Over the top	Unreliable results due to transplant shock

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26222	Fir, Douglas (Pseudotsuga menziesii) 'Blue'	Field Container	Marshall	MI	2007	Over the top	One application. No injury at 0.97, 1.94 and 3.88 lb ai per acre
26222	Fir, Douglas (Pseudotsuga menziesii) P. menziesii glauca	Field Container	Freiberger	NJ	2007	Over the top	No injury at 0.97, 1.94 and 3.99 lb ai per acre after 1st, significant at 1X after 2nd application
27797	Oak, Sawtooth (Quercus acutissima)	Field In-Ground	Beste/Frank (ARS)	MD	2008	Over the top	Slight injury with complete recovery at trial end at 0.97, 1.94 and 3.88 lb ai per acre; height reduction at 4X but all plants marketable.
27797	Oak, Sawtooth (Quercus acutissima)	Field In-Ground	Beste/Frank (ARS)	MD	2011	Over the top	Slight to moderate injury with 0.98, 1.97 and 3.94 lb ai per acre after 1st application, complete recovery in 3 months; reduced height at 1X and 4X but no reduced market value.
27797	Oak, Sawtooth (Quercus acutissima)	Field In-Ground	Beste/Frank (NER)	MD	2013	Over the top	No injury with 21 and 42, moderate injury with quick and complete recovery at 84 fl oz per acre, applied twice; slight growth reduction at 2X and 4X; no reduction in marketability.
27384	Oak, Sawtooth (Quercus acutissima)	Field Container	Gilliam	AL	2013	Over the top	Slight injury with complete recovery after 1st, no injury after 2nd, application with 21, 42 and 84 fl oz per acre applied twice; no growth reduction.
27421	Nuttall oak (Quercus nuttallii)	Field Container	Gilliam	AL	2013	Over the top	Acceptable injury after 1st, no injury after 2nd, application with 21, 42 and 84 fl oz per acre applied twice; no growth reduction.
19627	Oak, Northern Red (Quercus rubra)	Field In-Ground	Garrett	MD	2001		
27225	Oak (Quercus sp.) Q. alba	Field Container	Freiberger	NJ	2008	Over the top	Slight injury at 0.97 and 1.94, moderate at 3.88 lb ai per acre.
27225	Oak (Quercus sp.) Q. pinnata	Field Container	Mathers (OSU)	OH	2011	Over the top	Significant crop injury (leaf burn) and growth reduction with two sequential applications at 0.97, 1.94, 3.88 lb ai per acre.
27225	Oak (Quercus sp.) Q. rubra	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable.
27225	Oak (Quercus sp.) Q. rubra	Field Container	Mathers (OSU)	OH	2008	Over the top	Severe injury at 0.97, 1.94 and 3.88 lb ai per acre.
27230	Indian Hawthorn (Raphiolepis indica)	Field Container	Gilliam	AL	2008	Over the top	No injury at 0.97, 1.84 and 3.88 lb ai per acre with two applications.
27230	Indian Hawthorn (Raphiolepis indica) 'Indian Princess'	Field Container	Lieth	CA	2008	Over the top	Slight, acceptable injury at 4X but unacceptable growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
27230	Indian Hawthorn (Raphiolepis indica) 'Pink Lady'	Field Container	Uber	CA	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27230	Indian Hawthorn (Raphiolepis indica) R. umbellata 'Eleanor Taber'	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26287	Azalea (Rhododendron sp.) 'Amelia Rose'	Field Container	Gilliam	AL	2007	Over the top	Slight injury after 1st but none after 2nd application only at 3.88 lb ai per acre (4X); no growth reduction
30847	Azalea (Rhododendron sp.) 'Delaware Valley White'	Greenhouse	Ahrens/Mervosh	CT	2012	Over the top	Very slight injury with 21, 42 and 84 fl oz per acre applied once.
26287	Azalea (Rhododendron sp.) 'Fashion'	Field Container	Fraelich	GA	2007	Over the top	Slight injury (chlorosis) at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
26287	Azalea (Rhododendron sp.) 'Fashion'	Field Container	Neal	NC	2009	Over the top	No significant injury at 0.97 and 1.94, slight stunting at 3.88 lb ai per acre.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26287	Azalea (Rhododendron sp.) 'Gwenda'	Field Container	Wade	SC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26287	Azalea (Rhododendron sp.) R. x 'Crete' (Yakushmanum hybrid)	Field Container	Senesac	NY	2007	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre
26287	Azalea (Rhododendron sp.) 'Vulcan'	Field Container	Regan	OR	2007	Over the top	Moderate injury after the 2nd application and growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
28719	Evergreen currant (Ribes viburnifolium)	Field Container	Wilén	CA	2009	Over the top	No significant injury at 0.97, moderate at 1.94 and 3.88 lb ai per acre; root and shoot dry weights not significantly reduced.
26198	Rose (Rosa sp.) 'Nearly Wild'	Field Container	Czarnota	GA	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26198	Rose (Rosa sp.) 'Flower Carpet Red'	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26198	Rose (Rosa sp.) 'Nearly Wild'	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26198	Rose (Rosa sp.) 'Nearly Wild'	Field Container	Lieth	CA	2007	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26198	Rose (Rosa sp.) 'Nearly Wild'	Field Container	Wade	SC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26198	Rose (Rosa sp.) R. virginiana	Field Container	Senesac	NY	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre after 1st application; slight and moderate injury at 2X and 4X after 2nd application with complete recovery in 4 weeks
30673	Coneflower (Rudbeckia sp.) 'Goldsturm'	Field Container	Boydston	WA	2013	Over the top	Virtually no injury and no growth reduction with 0.98 and 1.97 lb ai per acre applied twice, moderate with 3.94 lb; 4X plants not saleable.
30673	Coneflower (Rudbeckia sp.) R. hirta	Field Container	Klett	CO	2013	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
27821	Palmetto Palm (Sabal minor)	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27821	Palmetto Palm (Sabal minor)	Field Container	Marble	FL	2015	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27821	Palmetto Palm (Sabal minor)	Field Container	Witcher	TN	2016	Over the top	No injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
30676	Pasture Sage (Salvia nemorosa) 'May Night'	Field Container	Beste/Frank (ARS)	MD	2012	Over the top	Slight injury, significant flower reduction with 0.98 and 1.97 and 3.94 lb ai per acre; reduction in marketability.
30676	Pasture Sage (Salvia nemorosa) 'May Night'	Field Container	Boydston	WA	2014	Over the top	No significant injury with 0.98, slight and moderate with some recovery at 1.97 and 3.94 lb ai per acre applied twice; slight growth reduction at 4X; all treated plants marketable.
30676	Pasture Sage (Salvia nemorosa) 'May Night'	Field Container	Reding	OH	2014	Over the top	Moderate to severe injury and growth reduction with 0.98, 1.47 and 3.94 lb ai per acre applied twice; treated plants not marketable.
30676	Pasture Sage (Salvia nemorosa) S. sylvestris 'Blue Hill'	Field Container	Senesac	NY	2013	Over the top	No to minor injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
30676	Pasture Sage (Salvia nemorosa) 'Sensation Deep Rose'	Field Container	Klett	CO	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre after 1st, very slight with 4X after 2nd applic.; slight and moderate growth reduction with 2X and 4X.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26201	Sage, Ramona (<i>Salvia sylvestris</i>) 'May Night'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury and growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26201	Sage, Ramona (<i>Salvia sylvestris</i>) 'May Night'	Field Container	Klett	CO	2007	Over the top	Two trials; slight injury (leaf discoloration) at 0.97, 1.94 and 3.88 lb ai per acre; no growth reduction
26201	Sage, Ramona (<i>Salvia sylvestris</i>) <i>S. nemorosa</i> 'Cardonna'	Field Container	Derr	VA	2007	Over the top	8, 14, and 53 % injury at 1, 2 and 4 lb ai per acre; 100 % control of rice flatsedge and fragrant flatsedge
26201	Sage, Ramona (<i>Salvia sylvestris</i>) <i>S. nemorosa</i> 'May Night'	Field Container	Mathers (OSU)	OH	2007	Over the top	Slight injury (burning and yellowing) at 0.97, 1.94 and 3.92 lb ai per acre after 1st application but plants quickly recovered; slight growth reduction at 4X
26201	Sage, Ramona (<i>Salvia sylvestris</i>) 'Snow Hill'	Field Container	Boydston	WA	2007	Over the top	No injury but slight growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; plants saleable
29977	Beach Naupaka (<i>Scaevola</i> sp.) 'Mauve Clusters'	Field Container	Wilén	CA	2012	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
29977	Beach Naupaka (<i>Scaevola</i> sp.) <i>S. aemula</i> 'Scala Blue'	Field Container	Pemberton	TX	2012	Over the top	No injury or growth reduction with 0.97, 1.94 and 3.88 lb ai per acre applied twice; delayed flowering up to 4 weeks.
29977	Beach Naupaka (<i>Scaevola</i> sp.) <i>S. 'Cajun Blue'</i>	Field Container	Derr	VA	2010	Over the top	Little to no crop injury with 1.0 lb ai per acre. 62 DAT and 14DAT2. Fair to very good control of longstalked phyllanthus, tasselflower, s. crabgrass.
30674	Stonecrop (<i>Sedum</i> sp.)	Field Container	Uber	CA	2013	Over the top	Moderate to severe injury increasing with rates (21, 42 and 84 fl oz per acre) applied twice.
30674	Stonecrop (<i>Sedum</i> sp.) <i>S. reflexum</i> 'Blue Spruce'	Field Container	Reding	OH	2014	Over the top	No significant injury or growth reduction with 0.98, moderate and severe with 1.47 and 3.94, lb ai per acre applied twice; 2X- and 4X-treated plants not marketable.
30674	Stonecrop (<i>Sedum</i> sp.) <i>S. ternatum</i> 'Larinem Park'	Field Container	Senesac	NY	2013	Over the top	No injury with 0.98 and 1.97, minor injury with complete recovery at 3.94, lb ai per acre applied twice.
26279	Bridal-Wreath (<i>Spiraea</i> sp.)	Field Container	Harvey	WA	2007	Over the top	No injury after the first application at 0.97, 1.94 and 3.88 lb ai per acre, significant injury only at 4X rate after the second application
26279	Bridal-Wreath (<i>Spiraea</i> sp.) <i>S. decumbens</i>	Field Container	Senesac	NY	2007	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre
26279	Bridal-Wreath (<i>Spiraea</i> sp.) <i>S. prunifolia</i>	Field Container	Freiberger	NJ	2007	Over the top	No injury at 0.97, 1.94 and 3.99 lb ai per acre after 1st, moderate after 2nd application
26279	Bridal-Wreath (<i>Spiraea</i> sp.) <i>S. thunbergii</i>	Field Container	Mickelbart	IN	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre with single application 3 weeks after transplanting.
26279	Bridal-Wreath (<i>Spiraea</i> sp.) <i>Spiraea x bumalda</i> 'Gold Mound'	Field Container	Ahrens/Mervosh	CT	2007	Over the top	Very slight injury at 0.97, 1.94 and 3.88 lb ai per acre
27280	Camellia, Mountain (<i>Stewartia</i> sp.) <i>S. pseudocamellia</i>	Field Container	Senesac	NY	2011	Over the top	No crop injury with two applications at 0.97, 1.94, and 3.88 lb aia.
27248	Lilac (<i>Syringa</i> sp.) 'Miss Kim'	Field Container	Harvey	WA	2008	Over the top	No injury at 0.97 and 1.94, minor injury at 3.88 lb ai per acre
27248	Lilac (<i>Syringa</i> sp.) <i>S x tribida</i> 'Lark Song'	Field Container	Mathers (OSU)	OH	2008	Over the top	No significant injury or growth reduction at 0.97 and 1.94, moderate injury at 3.88 lb ai per acre
27248	Lilac (<i>Syringa</i> sp.) <i>S. microphylla</i> 'Superba'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	Significant injury (stem and leaf distortion), no growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; 1X marginally marketable, higher rates unacceptable

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
30149	Lilac (<i>Syringa</i> sp.) S. 'Paliban'	Greenhouse	Mathers (OSU)	MI	2010	Over the top	Moderate crop injury with 32 oz/A. No ratings taken first two weeks due to dormancy.
28223	Bald Cypress (<i>Taxodium distichum</i>)	Field Container	Neal	NC	2015	Over the top	No injury after 1st applic. with 21, 42 and 84 fl oz per acre, slight with good recovery after 2nd applic.
26308	Yew (<i>Taxus</i> sp.)	Field Container	Harvey	WA	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26308	Yew (<i>Taxus</i> sp.) T. <i>baccata</i>	Field Container	Senesac	NY	2007	Over the top	Slight to moderate injury at 0.97, 1.94 and 3.88 lb ai per acre with complete recovery by 4 weeks after 2nd application
26308	Yew (<i>Taxus</i> sp.) T. <i>baccata</i> <i>fasigata</i>	Field Container	Freiberger	NJ	2007	Over the top	No injury at 0.97, 1.94 and 3.99 lb ai per acre
26308	Yew (<i>Taxus</i> sp.) T. <i>baccata</i> 'Repandens'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants marketable
27256	Ternstroemia (<i>Ternstroemia</i> sp.)	Field Container	Neal	NC	2010	Over the top	No crop injury with any rate during the evaluation period.
27256	Ternstroemia (<i>Ternstroemia</i> sp.) 'Leann'	Field Container	Gilliam	AL	2009	Over the top	No injury or growth reduction at 0.97, 1.84 and 3.88 lb ai per acre.
27256	Ternstroemia (<i>Ternstroemia</i> sp.) T. <i>gymnanthera</i>	Field Container	Gilliam	AL	2011	Over the top	No crop injury with 0.97, 1.94, 3.88 lb ai per acre but slight reduction in growth with 1x and 4x.
28723	Dwarf germander (<i>Teucrium chamaedrys</i>)	Field Container	Klett	CO	2013	Over the top	Severe injury and growth reduction with 21, 42 and 84 fl oz per acre applied twice.
28723	Dwarf germander (<i>Teucrium chamaedrys</i>)	Field Container	Peachey	OR	2013	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice.
28723	Dwarf germander (<i>Teucrium chamaedrys</i>)	Field Container	Wilen	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre; root dry weight significantly reduced at all rates.
26226	Cedar, Western Red (<i>Thuja plicata</i>)	Field Container	Boydston	WA	2007	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26226	Cedar, Western Red (<i>Thuja plicata</i>)	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26226	Cedar, Western Red (<i>Thuja plicata</i>) 'Stoneham Gold'	Field Container	Senesac	NY	2008	Over the top	Slight injury at 0.98 and 1.98, moderate at 3.94 lb ai per acre
26226	Cedar, Western Red (<i>Thuja plicata</i>) T. <i>plicata</i> 'Watnong Green'	Field Container	Ahrens/Mervosh	CT	2008	Over the top	Slight injury at 0.98 and 1.97, marginal at 3.94 lb ai per acre after 2nd application; 1st application rates were 3.8 times higher than protocol rates.
26275	Arborvitae (<i>Thuja</i> sp.)	Field Container	Mathers (OSU)	OH	2007	Over the top	No to slight injury at highest rate.
30154	Arborvitae (<i>Thuja</i> sp.) T. <i>occidentalis</i> 'Nigra'	Greenhouse	Mathers (OSU)	MI	2010	Over the top	Minor crop injury initially 1WAT decreasing with time until second application causing slight to minor (13-23%) crop injury with 32 oz per acre.
26275	Arborvitae (<i>Thuja</i> sp.) T. <i>occidentalis</i> 'Emerald Green'	Field Container	Ahrens/Mervosh	CT	2007	Over the top	No injury after the 1st application at 0.97, 1.94 and 3.88 lb ai per acre; slight injury after 2nd application at the higher rates
26275	Arborvitae (<i>Thuja</i> sp.) T. <i>occidentalis</i> 'Emerald Green'	Field Container	Lieth	CA	2007	Over the top	Unacceptable injury (burning of shoot tips) and distorted growth at 0.97, 1.94 and 3.88 lb ai per acre
26275	Arborvitae (<i>Thuja</i> sp.) T. <i>occidentalis</i> 'Nigra'	Field Container	Senesac	NY	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
26275	Arborvitae (Thuja sp.) T. orientalis 'Green Giant'	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
29169	Jasmine, Star;Confederate (Trachelospermum jasminoides)	Field Container	Koivunen	CA	2015	Over the top	No injury or growth reduction with 21, 42 and 84 fl oz per acre applied twice.
29169	Jasmine, Star;Confederate (Trachelospermum jasminoides)	Field Container	Marble	FL	2015	Over the top	No significant injury or growth reduction with 0.98 and 1.97, slight and acceptable with 3.94 lb ai per acre applied twice.
29169	Jasmine, Star;Confederate (Trachelospermum jasminoides)	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
26663	Hemlock, Western (Tsuga heterophylla)	Field In-Ground	Beste/Frank (ARS)	MD	2007	Over the top	Results not useful because of severe injury caused by very high temperature and drought conditions
26663	Hemlock, Western (Tsuga heterophylla)	Field In-Ground	Beste/Frank (ARS)	MD	2008	Over the top	Moderate to severe injury at 0.97, 1.94 and 3.88 lb ai per acre.
26237	Hemlock, Western (Tsuga heterophylla)	Field Container	Boydston	WA	2007	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre
26237	Hemlock, Western (Tsuga heterophylla)	Field Container	Boydston	WA	2008	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; treated plants saleable
27265	Hemlock (Tsuga sp.)	Field Container	Ahrens/Mervosh	CT	2011	Over the top	Variable crop injury with two sequential applications at 0.97, 1.94, 3.88 lb ai per acre.
27265	Hemlock (Tsuga sp.) T. canadensis	Field Container	Boydston	WA	2011	Over the top	Little to no crop injury with 0.97 and 1.94 lb ai per acre but 3.88 lb ai per acre caused significant injury and reduction in growth. 4x treated plants not saleable.
27265	Hemlock (Tsuga sp.) T. heterophylla	Field Container	Boydston	WA	2010	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all treated plants saleable.
31900	Tulip (Tulipa sp.)	Field In-Ground	Senesac	NY	2014	Over the top	Trial not completed due to crop failure; will be repeated in 2015.
31900	Tulip (Tulipa sp.) 'Ile de France'	Field In-Ground	Miller	WA	2015	Over the top	Moderate foliar injury with 21, 42 and 84 fl oz per acre applied twice; slight reduction of bulb number with 21 and 42, moderate with 84 fl oz.
31900	Tulip (Tulipa sp.) 'Flair'	Field In-Ground	Senesac	NY	2015	Over the top	No injury with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
27422	Viburnum (Viburnum nudum) 'Brandywine'	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	No significant injury at 0.97 and 1.94, severe at 3.88 lb ai per acre; significant growth reduction, unmarketable with 2X and 4X.
27422	Viburnum (Viburnum nudum) V. nudum 'Winterthur'	Field Container	Senesac	NY	2007	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre
27422	Viburnum (Viburnum nudum) 'Winterthur'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury at 0.97, unacceptable at 1.94 and 3.88 lb ai per acre; growth reduction at all rates; 1X plants marketable.
27422	Viburnum (Viburnum nudum) 'Winterthur'	Field Container	Trader	MS	2009	Over the top	Slight injury at 0.97 and 1.94, moderate at 3.88 lb ai per acre; growth reduction at all rates.
27423	Viburnum (Viburnum opulus)	Field Container	Grunwald	OR	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants saleable.
27423	Viburnum (Viburnum opulus) 'Roseum'	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	Slight to moderate injury and growth reduction at 0.97, 1.94 and 3.88 lb ai per acre.
27423	Viburnum (Viburnum opulus) 'Roseum'	Field Container	Trader	MS	2009	Over the top	No significant injury at 0.97, slight and moderate at 1.94 and 3.88 lb ai per acre; growth reduction at all rates.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
27424	Viburnum (<i>Viburnum plicatum</i>) 'Mariesii'	Field Container	Grunwald	OR	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants saleable.
27424	Viburnum (<i>Viburnum plicatum</i>) 'Summer Snowflake'	Field Container	Beste/Frank (ARS)	MD	2008	Over the top	No significant injury at 0.97 and 1.94, slight at 3.88 lb ai per acre; no growth reduction; all plants marketable.
27424	Viburnum (<i>Viburnum plicatum</i>) 'Summer Snowflake'	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre; significant growth reduction with 2X and 4X; all plants marketable.
27424	Viburnum (<i>Viburnum plicatum</i>) V. plicatum 'Shasta'	Field Container	Ahrens/Mervosh	CT	2007	Over the top	Slight injury at 0.97, 1.94 and 3.88 lb ai per acre
27424	Viburnum (<i>Viburnum plicatum</i>) V. plicatum 'Shoshoni'	Field Container	Beste/Frank (ARS)	MD	2007	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre; significant growth reduction but plants considered marketable
27424	Viburnum (<i>Viburnum plicatum</i>) V. plicatum tomentosum 'Shasta'	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
26254	Arrowwood (<i>Viburnum</i> sp.) V. odoratissimum	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
26254	Arrowwood (<i>Viburnum</i> sp.) V. opulus	Field Container	Regan	OR	2007	Over the top	Very slight injury after the first application at 0.97, 1.94 and 3.88 lb ai per acre but slight to moderate injury and growth reduction after the second application.
26254	Arrowwood (<i>Viburnum</i> sp.) V. x 'Juddi'	Field Container	Mickelbart	IN	2008	Over the top	No injury at 2.65, 5.3 and 10.6 lb ai per acre with single application 3 weeks after transplanting.
26254	Arrowwood (<i>Viburnum</i> sp.) V. x pragense	Field Container	Neal	NC	2007	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre
27425	Viburnum (<i>Viburnum tinus</i>)	Field Container	Beste/Frank (ARS)	MD	2009	Over the top	No significant injury at 0.97, slight with complete recovery at 1.94 and 3.88 lb ai per acre; no growth reduction.
27425	Viburnum (<i>Viburnum tinus</i>)	Field Container	Czarnota	GA	2008	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre.
27425	Viburnum (<i>Viburnum tinus</i>) 'Compactum'	Field Container	Neal	NC	2009	Over the top	No injury at 0.97, 1.94 and 3.88 lb ai per acre after 1st, moderate stunting at 4X after 2nd application.
27425	Viburnum (<i>Viburnum tinus</i>) 'Spring Bouquet'	Field Container	Grunwald	OR	2009	Over the top	No injury or growth reduction at 0.97, 1.94 and 3.88 lb ai per acre; all plants saleable.
32954	Palm, Mexican Fan (<i>Washingtonia robusta</i>)	Field Container	Rivera	PR	2014	Over the top	Minor initial injury with quick recovery at 0.98, 1.97 and 3.94 lb ai per acre applied twice; no growth reduction.
29170	Wisteria (<i>Wisteria</i> sp.) W. sempervirens	Field Container	Uber	CA	2009	Over the top	No significant injury at 0.97, 1.94 and 3.88 lb ai per acre.
29170	Wisteria (<i>Wisteria</i> sp.) W. sinensis	Field Container	Persad	OH	2014	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.
29170	Wisteria (<i>Wisteria</i> sp.) W. sinensis	Field Container	Uber	CA	2014	Over the top	No injury with 21, 42 and 84 fl oz per acre applied twice.
27285	Japanese Zelkova (<i>Zelkova serrata</i>)	Field Container	Freiberger	NJ	2009	Over the top	Virtually no injury at 0.97 and 1.94, slight at 3.88 lb ai per acre.
27285	Japanese Zelkova (<i>Zelkova serrata</i>)	Field Container	Persad	OH	2014	Over the top	No significant injury or growth reduction with 0.98, 1.97 and 3.94 lb ai per acre applied twice.

Label Suggestions

For Tower EC, it is suggested that 6 crops not currently on the EPA label be added for over-the-top applications.

Euonymus sp.

Gaura lindheimeri

Leucanthemum maximum

Paeonia lactiflora

Sabal minor

Ternstroemia sp.

It is also suggested that the 5 plant species exhibiting significant injury with over-the-top applications be added to the sensitive ornamental species list.

Catharanthus roseus

Cladrastis sp.

Echeveria sp.

Teucrium chamaedrys

Viburnum opulus

Appendix 1: Contributing Researchers

Dr. John Ahrens <i>(retired)</i>	Connecticut Agricultural Experiment Station Valley Laboratory 153 Cook Hill Road, P.O. Box 248 Windsor, CT
Dr. Ed Beste	University of Maryland LESREC – Salisbury Facility 27664 Nanticoke Road Salisbury, MD 21801
Dr. Rick Boydston	USDA-ARS IAREC 24106 N Bunn Road Prosser, WA 99350
Mr. Luke Case	The Ohio State University Dept. Hort. and Crop Science 2001 Fyffe Ct. Columbus, OH 43210
Dr. Diana Cochran	Iowa State University Department of Horticulture 125 Horticulture Hall Ames, IA 50011
Dr. Mark Czarnota	University of Georgia Department of Horticulture 1109 Experiment St. Griffin, GA 30223
Mr. Joe DeFrancesco	Oregon State University 2040 Cordley Hall Corvallis, OR 97331
Dr. Geoffrey Denny	Mississippi State University Plant & Soil Science Department 246 Dorman Hall Mississippi State, MS 39762
Dr. Jeffrey Derr	Hampton Roads Ag. Exp. Station 1444 Diamond Springs Road Virginia Beach, VA 23455

Mr. Ben Fraelich	USDA-ARS CPES P.O. Box 748 Tifton, GA 31793
Mr. Ray Frank <i>(retired)</i>	6916 Boyers Mill Road New Market, MD 21774
Mr. Tom Freiburger	Rutgers University Cream Ridge Experiment Station 283 Rt. 539 Cream Ridge, NJ 08514
Dr. Charles Gilliam	Auburn University Department of Horticulture 101 Funchess Hall Auburn, AL 36849
Dr. Niklaus Grunwald	USDA-ARS Research Plant Pathologist 3220 NW Orchard Avenue Corvallis, OR 97330
Dr. Brad Hanson	University of California, Davis Department of Plant Sciences One Shields Ave Davis, CA 95616
Mr. Paul Harvey	USDA-ARS 5230 Konnawac Pass Road Wapato, WA, 98951
Dr. Grant Jones <i>(past affiliate)</i>	Davey Tree Expert Company 396 Fenton Lane Suite 608 West Chicago, IL 60185
Dr. Jim Klett	Colorado State University Department of Horticulture and Landscape Architecture Fort Collins, CO 80523
Dr. Heiner Lieth	Department of Plant Sciences University of California One Shield Avenue Davis, CA 95616

Ms. Carrie Mansue (<i>past affiliate</i>)	Rutgers University Adelphia Research Station 594 Halls Mill Road Freehold, NJ 07728
Dr. Chris Marble	University of Florida Mid-Florida Research and Education Center 2725 S. Binion Rd. Apopka, FL 32703
Mr. Michael Marshall (<i>past affiliate</i>)	Michigan State University Department of Horticultural Science East Lansing, MI 48823
Dr. Hannah Mathers (<i>past affiliate</i>)	The Ohio State University Dept. Hort. and Crop Science 2001 Fyffe Ct. Columbus, OH 43210
Dr. Todd Mervosh	Connecticut Agricultural Experiment Station Valley Laboratory 153 Cook Hill Road, P.O. Box 248 Windsor, CT
Dr. Michael Mickelbart	Purdue University Dept. Hort. and Landscape Architecture 625 Agriculture Mall Dr. West Lafayette, IN 47907
Dr. Tim Miller	Washington State University 16650 State Route 536 Mount Vernon, WA 98273-9761
Dr. Joe Neal	North Carolina State University Department of Horticultural Science 262 Kilgore Hall Box 7609, NCSU Raleigh, NC 27695-7609
Dr. Edward Peachey	Dept. of Horticulture Oregon State University 4017 Ag. and Life Sciences Bldg. Corvallis, OR 97331-7304

Dr. Brent Pemberton	Texas A & M University Agricultural Research and Education Center P. O. Box E Overton, TX 75684
Dr. Anand Persad	The Davey Institute 1500 Mantua St Kent, OH 44240
Dr. Michael Reding	USDA-ARS Hort Insects Lab 1680 Madison Ave. Wooster, OH, 44691
Dr. Rich Regan	Oregon State University North Willamette Res. & Ext. Ctr. 15210 NE Miley Rd. Aurora, OR 97002
Dr. Dania Rivera	University of Puerto Rico Recinto Universitario de Mayagüez Servicio de Extensión Agrícola, Call Bo Mayagüez, PR 00681-9000
Dr. Andy Senesac	Long Island Horticultural Research Laboratory 39 Sound Avenue Riverhead, NY 11901
Dr. Brian Trader (<i>past affiliate</i>)	Mississippi State University 158 Dorman Hall, Box 9555 Mississippi State, MS 39762
Mr. Buzz Uber	Crop Inspection Service 31130 Hilltop Drive Valley Center, CA 92082
Dr. Lucia Villavicencio (<i>past affiliate</i>)	Center for Applied Horticultural Research 3742 Blue Bird Canyon Road Vista, CA 92084
Mr. Paul Wade	USDA-ARS US Vegetable Laboratory 2700 Savannah Highway Charleston, SC 29414

Dr. Cheryl Wilen

University of California, San Diego
4444 Overland Ave., Bldg. 2
San Diego, CA 92123

Dr. Anthony Witcher

Tennessee State University
Otis L. Floyd Nursery Research Center
472 Cadillac Lane
McMinnville, TN 37110