



[Environment Horticulture Program Research Summaries](#)

IR-4 Environmental Horticulture Program Management of Borers, Beetles, and White Grubs:

Ambrosia Beetles (*Xylosandrus crassiusculus*, *X. germanus*)
Banded Ash Clearwing Borer (*Podosesia aureocincta*)
Black Vine Weevil (*Otiorhynchus sulcatus*)
Bronze Birch Borer (*Agrilus anxius*)
Flat-headed Apple Tree Borer (*Chrysobothris femorata*)
European Elm Flea Weevil (*Orchestes alni*)
Flea Beetle (*Epitrix* sp.)
Japanese Beetle (*Popillia japonica*)
May/June Beetles (*Phyllophaga* spp.)
Peachtree Borer (*Synanthedon exitiosa*)
Redheaded (Cranberry) Flea Beetle (*Systema frontalis*)
Sri Lankan Weevil (*Myloccerus undatus*)
Strawberry Rootworm (*Paria fragariae* ssp. *fragariae*)
Viburnum Leaf Beetle (*Pyrrhalta viburni*)

Authors: Cristi Palmer and Ely Vea
Date: June 28, 2022

Acknowledgements

Susan Bierbrunner
Diane Infante
Lori Harrison
Karen Sims

Table of Contents

Table of Contents	2
Table of Tables	3
Abstract	8
Introduction	9
Materials and Methods	9
Results: Foliar Feeding Beetles	16
Comparative Efficacy on Black Vine Weevil Adults (<i>Otiorhynchus sulcatus</i>)	16
Comparative Efficacy on Japanese Beetle Adults (<i>Popillia japonica</i>)	16
Comparative Efficacy on Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>)	38
Comparative Efficacy on Red headed Flea Beetle (<i>Systema frontalis</i>)	45
Comparative Efficacy on European Elm Flea Weevil (<i>Orchestes alni</i>)	67
Comparative Efficacy on Sri Lankan Weevil (<i>Myloccerus undatus</i>)	70
Results: Soil Dwelling Immatures	74
Comparative Efficacy on Black Vine Weevil (<i>Otiorhynchus sulcatus</i>)	74
Comparative Efficacy on Japanese Beetle Grubs (<i>Popillia japonica</i>)	84
Comparative Efficacy on May/June Beetle Grubs (<i>Phyllophaga</i> spp.)	86
Comparative Efficacy on Oriental Beetle (<i>Anomala orientalis</i>)	87
Comparative Efficacy on Strawberry Rootworm (<i>Paria fragariae</i>)	94
Results: Trunk and Stem Borers	96
Comparative Efficacy on Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>)	96
Comparative Efficacy on Ambrosia Beetle (<i>Xylosandrus germanus</i>)	106
Comparative Efficacy on Bronze Birch Borer (<i>Agrilus anxius</i>)	107
Comparative Efficacy on Flatheaded Apple Tree Borer (<i>Chrysobothris femorata</i>)	107
Comparative Efficacy on Banded Ash Clearwing Borer (<i>Podosesia aureocincta</i>)	108
Comparative Efficacy on Peachtree Borer (<i>Synanthedon exitiosa</i>)	109
Results: Efficacy Summary by Product	110
Phytotoxicity	113
Label Suggestions	176
Appendix 1: Contributing Researchers	177

Table of Tables

Table 1. List of Products and Rates Tested from 2005 to 2020.....	10
Table 2. Efficacy of several insecticides for black vine weevil adults (<i>Otiorhynchus sulcatus</i>) on Yew (<i>Taxus</i> sp.) or Rhododendron (<i>Rhododendron</i> sp.), Nielsen, 2007.	16
Table 3. Summary of Japanese Beetle Adult (<i>Popillia japonica</i>) Efficacy.....	18
Table 4. Efficacy of several insecticides for <i>Popillia japonica</i> on Sargent Cherry, <i>Prunus sargentii</i> , Alm, 2006a.....	20
Table 5. Efficacy of several insecticides for Japanese Beetle (<i>Popillia japonica</i>) on Shamrock Linden (<i>Tilia cordata</i>) ‘Bailyei’, Alm, 2006b.....	20
Table 6. Efficacy of DPX-E2Y45 and Onyx for Japanese Beetle (<i>Popillia japonica</i>) on Sargent Cherry (<i>Prunus sargentii</i>), Alm, 2007.	21
Table 7. Efficacy of BAS 320i, Onyx, Safari and Tick-EX for Japanese Beetle (<i>Popillia japonica</i>) on <i>Rosa</i> sp., Alm, 2008.	21
Table 8. Efficacy of several insecticides for <i>Popillia japonica</i> on black pussy willows (<i>Salix gracilistyla</i>) ‘Melanostachys’, Braman, 2006.	22
Table 9. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on willow leaves (<i>Salix hakuro nishiki</i>), Braman, 2007.....	23
Table 10. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on rose (<i>Rosa</i> sp.) ‘Blushing’, Braman, 2008.....	23
Table 11. Efficacy of several insecticides on Japanese beetle adults feeding on rose (<i>Rosa</i> sp.), Davis, 2009.....	25
Table 12. Efficacy of several insecticides for <i>Popillia japonica</i> adults on Hibiscus (<i>Hibiscus syriacus</i>), Reding, 2006.	26
Table 13. Efficacy of several insecticides for <i>Popillia japonica</i> adults on Hibiscus (<i>Hibiscus syriacus</i>), Reding, 2007.	27
Table 14. Efficacy of several insecticides for <i>Popillia japonica</i> adults on ‘Julia Child TM. Butter Gold’ Rose, Schultz, 2007.....	30
Table 15. Efficacy of several insecticides for <i>Popillia japonica</i> adults on ‘Pink Velour’ crape myrtle (<i>Lagerstroemia indica</i>), Addesso, 2016.....	30
Table 16. Efficacy of several insecticides for <i>Popillia japonica</i> adults on Rose (<i>Rosa</i> sp.) ‘Louis Phillippe’, Addesso, 2017.....	31
Table 17. Efficacy of several insecticides for <i>Popillia japonica</i> adults on ‘Hopi’ and ‘Pink Velour’ crape myrtle (<i>Lagerstroemia indica</i>), Addesso, 2018.	31
Table 18. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on rose (<i>Rosa</i> sp.), Braman, 2017.....	32
Table 19. Efficacy of several insecticides for <i>Popillia japonica</i> adults on ‘Radrazz’ Knockout Rose (<i>Rosa</i> sp.), Persad, 2017.....	32
Table 20. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on littleleaf linden (<i>Tilia cordata</i>), Potter, 2019.	33
Table 21. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on littleleaf linden (<i>Tilia cordata</i>), Potter, 2020.	34
Table 22. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on crapemyrtle (<i>Lagerstroemia</i> sp.) ‘Hopi’, Addesso, 2020	35
Table 22. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on crapemyrtle (<i>Lagerstroemia</i> sp.) ‘Hopi’, Addesso, 2020	35
Table 22. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on Linden (<i>Tilia cordata</i>), Potter, 2021a	37
Table 22. Efficacy of several insecticides for <i>Popillia japonica</i> adults feeding on Linden (<i>Tilia cordata</i>), Potter, 2021b	37

Table 23. Efficacy of several insecticides for Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>) larval management on Arrowwood viburnum (<i>Viburnum dentatum</i>) – Defoliation Severity, Costa, 2006.	39
Table 24. Efficacy of several insecticides for Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>) larval management on Arrowwood viburnum (<i>Viburnum dentatum</i>) – Defoliation Extent, Costa, 2006..	39
Table 25. Efficacy of several insecticides for Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>) larval management on Arrowwood viburnum (<i>Viburnum dentatum</i>) – Defoliation Severity, Costa, 2007.	39
Table 26. Efficacy of several insecticides for Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>) larval management on Arrowwood viburnum (<i>Viburnum dentatum</i>) – Defoliation Extent, Costa, 2007..	40
Table 27. Foliar damage due to Viburnum Leaf Beetle on <i>Viburnum trilobum</i> treated with thiamethoxam (Actara 25WG) Experiment 1, Holmes, 2007 a.....	40
Table 28. Foliar damage due to Viburnum Leaf Beetle on <i>Viburnum trilobum</i> treated with thiamethoxam (Actara 25WG) Experiment 2, Holmes, 2007 b	41
Table 29. Foliar damage due to Viburnum Leaf Beetle on <i>Viburnum trilobum</i> treated with Rynaxapyr (DPX 2EY45 20SC) Experiment 1, Holmes, 2007 a.....	41
Table 30. Foliar damage due to Viburnum Leaf Beetle on <i>Viburnum trilobum</i> treated with Rynaxapyr (DPX 2EY45 20SC) Experiment 2, Holmes, 2007 b.....	41
Table 31 Summary of Damage ratings (0 – 10) for <i>Viburnum opulus nanum</i> infested with VLB, treated with thiamethoxam (Actara 25WG) Experiment 1, Isaacson, 2007 a	43
Table 32 Summary of Damage ratings (0 – 10) for <i>Viburnum opulus nanum</i> infested with VLB, treated with thiamethoxam (Actara 25WG) Experiment 2, Isaacson, 2007 b	43
Table 33 Summary of Damage ratings (0 – 10) for <i>Viburnum opulus nanum</i> infested with VLB, treated with Rynaxapyr (DPX 2EY45 20SC) Experiment 1, Isaacson, 2007 a.....	43
Table 34 Summary of Damage ratings (0 – 10) for <i>Viburnum opulus nanum</i> infested with VLB, treated with Rynaxapyr (DPX 2EY45 20SC) Experiment 2, Isaacson, 2007 b.....	43
Table 35. Efficacy of several insecticides for Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>) management on Arrowwood viburnum (<i>Viburnum dentatum</i>), Weston, 2007.....	44
Table 36. Detached Leaf Study for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Virginia Sweetspire (<i>Itea virginica</i>) 'Henry's Garnet' and Hydrangea (<i>Hydrangea</i> sp.) 'White Diamonds', Braman, 2012.....	49
Table 37. Damage Rating for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Virginia Sweetspire (<i>Itea virginica</i>) 'Henry's Garnet' and Hydrangea (<i>Hydrangea</i> sp.) 'White Diamonds', Braman, 2012.....	50
Table 38. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Virginia Sweetspire (<i>Itea virginica</i>), Braman, 2013.....	51
Table 39. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Virginia Sweetspire (<i>Itea virginica</i>), Frank, 2013.....	52
Table 40. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Virginia Sweetspire (<i>Itea virginica</i>) 'Little Henry', Sage (<i>Salvia nemorosa</i>) 'Blue Hill', and Stonecrop (<i>Sedum telephium</i>) 'Autumn Joy', Kunkel, 2014.	53
Graph 41. Foliar damage from Jul 1 to Oct 7 caused by Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Virginia Sweetspire (<i>Itea virginica</i>), Cloyd, 2016.....	55
Table 42. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on 'Pee Gee' Hydrangea (<i>Hydrangea paniculata</i> 'Grandiflora'), Gilrein, 2016.	56
Table 43. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Hydrangea (<i>Hydrangea paniculata</i>) 'Baby Lace', Chong, 2017.....	57
Table 44. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Forsythia (<i>Forsythia</i> sp.), Kunkel, 2017a.	58

Table 45. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Forsythia (<i>Forsythia</i> sp.), Kunkel, 2017a.	59
Table 46. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Panicle Hydrangea (<i>Hydrangea paniculata</i>) ‘Vanilla Strawberry’, Frank, 2017.	60
Table 47. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Panicle Hydrangea (<i>Hydrangea paniculata</i> ‘Bokrathirteen’) ‘Sweet Summer’, Gilrein, 2018... ..	61
Table 48. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Panicle Hydrangea (<i>Hydrangea paniculata</i>) ‘Vanilla Strawberry’, Frank, 2019.	62
Table 48. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Panicle Hydrangea (<i>Hydrangea paniculata</i>) ‘Moonrock’, Gilrein, 2019.	63
Table 48. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Panicle Hydrangea (<i>Hydrangea paniculata</i>) ‘Moonrock’, Gilrein, 2020.	64
Table 49. Efficacy of several insecticides applied as foliar sprays for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Panicle Hydrangea (<i>Hydrangea paniculata</i>) ‘Limelight’, Sadof, 2021.	65
Table 48. Efficacy of several insecticides for Redheaded Flea Beetle (<i>Systema frontalis</i>) adults feeding on Panicle Hydrangea (<i>Hydrangea paniculata</i>) ‘Fire and Ice’, Gilrein, 2021.	66
Table 50. Efficacy of several insecticides for European Elm Flea Weevil (<i>Orchestes alni</i>) feeding on Elm (<i>Ulmus</i> sp.) ‘Patriot’ - % leaf area affected, Jones, 2013.	68
Table 51. Efficacy of several insecticides for European Elm Flea Weevil (<i>Orchestes alni</i>) feeding on Elm (<i>Ulmus</i> sp.) ‘Patriot’ - % tree canopy affected, Jones, 2013.	68
Table 52. Efficacy of several insecticides for European Elm Flea Weevil (<i>Orchestes alni</i>) feeding on Elm (<i>Ulmus</i> sp.) ‘Patriot’ - presence of leafmine activity, Jones, 2013.	69
Table 53. Efficacy of several insecticides for Sri Lankan Weevil (<i>Mytillocerus undatus</i>) feeding on Hibiscus (<i>Hibiscus rosa-sinensis</i>) ‘Double Peach’ - % leaf area affected, Dale, 2016.	71
Table 54. Efficacy of several insecticides for Sri Lankan Weevil (<i>Mytillocerus undatus</i>) feeding on Hibiscus (<i>Hibiscus rosa-sinensis</i>) ‘Double Peach’ - % herbivory per 20 leaves, Dale, 2016.	72
Table 55. Efficacy of several insecticides for Sri Lankan Weevil (<i>Mytillocerus undatus</i>) feeding on Hibiscus (<i>Hibiscus rosa-sinensis</i>) ‘Double Peach’ – proportion of leaves with damage, Dale, 2016.	73
Table 56. Summary of Black Vine Weevil (<i>Otiorynchus sulcatus</i>) Efficacy.	75
Table 57. Efficacy of several insecticides for Black Vine Weevil on Yew (<i>Taxus media</i>) ‘Nigra’, Alm, 2006.	77
Table 58. Efficacy of several insecticides for Black Vine Weevil on Strawberry (<i>Fragaria</i> sp.), Cowles, 2007.	78
Table 59. Efficacy of several insecticides for Black Vine Weevil on Strawberry (<i>Fragaria</i> sp.), Cowles, 2008.	80
Table 60. Efficacy of several insecticides for curative control of Black Vine Weevil on Astilbe (<i>Astilbe simplicifolia</i>) ‘Pink Sprite’, Gilrein, 2005*.	81
Table 61. Efficacy of several insecticides for Black Vine Weevil on Yew (<i>Taxus media densiformis</i>), Nielsen, 2006.	82
Table 62. Efficacy of Safari 20SG for Black Vine Weevil on Rhododendron (<i>Rhododendron</i> sp.) ‘Nova Zembla’, Reding, 2004.	83
Table 63. Efficacy of Safari 20SG drenches for Black Vine Weevil on Sedum (<i>Sedum</i> sp.) ‘Vera Jameson’, Reding, 2004.	83
Table 64. Efficacy of Safari 20SG drenches for Black Vine Weevil on Yew (<i>Taxus</i> sp.) ‘Brownii’, Reding, 2004.	84
Table 65. Efficacy of several insecticides for Black Vine Weevil on Sedum (<i>Sedum spurium</i>) ‘Vera Jameson’, Reding, 2006.	84

Table 66. Efficacy of several insecticides for <i>Popillia japonica</i> on rose (<i>Rosa</i> sp.) ‘Caldwell Pink’, Braman, 2006.....	85
Table 67. Efficacy of several insecticides for <i>Popillia japonica</i> on rose (<i>Rosa</i> sp.) ‘Caldwell Pink’, Braman, 2007.....	85
Table 68. Efficacy of several insecticides for May/June Beetle Grubs (<i>Phyllophaga</i> spp.) on Live Oak (<i>Quercus virginiana</i>) ‘Highrise’, Buss, 2008*.	87
Table 69. Summary of Oriental Beetle Grub (<i>Anomala orientalis</i>) Efficacy.....	88
Table 70. Efficacy of several insecticides for Oriental Beetle grubs on Yew (<i>Taxus media</i>) ‘Nigra’, Alm, 2006.....	89
Table 71. Efficacy of several insecticides for Oriental Beetle grubs on Rhododendron (<i>Rhododendron</i> sp.) ‘Scintillation’, Alm, 2008.	89
Table 72. Efficacy of several insecticides for Oriental Beetle on Arborvitae (<i>Thuja</i> sp.) ‘Emerald Giant’ and holly (<i>Ilex</i> sp.) ‘Blue Girl’ – Number of Grubs, Freiberger, 2005.....	90
Table 73. Efficacy of several insecticides for Oriental Beetle on Arborvitae (<i>Thuja</i> sp.) ‘Emerald Giant’ and holly (<i>Ilex</i> sp.) ‘Blue Girl’ – Plant Height, Freiberger, 2005.....	90
Table 74. Efficacy of Seven Products to Reduce Oriental Beetle Populations on Arborvitae, Freiberger, 2009.....	91
Table 75. Efficacy of several insecticides for Oriental Beetle (<i>Anomala orientalis</i>) on lawn type grass, Gilrein, 2005*.	92
Table 76. Efficacy of several insecticides for Oriental Beetle (<i>Anomala orientalis</i>) on lawn type grass, Gilrein, 2007.	92
Table 77. Efficacy of several insecticides for Oriental Beetle on Lilac (<i>Syringa vulgaris</i>), Reding, 2006.	93
Table 78. Efficacy of several insecticides for Oriental Beetle on Lilac (<i>Syringa x chinensis</i>) and White Oak (<i>Quercus alba</i>), Reding, 2009.	93
Graph 79. Efficacy of several insecticides for Strawberry Rootworm larvae (<i>Paria fragariae</i>) on Azalea (<i>Rhododendron</i> sp.), Hesselein, 2005.	94
Table 80. Efficacy of several insecticides for Strawberry Rootworm (<i>Paria fragariae</i>) on Azalea (<i>Rhododendron</i> sp.), Hesselein, 2007.	95
Table 81. Efficacy of Onyx and Talstar for <i>Xylosandrus crassiusculus</i> on Redbud (<i>Cercis canadensis</i>), Ludwig, 2004a.	97
Table 82. Efficacy of Onyx and Talstar for <i>Xylosandrus crassiusculus</i> on Redbud (<i>Cercis canadensis</i>), Ludwig, 2004b.	97
Table 83. Efficacy of Onyx and Talstar for <i>Xylosandrus crassiusculus</i> on Bradford Pear (<i>Pyrus</i> sp), Ludwig, 2004.	97
Table 84. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 1, Mizell, 2005.	98
Table 85. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 2, Mizell, 2005.	98
Table 86. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 3, Mizell, 2005.	98
Table 87. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 4, Mizell, 2005.	98
Table 88. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 5, Mizell, 2005.	99
Table 89. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 1, Mizell, 2006.	100
Table 90. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 2, Mizell, 2006.	100

Table 91. Efficacy of Onyx with and without surfactants for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 1, Mizell, 2007.	101
Table 92. Efficacy of <i>Metarhizium</i> for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 2, Mizell, 2007.	101
Table 93. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Mimosa (<i>Albizia julibrissin</i>) bolts – Experiment 3, Mizell, 2007.	101
Table 94. Efficacy Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Magnolia, Schultz, 2009.	102
Table 95. Efficacy Efficacy of insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Eastern Redbud (<i>Cercis canadensis</i>) 'Forest Pansy', Chong, 2010.....	102
Table 96. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Redbud (<i>Cercis canadensis</i>), Ludwig 2010.....	103
Table 97. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on Redbud (<i>Cercis canadensis</i>), 'Forest Pansy' Chong, 2018.	103
Table 98. Efficacy of several insecticides for Asian Ambrosia Beetle (<i>Xylosandrus crassiusculus</i>) on flooded Flowering Dogwood (<i>Cornus florida</i>), 'Cherokee Princess', Chong, 2019.	105
Table 99. Efficacy of several insecticides for Ambrosia Beetle (<i>Xylosandrus germanus</i>) on Magnolia (<i>Magnolia virginiana</i>), Reding, 2009.	106
Table 100. Efficacy of several insecticides for Bronze Birch Borer (<i>Agrilus anxius</i>) on Weeping European white birch (<i>Betula pendula</i>) – Experiment 1, Nielsen, 2006.	107
Table 101. Efficacy of several insecticides for Bronze Birch Borer (<i>Agrilus anxius</i>) on Weeping European white birch (<i>Betula pendula</i>) – Experiment 2, Nielsen, 2006.	107
Table 102. Efficacy of Discus and Flagship Drenches for Flatheaded Apple Tree Borer (<i>Chrysobothris femorata</i>) on Red Maples (<i>Acer rubrum</i>), Potter, 2005*.....	108
Table 103. Efficacy of several insecticides for Banded Ash Clearwing Borer (<i>Podosesia aureocincta</i>) on Green Ash (<i>Fraxinus pennsylvanica</i>); Nielsen, 2008.	109
Table 104. Efficacy of several insecticides for Peachtree Borer (<i>Synanthedon exitiosa</i>) on Sand Cherry (<i>Prunus cistina</i>), Nielsen, 2009.	109
Table 105. Summary of Efficacy By Product – Borers and Foliar Feeding Beetles	114
Table 106. Summary of Efficacy By Product – White Grubs and Weevils	163

Abstract

Collectively, managing coleopteran insects can be challenging because the adult and larval stages may both cause damage and sometimes occur on different hosts or on different plant parts. While organophosphates, pyrethroids, and neonicotinoids can provide good to excellent control of coleopteran insects, not all products work equally well in all situations. Treatments for borers are very different than treatments targeting white grubs. Developing newer classes of chemistry are important to reduce the environmental consequences and to minimize the development of resistance. Starting with the 2004 Annual Workshop, screening a number of products to manage coleopteran insects became one of the high priority projects for entomology. From 2005 through 2021, 91 products representing 58 different active ingredients were tested for management of adult and larval stages of coleopteran insects. In addition, 10 products representing 10 active ingredients were evaluated for lepidopteran clearwing borers in 2008 and 2009. These products represented both biological and chemical tools. Some products were already registered but more data were needed, or they were considered standards to measure the level of efficacy achieved with other materials. Other products were in development but have not yet been registered with the EPA. While a number of coleopteran and lepidopteran species were tested, only enough experiments were able to be completed on the coleopteran species black vine weevil, Japanese beetle, oriental beetle, red headed flea beetle, Sri Lankan weevil, and viburnum leaf beetles to recommend actions to register or amend labels for these pests.

Introduction

Coleopteran insects have represented some of the most pervasive invasive insects imported into the United States. While not all coleopteran insects causing damage are invasive, a large number impacting growers and landscapes originated outside the US. Collectively, managing coleopteran insects can be challenging because the adult and larval stages may both cause damage and sometimes occur on different hosts or on different plant parts. While organophosphates, pyrethroids, and neonicotinoids can provide good to excellent control of coleopteran insects, not all products work equally well in all situations. Treatments for borers are very different than treatments targeting white grubs. Developing newer classes of chemistry are important to reduce the environmental consequences and to minimize the development of resistance. At the 2004 Annual Workshop, screening a number of products to manage coleopteran insects became one of the high priority projects for entomology. The following research was conducted between 2005 and 2021. Additional research for managing lepidopteran clearwing borers was conducted in 2008 and 2009.

Materials and Methods

Ninety-one insecticides were tested against six species of soil dwelling larvae, six species of borers, and five species of foliar feeding adults and larvae. However, not all products were tested against all species. Depending upon product characteristics, foliar, trunk spray, drench applications, soil incorporation or other application methods were made. A minimum of four plants (replicate treatments) were required with most researchers exceeding this minimum. Insect counts were recorded at timings suitable for each pest. Phytotoxicity when observed was recorded. The following protocols were used: 08-008, 09-017, 10-023, 12-006, 12-019, 13-006, 16-007, 17-007, 18-008, 18-009, 19-008, 19-009 19-022, 20-001, 20-002, and 21-002. For more detailed materials and methods, including application rates for various products, please visit <https://www.ir4project.org/ehc/ehc-registration-support-research/env-hort-researcher-resources/#Protocols> to view and download these protocols.

Products were supplied to researchers (See list of researchers in Appendix 1) by their respective manufacturers.,

Table 1. List of Products and Rates Tested from 2005 to 2021.

Product	Active Ingredient(s)	Manufacturer	Application Method & Rate(s) – <i>per 100 gal water unless otherwise specified</i>		# Trials
Mainspring (A20520A / DPX-HGW86)	Cyantraniliprole	DuPont	Drench	32 fl oz	1
				100 fl oz	1
				0.2 fl oz/ft ht	1
			Foliar	4 fl oz	1
				8 fl oz	2
				16 fl oz	5
			Trunk Spray	32 fl oz	1
Acelepryn (DPX- E2Y45 1.67SC)	Chlorantraniliprole	DuPont	Dipping bolt	15.9 fl oz (47 g ai)	1
				32 fl oz	1
				64 fl oz (189.4 g ai)	1
				255 fl oz (754.4 g ai)	1
			Drench	0.5 fl oz/inch DBH (14.8 ml)	1
				0.8 fl oz	10
				3.2 fl oz	1
				6.47 fl oz	3
				8 fl oz	1
				11.5 fl oz	1
				8 fl oz	1
				16 fl oz	2
				23 fl oz	1
				32 fl oz	2
				46 fl oz	1
				47.9 fl oz	1
				100 fl oz	1
			Foliar	2 fl oz	1
				4 fl oz	1
				6 fl oz	1
				8 fl oz	2
				10 fl oz	11
				16 fl oz	4
				47.9 fl oz	1
			Soil incorporation	1 ppm	1
				2 ppm	1
				4 ppm	1
				5 ppm	1
				10 ppm	1
				20 ppm	1
			Trunk spray	10 fl oz	1
				32 fl oz	2
Aloft	Clothianidin/Bifenthrin	Arysta	Foliar	8 fl oz	1
				15 fl oz	3
			Trunk spray	32 fl oz	1
Ammo 2.5 EC	Cypermethrin	Helena	Dipping bolt	15 fl oz (66.3 g ai)	1
				32.7 fl oz (144.9 g ai)	1
Ancora	<i>Isaria fumosoroseus</i>	OHP	Foliar	16 oz	2
Arena 50WDG	Clothianidin	Arysta	Drench	1.28 oz	1
				1.9 g/inch DBH	1
			Soil incorporation	49 mg/pot	1
Asana XL	Esfenvalerate	DuPont	Dipping bolt	6.8 fl oz (8 g ai)	1
				65.4 fl oz (76.5 g ai)	1
				82 fl oz (95.9 g ai)	1

Product	Active Ingredient(s)	Manufacturer	Application Method & Rate(s) – <i>per 100 gal water unless otherwise specified</i>		# Trials
				131.3 fl oz (153.5 g ai)	1
AzaGuard	Azadirachtin	BioSafe	Foliar	16 fl oz	2
Azatin XL	Azadirachtin	OHP	Dipping bolt	256 fl oz	1
BAS 320i	Metaflumizone	BASF	Dipping bolt	16 fl oz	1
			Drench	12 fl oz	1
				16 fl oz	5
				50 ppm	1
				100 ppm	1
			Foliar	4.5 fl oz	1
				16 fl oz	7
			Soil incorporation	1 ppm	1
				2 ppm	1
				4 ppm	1
				25 ppm	1
				50 ppm	1
				100 ppm	1
				200 ppm	1
BCS 507	BCS 507	Bayer	Foliar	16 oz	3
			Trunk spray	16 oz	2
BeetleGONE	<i>Bacillus thuringiensis galleriae</i> strain SDS-502	Phyllom	Foliar	16 oz	1
				16 lb	4
				10 g/gal	1
Bifenthrin 8% ME	Bifenthrin		Dipping bolt	36.3 fl oz (42.5 g ai)	1
				72.7 fl oz (85 g ai)	1
				145.4 fl oz (170 g ai)	1
Botanigard ES	<i>Beauveria bassiana</i>	BioWorks	Drench	39 uL/pot	1
			Foliar	32 fl oz	4
			Soil incorporation	39 uL/pot	1
Botanigard WP	<i>Beauveria bassiana</i>	Laverlam	Drench	18.7 mg/pot	1
			Soil incorporation	18.7 mg/pot	1
Botaniguard MAXX	Pyrethrins + <i>Beauveria bassiana</i> strain GHA	BioWorks	Foliar	32 fl oz	2
			Trunk spray	32 fl oz	3
BW133		BioWorks	Foliar	5 lb	1
BW238 ES		BioWorks	Foliar	2 qt	1
Cal-Agri 50 1 %	Potassium phosphate	Cal-Agri Products	Drench	128 fl oz	1
Celero 16WSG	Clothianidin	Valent	Dipping bolt	8 oz	1
				8.6 oz (38.8 g ai)	1
				17.1 oz (77.6 g ai)	1
				37.5 oz (170 g ai)	1
			Drench	0.5 oz	3
				1.2 oz	2
				4 oz	2
				12 oz	1
				16 oz	1
				20 oz	1
				4 oz/1320 pots	1
			Foliar	4 oz	8
				6 oz	2
CoreTec	Imidacloprid	Bayer	Soil treatment	3 tablets/inch trunk diameter	1
DEET	N, N-diethyl-m-toluamide		Dipping bolt	40 %	2
				90 %	1
Discus		OHP	Dipping bolt	100 fl oz	1

Product	Active Ingredient(s)	Manufacturer	Application Method & Rate(s) – per 100 gal water unless otherwise specified		# Trials
	Imidacloprid + cyfluthrin			180 fl oz (83.3 + 19.8 g ai)	2
				364 fl oz (169.6 + 39.6 g ai)	1
			Drench	10 fl oz	2
				13 fl oz	1
				1.5 fl oz/inch DBH (44 ml)	2
			Foliar	50 fl oz	2
DuraGuard	Chlorpyrifos	Whitmire	Drench	50 fl oz	1
Dursban 2E	Chlorpyrifos	Dow	Dipping bolt	32 fl oz	1
Dursban Turf (4E)	Chlorpyrifos	Dow	Dipping bolt	13.1 fl oz (92.8 g ai)	1
			Foliar	16 oz	1
Dylox 80S	Trichlorfon	Bayer	Drench	3.75 oz	1
ESP 715	ESP 715	Bayer	Drench	8.55 fl oz	1
			Trunk spray	8.55 fl oz	1
Flagship 0.22G	Thiamethoxam	Syngenta	Top dress	4.25 g/9 sq ft	1
				6 g/pot	1
Flagship 25WG	Thiamethoxam	Syngenta	Drench	0.4 oz	2
				5 oz	1
				8 oz	8
				17 oz	1
				24 oz	1
				0.18 oz per 1000 sq ft	1
			Foliar	8 oz	1
			Top dress	0.09 g /linear foot	1
				0.1 g/9 sq ft	1
			Trunk spray	16 oz/acre	1
Hachi-Hachi 15SC	Tolfenpyrad	SePro	Foliar	21 fl oz	7
				27 fl oz	7
				32 fl oz	5
Hexacide	Rosemary oil	EcoSmart	Drench	1.5 qt	1
ISM-555	ISM-555	Syngenta	Foliar	5.76 fl oz	5
Kontos	Spirotetramat	OHP	Spray	3.4 fl oz	1
KOC22018-8	botanical oil blend	Kemin	Foliar	128 fl oz	3
Lorsban 4E	Chlorpyrifos	Dow	Dipping bolt	66.9 fl oz (474 g ai)	1
				108.3 fl oz (767.5 g ai)	1
Lynx	Pyrethrins	Laverlam	Foliar	16 fl oz	1
Mach 2 2SC	Halofenozide	Dow	Drench	2 lb ai/A	1
				2.9 fl oz per 1000 sq ft	1
Marathon	Imidacloprid	OHP	Drench	20 g per 3000 sq ft	1
Marathon 1G	Imidacloprid	OHP	Soil incorporation	0.1 g ai/gal	1
				7 g/pot	1
Marathon II 2F	Imidacloprid	OHP	Drench	20 g/650 pots	2
				20 g/244 pots	1
				0.2 fl oz/in diam	1
			Drench	0.2 fl oz	1
			Foliar	1.7 fl oz	2
Grandevo (MBI- 203 DF)	<i>Chromobacterium subtsugae</i> strain PRAA4-1T	Marrone BioInnovations	Foliar	1 lb	4
				2 lb	5
				3 lb	1
MBI-203 SC2	<i>Chromobacterium subtsugae</i> strain PRAA4-1T	Marrone BioInnovations	Foliar	128 fl oz	4
MBI-306		Marrone BioInnovations	Foliar	2.5 fl oz	3
				5.0 fl oz	5
Merit 75	Imidacloprid	Bayer	Foliar	10 tsp	1
			Drench	6.4 oz	1

Product	Active Ingredient(s)	Manufacturer	Application Method & Rate(s) – per 100 gal water unless otherwise specified		# Trials
Met 52	<i>Metarhizium anisopliae</i>	Novozymes	Drench	58 oz	2
			Soil incorporation	20 g / 4 gal media	1
<i>Metarhizium anisopliae</i> Strain F52	<i>Metarhizium anisopliae</i>	Novozymes	Dipping bolt	1.3 x 10 ⁹	1
				3.9 x 10 ⁸	1
				3.9 x 10 ⁹	1
			Drench	2.9 g/pot	1
				14.04 cfu/pot	2
				28.08 cfu/pot	2
				56.16 cfu/ pot	1
			Foliar	29 fl oz	3
			Soil incorporation	4.5 x10 ⁸ spores/L	1
				2.9 g/pot	1
				6.25 g/pot	1
NEI 25925	Acetamiprid	Cleary	Trunk spray	4 ml/inch DBH	1
Onyx 2EC	Bifenthrin	FMC	Dipping bolt	32 oz	4
				60 fl oz (212.3 g ai)	5
			Foliar	6.4 fl oz	2
				7.2 fl oz	1
				12.8 fl oz	7
				32 fl oz	1
			Trunk spray	6.4 fl oz	3
				12.8 fl oz	4
				16 fl oz	3
				32 fl oz	2
				102 fl oz	1
Ornazin	Azadirachtin	SePro	Drench	10 oz per 100 gal	1
Orthene	Acephate	Arysta	Drench	12 oz per 100 gal	1
Permethrin 2.5 EC	Permethrin	Bonide Products	Foliar	128 fl oz	2
Perm-UP	Permethrin	UPI	Trunk spray	8 fl oz	2
Precise G & N	Acephate	Purcell Technologies	Soil incorporation	6 g product/can	1
			Top Dress	3 tsp per pot	2
Preferal	<i>Isaria fumosoroseus</i>	SEPro	Foliar	8 oz	1
				16 oz	5
Proclaim	Emamectin benzoate	Syngenta	Dipping bolt	256 fl oz	1
Pyridium	Beta-cyfluthrin	Bayer	Foliar	2.1 fl oz	2
			Trunk spray	2.1 fl oz	1
RTSA 721	RTSA 721	Rainbow Tree Care	Drench	10 ml + 120 bl water / in caliper	3
			Foliar	650 ml	3
			Trunk spray	100 ml per gal	3
Safari 2G	Dinotefuran	Valent	Soil incorporation	1.23 g/pot	1
			Top dress	2.2 g/gal media	3
				60 g/plant	1
Safari 20SG	Dinotefuran	Valent	Drench	12 g/inch DBH	3
				12 oz	4
				24 oz	23
				48 oz	3
				6 g/ft shrub height	1
			Foliar	8 oz	14

Product	Active Ingredient(s)	Manufacturer	Application Method & Rate(s) – <i>per 100 gal water unless otherwise specified</i>		# Trials
			Soil incorporation	24 oz	1
			Basal spray	8 oz	1
			Trunk spray	24 oz	1
Sarissa (IKI-3106)	Cyclaniliprole	ISK	Foliar	22 fl oz	12
				27 fl oz	9
Scimitar GC	Lambda-cyhalothrin	Syngenta	Drench	60 fl oz	1
				0.5 fl oz/ft ht	1
			Foliar	3.2 fl oz	1
				5 fl oz	4
			Trunk spray	5 fl oz	1
Sevin XLR 4F	Carbaryl	Bayer	Drench	6 fl oz/1000 sq ft	2
Sevin SL	Carbaryl	Bayer	Foliar	150 fl oz	1
Sevin (zeta-cyp.)	Zeta-cypermethrin		Foliar	4 fl oz per gal	2
SP3014	SP3014	SePRO	Foliar	13 fl oz	3
				26 fl oz	5
Talstar 0.2G	Bifenthrin	FMC	Soil incorporation	10 ppm	1
				25 ppm	1
				33.7 g / 4 gal media	1
Talstar F	Bifenthrin	FMC	Dipping bolt	32 fl oz	1
				35 fl oz (41 g ai)	1
				115.7 fl oz (135.3 g ai)	1
				183 fl oz (214 g ai)	1
			Drench	25 fl oz	1
				80 fl oz	1
			Foliar	10 fl oz	1
				40 fl oz	2
			Trunk spray	40 fl oz	3
Talstar One	Bifenthrin	FMC	Foliar	21.7 fl oz	1
			Trunk Spray	21.7 fl oz	1
Tempo 2	Cyfluthrin	Bayer	Dipping bolt	32 fl oz	1
Thiodan 3EC	Endosulfan	UCPA	Dipping bolt	15.6 fl oz (83 g ai)	1
TickEx EC	<i>Metarhizium anisopliae</i>	Novozymes	Drench	21 fl oz	1
				29 fl oz	1
			Foliar	29 fl oz	5
TetraCURB Conc.	rosemary oil	Kemin	Foliar	128 fl oz	2
TetraCURB Org.	rosemary oil	Kemin	Foliar	128 fl oz	2
Tolfenpyrad 15EC	Tolfenpyrad	Nichino	Foliar	14 fl oz	1
				21 fl oz	5
			Soil incorporation	10 ppm	1
			Trunk spray	21 fl oz	3
				24 fl oz	1
TriStar 30SG	Acetamiprid	Cleary	Trunk spray	4 oz/inch DBH	1
			Foliar	8 g	1
				12.0 oz	2
TriStar 8.5 SL	Acetamiprid	Cleary	Foliar	25.3 fl oz	6
				32 fl oz	2
TriStar 70WSP	Acetamiprid	Cleary	Foliar	96 g (3.38 oz)	6
				8 oz	2
				12 oz	1
UP-Star Gold	Bifenthrin	UPI	Foliar	8 fl oz	1
V-10433	V-10433	Valent	Foliar	5.5 fl oz	1
				11 fl oz	5

Product	Active Ingredient(s)	Manufacturer	Application Method & Rate(s) – <i>per 100 gal water unless otherwise specified</i>		# Trials
			Trunk spray	13 fl oz	1
				22 fl oz	4
				5.5 fl oz	2
				11 fl oz	2
Velifer		BASF	Foliar	13 fl oz	1
Venerate	<i>Burkholderia rinojensis</i> A396	Marrone	Foliar	1 gal	1
VST-06330	VST-06330	Vesteron	Foliar	16 lb	1
VST-06350	VST-06350	Vesteron	Foliar	3.33 L	3
Xpectro OD	Pythrerins + <i>Beauveria bassiana</i>	Laverlam	Foliar	25.3 fl oz	1
				32 fl oz	3
XXpire / GF-2860 40WP	spinoteram/sulfoxaflo r	Dow	Foliar	2.75	1
				3.5 oz	4
				7 oz	1
Xytect 2F	Imidacloprid		Drench	0.2 fl oz/inch DBH	1

Results: Foliar Feeding Beetles

Comparative Efficacy on Black Vine Weevil Adults (Otiorynchus sulcatus)

Black vine weevil (*Otiorynchus sulcatus*) is a serious pest of ornamental nursery crops (field and container-grown), vineyards, strawberries and hops. Even though it is suspected the black vine weevil (BVW) originated in northern Europe, it was first identified in North America in 1835 and became a notable pest in Missouri by 1871. It is found predominantly in the northern portions of the United States, but its range extends into Virginia and out to the Pacific Northwest.

Throughout Asia, Europe, and North America, black vine weevil adults feed on the foliage and larvae feed on the roots damaging a tremendous variety of species, including azalea, strawberry, begonia, blackberry, blueberry, and cranberry, cyclamen, euonymus, forsythia, fuchsia, hemlock, impatiens, primrose, rhododendron, sedum and yew (http://www.mortonarb.org/res/CLINIC_pests_BlackVineWeevil.pdf; <http://www.entomology.umn.edu/cues/blackvw/blackvh.html>).

IR-4 sponsored a single study on adult BVW and several studies on the larvae [See Comparative Efficacy on Black Vine Weevil (*Otiorynchus sulcatus*)].

Nielsen 2007

In 2007, Nielsen tested six products for their residual efficacy in controlling black vine weevil adults on foliage. Five products were applied to yew and one was applied to rhododendron. Foliage of rhododendron or yew was sprayed Aug 13, and then adults were caged with treated leaves at 1, 3, 7, and 13 DAT. After exposure for 72 hours, the number of dead weevils was counted and any moribund adult was moved to Nontreated foliage and reevaluated 3-days later.

Only the standard Talstar and BAS 320i treatments provided any mortality of adult black vine weevils. Data were similar for all evaluation dates. (Table 2).

No phytotoxicity was observed.

Table 2. Efficacy of several insecticides for black vine weevil adults (*Otiorynchus sulcatus*) on Yew (*Taxus* sp.) or Rhododendron (*Rhododendron* sp.), Nielsen, 2007.

Treatment ^z	Rate Per 100 Gal	Plant Host	Percent Mortality after 3 day exposure ^y
Acelepryn / DPX-E2Y45(chlorantraniliprole)	10 oz	Yew	0
BAS 320i (metaflumizone)	16 oz	Rhododendron	100
<i>Metarhizium</i>	29 oz	Yew	0
Safari 20SG (dinotefuran)	8 oz	Yew	0
Talstar F (bifenthrin)	40 fl oz	Yew	100
Tolfenpyrad	21 oz	Yew	0
Nontreated		Yew	0

^z Treatments were applied August 13, 2007 and evaluated through 13DAT. Four plants per treatment were used.

^y Exposed 5 weevils/replicate in plastic cups with treated foliage.

Comparative Efficacy on Japanese Beetle Adults (Popillia japonica)

The Japanese beetle (*Popillia japonica*) is a widespread and destructive exotic pest of turf, landscape, and ornamental plants in the United States. Outside of its native Japan, it is also found in China, Russia, Portugal, and Canada. Since the first detection in the US in a nursery near Riverton, New Jersey in 1916, it has spread to many states east of the Mississippi River, as well as parts of Wisconsin, Minnesota, Iowa, Missouri, Nebraska, Kansas, Arkansas and Oklahoma. Despite regulatory efforts, it has become established in at least 30 states by 2002. Occasional introductions are made into western states such as California and Oregon when the adult beetles or larvae are shipped in commerce.

The Japanese beetle has a total host range of more than 400 plant species, including turf, ornamentals, fruits, and vegetables. Currently the Japanese beetle is the most widespread pest of turf and costs the turf and ornamental industry approximately \$450 million each year in management alone (<http://ohioline.osu.edu/hyg-fact/2000/2504.html>, <http://edis.ifas.ufl.edu/IN630>).

IR-4 sponsored several studies on adult JB and a couple studies on the larvae [See Comparative Efficacy on Japanese Beetle Grubs (*Popillia japonica*)].

From 2006 to 2019, seven researchers examined the efficacy of 29 products and unregistered materials for managing Japanese beetle adults. The products tested included Acelepryn, Aloft, BAS 320i, BotaniGard ES, BotaniGard MAXX, Celero 16WSG, Flagship 25WG, IKI-3106, Marathon II, Onyx 2EC, Precise, Pyridium, Safari 2G, Safari 20SG, Scimitar, Talstar, TickEx EC, Tolfenpyrad, TriStar, V-10433, VST-006350, and Xpectro. In these experiments, the assessment typically made was percent leaf damaged by adult beetle feeding. Even though Reding (Table 12 and Table 13) and Adesso (Table 15) were unable to achieve statistical separation in 3 experiments, in the 13 experiments conducted by Adesso, Alm, Braman, Davis, Persad, and Schultz, there were clear differences in efficacy. As the standards, Sevin, Tristar SL, and bifenthrin in Onyx and Talstar, provided good to excellent control. Acelepryn performed well achieving greater than 95% control in seven out of nine experiments. Aloft provided good to excellent control in 2 tests, IKI-3106 in 6 tests, and BAS 320i, in 2 of 3 tests.

Table 3. Summary of Japanese Beetle Adult (*Popillia japonica*) Efficacy.

Treatment	Rose							Sargent Cherry	Sargent Cherry	Shamrock Linden	Black Pussy Willow	Willow	Crape Myrtle	Littleleaf Linden	Littleleaf Linden
	14 DAT	14 DAT	19 DAT	4 DAT	7 DAT	14 DAT		19 DAT	17 DAT	19 DAT	2 WAT	4 DAT	7 DAT	7 DAT	1 DAT
	Schultz 2007	Alm 2008	Braman 2008	Davis 2008	Addesso 2017	Braman 2017	Persad 2017	Alm 2006a	Alm 2007	Alm 2006b	Braman 2006	Braman 2007	Addesso 2018	Potter 2019a	Potter 2019b
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	++			+				++	++	++	+/-	++		++	++
Aloft (clothianidin+bifenthrin)			++	+											
AzaGuard (azadirachtin)														-	-
BAS 320i (metaflumizone)	++	-	++												
BCS 507														+	++
BotaniGard ES					-	-	+/-								
BotaniGard MAXX (pythrerins + <i>Beauveria bassiana</i>)													-	-	++
Celero 16WSG (clothianidin)	++							-		+	-	+/-			
Flagship 25WG (thiamethoxam)				+							-				
IKI-3106 (cyclaniliprole)					+	++	++						+	++	++
KOC22018-8 (botanical oil blend)														-	+
Marathon II (imidacloprid)						++									
Onyx 2EC (bifenthrin)		+	++					++	++	+					
Precise (acephate)											-				
Pyridium (beta-cyfluthrin)													+		
Safari 2G (dinotefuran)		+/-		+/-											
Safari 20SG (dinotefuran)	+		+/-	+/-				-		+/-	-	+/-			
Scimitar (lambda- cyhalothrin)				+							+				

Treatment	Rose							Sargent Cherry	Sargent Cherry	Shamrock Linden	Black Pussy Willow	Willow	Crape Myrtle	Littleleaf Linden	Littleleaf Linden
	14 DAT	14 DAT	19 DAT	4 DAT	7 DAT	14 DAT		19 DAT	17 DAT	19 DAT	2 WAT	4 DAT	7 DAT	7 DAT	1 DAT
	Schultz 2007	Alm 2008	Braman 2008	Davis 2008	Addesso 2017	Braman 2017	Persad 2017	Alm 2006a	Alm 2007	Alm 2006b	Braman 2006	Braman 2007	Addesso 2018	Potter 2019a	Potter 2019b
Sevin (carbaryl)														++	++
Talstar F (bifenthrin)	++		++				++								
TetraCURB Conc. (rosemary oil)														-	-
TetraCURB Org (rosemary oil)														-	-
TickEx (<i>Metarhizium anisopliae</i>)	- (by 28 DAT 100% contr ol was achie ved)	-	-									-			
Tolfenpyrad EC	++		-	-											
TriStar 8.5 SL (acetamiprid)					+								++		
TriStar 70WP (acetamiprid)								++		++	+/-	+			
V-10433													+/-	-	-
VST-006350					-										
Xpectro OD (pythrerins + <i>Beauveria bassiana</i>)						-	+/-								

¹ Rating Scale: ++ = clearly statistically better than Nontreated and greater than 95% control; + = statistically better than Nontreated and between 85 and 95% control; +/- statistically better than Nontreated with control between 70 and 85%; - = statistically equivalent to Nontreated and/or efficacy less than 70%.

² Where more than one rate or application type for a product was included in the experiment and each performed statistically different, the better rating is provided in this table.

Alm 2006

In 2006, Alm conducted two experiments to determine efficacy of five products to control Japanese beetle adults on foliage of Sargent cherry and shamrock linden. In both experiments, foliar applications were made and then a single terminal branch was selected from each plant with the top five treated leaves rated for percent Japanese beetle feeding damage on Jul 10, Jul 19, and Jul 31 (10, 19 and 31 DAT, respectively). The mean percent feeding damage was calculated for the top five leaves. An overall mean percent feeding damage was calculated from four replicates.

For Sargent cherry, there was no statistically significant Japanese beetle feeding damage to any of the treated trees at 10 DAT (Table 4), but by 19 and 31 DAT there was significantly more feeding damage on the Nontreated and the Safari treated trees than any of the other treatments. Acelepryn, Celero, Onyx, and TriStar 70WP significantly reduced feeding; however, Celero only provided approximately 50% control whereas the other treatments provided 95% control or better.

For shamrock linden, all treatments provided statistically significant control of Japanese beetle feeding by the time trees were rated on 10 and 19 DAT (Table 5). When trees were rated on 31 DAT, there was significant feeding damage on the Safari treated trees compared to the other treatments. With the exception of Safari at 31 DAT, all treatments provided 82% control or better based upon percent feeding damage.

No phytotoxicity was observed.

Table 4. Efficacy of several insecticides for *Popillia japonica* on Sargent Cherry, *Prunus sargentii*, Alm, 2006a.

Treatment	Rate Per 100 Gal	Average Percent Leaf Damage ^z		
		10 July 10 DAT	19 July 19 DAT	31 July 31 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz	0.0 a (100%)	1.0 bc (98%)	2.0 c (98%)
Celero 16 WSG (clothianidin)	6 oz	0.0 a (100%)	23.0 b (59%)	40.0 b (50%)
Onyx 2EC (bifenthrin)	12.8 fl oz	0.0 a (100%)	2.5 bc (95%)	2.0 c (98%)
Safari 20SG (dinotefuran) + Capsil	8 oz + 6 fl oz	2.5 a (0%)	49.0 a (12%)	65.0 a (19%)
TriStar 70WP (acetamiprid)	3.38 oz	0.0 a (100%)	0.0 c (100%)	0.0 c (100%)
Nontreated		2.5 a (0%)	55.5 a (0%)	80.5 a (0%)

^z Means in the same column followed by the same letter are not significantly different, ($P = 0.05$, LSD test).

Table 5. Efficacy of several insecticides for Japanese Beetle (*Popillia japonica*) on Shamrock Linden (*Tilia cordata*) ‘Bailyei’, Alm, 2006b.

Treatment	Rate Per 100 Gal	Average Percent Leaf Damage ^z		
		10 July 10 DAT	19 July 19 DAT	31 July 31 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz	0.5 b (93%)	0.0 b (100%)	3.5 c (96%)
Celero 16 WSG (clothianidin)	6 oz	0.5 b (93%)	3.5 b (88%)	14.5 c (82%)
Onyx 2EC (bifenthrin)	12.8 fl oz	0.0 b (100%)	3.5 b (88%)	11.0 c (86%)
Safari 20SG (dinotefuran) + Capsil	8 oz + 6 fl oz	2.0 b (71%)	5.0 b (83%)	41.0 b (48%)
TriStar 70WP (acetamiprid)	3.38 oz	0.0 b (100%)	0.0 b (100%)	0.0 c (100%)
Nontreated		7.0 a (0%)	29.5 a (0%)	78.5 a (0%)

^z Means in the same column followed by the same letter are not significantly different, ($P = 0.05$, LSD test).

Alm 2007

In 2007, Alm compared the efficacy of Acelepryn with Onyx to control Japanese beetle adults on foliage of Sargent cherry (Table 6). In this experiment, after foliar applications two terminal branches were selected from each plant, and ten treated leaves were rated for percent Japanese beetle feeding damage on Jul 18, Jul 27, and

Aug 14 (8, 17 and 34 DAT, respectively). The mean percent feeding damage was calculated for these ten leaves. An overall mean percent feeding damage was calculated from five replicates.

In this experiment, Acelepryn and Onyx significantly reduced Japanese beetle adult feeding by 8 DAT. By 17 DAT, percent control was 91% or greater for all rates of both products. This level of management remained through 34 DAT, the last reading date.

No phytotoxicity was observed.

Table 6. Efficacy of DPX-E2Y45 and Onyx for Japanese Beetle (*Popillia japonica*) on Sargent Cherry (*Prunus sargentii*), Alm, 2007.

Treatment	Rate Per 100 Gal	Average Percent Leaf Damage (% Control) ^z		
		18 July 8 DAT	27 July 17 DAT	14 Aug 34 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	2 fl oz	6.4 b (56%)	2.3b (94%)	3.3b (92%)
Acelepryn (DPX-E2Y45; chlorantraniliprole)	4 fl oz	4.4bc (69%)	3.4 b (91%)	4.2b (90%)
Acelepryn (DPX-E2Y45; chlorantraniliprole)	8 fl oz	2.0c (86%)	0.7b (98%)	1.6b (96%)
Onyx 2EC (bifenthrin)	12.8 fl oz	2.4bc (83%)	1.9b (95%)	2.0b (95%)
Nontreated		14.4a (0%)	39.0a (0%)	41.3a (0%)

^z Means in the same column followed by the same letter are not significantly different, ($P = 0.05$, LSD test).

Alm 2008

In 2008, Alm compared the efficacy of several insecticides to control Japanese beetle adults on foliage of *Rosa* sp. In this study, BAS 320i, Onyx and Tick-EX were sprayed to runoff while Safari 2G was broadcast on media and watered in. Treatments were evaluated 7 and 14 days after treatment (DAT) by rating percent Japanese beetle feeding damage on the entire plant. An overall mean percent feeding damage was calculated from seven replicates.

Onyx and Safari treatments significantly prevented feeding damage by Japanese beetle adults (Table 7). At 7 and 14 DAT, percent control was 94-96 % from Onyx and 84-86 % from Safari. BAS 320i and Tick-EX significantly reduced feeding damage but not at a commercially acceptable level.

No phytotoxicity was observed.

Table 7. Efficacy of BAS 320i, Onyx, Safari and Tick-EX for Japanese Beetle (*Popillia japonica*) on *Rosa* sp., Alm, 2008.

Treatment	Rate Per 100 Gal	Average Percent Leaf Damage (% Control) ^z	
		7 DAT	14 DAT
BAS 320i (metaflumizone)	16 fl oz	33.6 b (45)	35.7 b (43)
Onyx 2EC (bifenthrin)	6.4 fl oz	2.7 c (96)	3.7 c (94)
Safari 2G (dinotefuran)	60 g/plant	8.6 c (86)	10.0 c (84)
Tick-EX EC ^a	29 fl oz	37.9 b (38)	40.0 b (36)
Nontreated		61.4 a (0)	62.9 a (0)

^a*Metarhizium anisopliae* strain F52

^z Means in the same column followed by the same letter are not significantly different, ($P = 0.05$, LSD test).

Braman 2006

In 2006, Braman compared seven treatments for the impact on the number of Japanese beetles on black pussy willows and the percent defoliation. Plants were sprayed on June 13, 2006 and arranged in an area with

historically heavy Japanese beetle populations in southeastern Spalding County, GA. Beetle density was recorded at one week, two weeks, and one month post application (1, 2, and 4 WAT, respectively). A final damage (% defoliation) assessment was made one month after application. All data were subjected to analysis of variance using the GLM procedure of SAS and means were separated using LSD.

In this test, the Japanese beetle pressure was high. Within one week beetle density varied by treatment, with Scimitar, Flagship, TriStar and Acelepryn providing significant reductions relative to the Nontreated control (Table 8). Celero, Safari and Precise were statistically similar to the Nontreated control 1 WAT. At two weeks post treatment all except Safari had significantly reduced beetle densities relative to the control. Defoliation was reduced in all treatments relative to the control and was least on plants treated with Scimitar, Precise, TriStar or Acelepryn.

Throughout the experiment, no phytotoxicity was observed. At the completion of the study, there was no discernable difference in growth among the treatments (data not shown).

Braman 2007

In 2007, Braman compared five treatments for efficacy on Japanese beetle adults on willow. In this experiment, 5 adult beetles were caged on treated foliage using nylon screen bags; at 1 and 4 days after exposure (6 and 9 DAT), the number of surviving beetles were counted, and at Day 4 foliar feeding was estimated as percent feeding damage. Data were subjected to analysis of variance using the GLM procedure of SAS and mean separation is by LSD.

Beetle survival was high for all treatments one day after caging on the willows (Table 9). After 4 days, however, survival was significantly reduced in all but the Tick Ex cages. Fewest beetles survived in the Acelepryn treatment. TriStar and Acelepryn provided the best reduction in damage.

Table 8. Efficacy of several insecticides for *Popillia japonica* on black pussy willows (*Salix gracilistyla*) ‘Melanostachys’, Braman, 2006.

Treatment	Rate Per 100 Gal	Mean no. Beetles per plant ^z			Mean % defoliation 4 WAT
		1 WAT	2 WAT	4 WAT	
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 oz	3.8 bc (83)	6.3 cd (77)	0.0 b (100)	20.5 cd
Celero 16WSG (clothianidin)	4 oz	15.6 ab (28)	16.1 bc (41)	0.0 b (100)	44.5 b
Flagship 25WG (thiamethoxam)	8 oz	5.9 bc (73)	12.6 bcd (54)	0.0 b (100)	29.0 c
Precise (acephate)	3 tsp/pot	22.4 a (0)	11.5 bcd (58)	0.0 b (100)	18.5 cd
Safari 20SG (dinotefuran)	8 oz	14.4 abc (34)	21.1 ab (22)	0.0 b (100)	44.5 b
Scimitar (lambda-cyhalothrin)	5 oz	0.4 c (98)	2.9 d (89)	0.0 b (100)	18.5 cd
Tristar 70WSP (acetamiprid)	96 g	1.3 bc (94)	5.1 d (81)	0.0 b (100)	14.5 d
Nontreated		21.8 a (0)	27.1 a (0)	0.3 a (0)	61.0 a

^z Means followed by the same letter are not significantly different, P> 0.05

Table 9. Efficacy of several insecticides for *Popillia japonica* adults feeding on willow leaves (*Salix hakuro nishiki*), Braman, 2007.

Treatment ^z	Rate Per 100 Gal	Survival (5 adult beetles caged on leaves) ^y		Percent Damage Day 4 (9 DAT)
		Day 1 (6 DAT)	Day 4 (9 DAT)	
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 oz	4.40 a	1.56 c	1.44 d
Celero 16WSG (clothianidin)	4 oz	4.29 a	2.30 bc	21.10 b
Safari 20SG (dinotefuran)	8 oz	4.56 a	2.70 b	17.20 bc
Tick Ex EC (<i>Metarhizium anisopliae</i>)	29 oz	4.86 a	4.40 a	57.50 a
Tristar 70WSP (acetamiprid)	96 g	4.50 a	2.50 bc	4.90 cd
Nontreated		4.56 a	4.40 a	56.00 a

^z Treatments were applied on July 17, 2007.

^y Means followed by the same letter are not significantly different, P> 0.05

Braman 2008

In 2008, Braman compared eight treatments for efficacy on Japanese beetle adults on rose. In this experiment, 5 adult beetles were caged on treated foliage using BugDorm insect rearing sleeves. At 7 and 19 days after treatment and caging (7 and 19 DAT), the number of surviving beetles were counted, and at Day 19 total damage was recorded using a rating scale from 0 to 10, with 0= no damage and 10= 100% defoliation. Data were subjected to analysis of variance using the GLM procedure of SAS and means were separated using LSD.

Acelepryn, Aloft, BAS 320i, Onyx and Talstar provided excellent control of Japanese beetle adults, based on number of survival at 7 and 19 DAT (Table 10). This resulted in virtually no defoliation on roses treated with these products. Safari provided significant but less effective control. Tolfenpyrad showed essentially similar beetle survival as the Nontreated but significantly reduced leaf feeding damage. Tick-Ex was non-effective, showing beetle survival and feeding damage similar to the Nontreated check.

Table 10. Efficacy of several insecticides for *Popillia japonica* adults feeding on rose (*Rosa* sp.) ‘Blushing’, Braman, 2008.

Product	Rate Per 100 Gal	No. feeding beetles 7 DAT	No. living beetles 19 DAT	Leaf Damage Rating ^a 19 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz	0.4 c	0 b	0.1 c
Aloft (clothianidin+bifenthrin)	8 fl oz	0 c	0 b	0 c
BAS 320i (metaflumizone)	16 fl oz	0 c	0 b	0 c
Onyx (bifenthrin)	12.8 fl oz	0.2 c	0 b	0 c
Safari 20SG (dinotefuran)	8 oz	0.8 bc	0.8 b	1.0 c
Talstar One (bifenthrin)	21.7 fl oz	0 c	0 b	0 c
Tick-Ex (<i>Metarhizium anisopliae</i>)	29 oz	1.4 ab	2.0 a	4.8 a
Tolfenpyrad	21 fl oz	1.4 ab	2.4 a	3.6 b
Nontreated		2.0 a	2.2 a	5.0 a

^z Means followed by the same letter are not significantly different, P> 0.05

^a Rating: 1 = 10 % defoliation, 10 = 100 % defoliation.

Davis 2009

In 2009, Davis initiated an outdoor assessment of products on the feeding of Japanese beetle adults. Container roses were positioned next to a planting of Linden where JB adults were present in previous years. At no time were there any differences between any of the treatments or the Nontreated check on any of the sample days while the plants were in the field. This is primarily due to the low numbers of JB adults in the area and the number of other suitable hosts nearby.

To enable usable data to be generated, treated leaves were placed into arenas and the amount of leaf tissue consumed by adult JB was measured. At 3 days after the arenas were set-up, the Acelepryn, Safari drench, Safari and Tolfenpyrad treatments were not significantly different from the Nontreated check. The number of adults left alive in the arenas was significantly different from the Nontreated check in the BAS 320i, Flagship drench and Scimitar treatments. The arenas were evaluated again the next day. Mortality had increased in all of the treatments. All of the treatments except for Safari and Tolfenpyrad were significantly different from the Nontreated check treatment. The 3 applications of BAS 320i, 3 applications of Scimitar and single drench application of Flagship were the superior treatments. Four days after the arenas were set-up, the Nontreated check had 35% of the foliage in the arena consumed. All of the treatments were significantly different from the Nontreated check with regards to amount of foliage consumed. The Flagship and Scimitar treatments protected the foliage the best.

Table 11. Efficacy of several insecticides on Japanese beetle adults feeding on rose (*Rosa* sp.), Davis, 2009.

Treatment	Rate	Application Type	Lab assay 5 JB initial,after 3 days	% Skeletonized after 3 days	Lab assay 5 JB initial,after 4 days	% Skeletonized after 4 days
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz/100 gal	foliar	2.86b	9.29b	1.57 ab	9.29b
BAS 320i 22% liquid (metaflumizone)	16 fl oz/100 gal	foliar	1.72 a	9.29b	1.00 ab	9.29b
Flagship 25WG (thiamethoxam)	24 oz/100 gal - 43 oz soln/gal/media	drench	1.57 a	3.71 a	1.00 ab	2.86 a
Safari 20SG (dinotefuran)	24 oz/100 gal - 4 oz soln/gal/media	drench	2.86b	6.43 a	2.00bc	7.86b
Safari 2G (dinotefuran)	2.2 g/gal/media	top of potting soil	3.72b	5.00 ab	3.00cd	5.71 ab
Scimitar CS	5 fl oz/100 gal	foliar	1.72 a	5.00 ab	0.86 a	5.00 ab
Tolfenpyrad 15EC & adjuvant (tolfenpyrad)	21 fl oz/100 gal & 0.25% v/v	foliar	4.00b	17.86c	3.71d	19.29c
Nontreated			4.00 b	26.43 c	3.86 d	35.00 d

Reding 2006

In this experiment, 4 products (Acelepryn, Celero, Safari, and Tristar) were tested in a no choice feeding bioassay along with a greenhouse study. Leaves were collected 7 days after application, and stems were inserted into water soaked oasis cubes and placed into lidded containers with two Japanese beetle adults per leaf sample. Leaf area measurements were taken after 12 days of feeding. There was no mortality during the trial. Square centimeters consumed were compared using analysis of variance (ANOVA) with no differences found between treatments (Table 12).

In addition to running a bioassay, insect counts and feeding damage were recorded 2 weeks after Japanese beetles were observed feeding on test plants or 17 days after treatment (17 DAT). Insect counts did not prove to be a good source of efficacy data due to the continual flight of adults during the counting process. Percent of leaves damaged from feeding and leaves with more than 25% of the leaf eaten were compared using analysis of variance (ANOVA) with no differences found between treatments (Table 12).

Throughout the experiment, no phytotoxicity was observed. All plants were marketable at the completion of the study, and there was no discernable difference in growth among the treatments.

Table 12. Efficacy of several insecticides for *Popillia japonica* adults on Hibiscus (*Hibiscus syriacus*), Reding, 2006.

Treatment	Rate Per 100 Gal	Mean leaf area removed in Feeding Bioassay	Feeding Data	
			% Leaves with Damage	% Leaves with >25% Leaf Area Removed
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	47.9 oz	1.91	34.7	10.4
Celero 16WSG (clothianidin)	4 oz	1.49	38.9	7.5
Safari 20SG (dinotefuran)	8 oz	1.24	32.9	4.6
Tristar 70WSP (acetamiprid)	96 g	1.17	34.6	5.7
Nontreated		0.88	36.2	5.0

Reding 2007

In this experiment, four products were tested to assess efficacy: Acelepryn, BAS 320i, Celero, and TickEx. Feeding damage on leaves was recorded in the field 7 and 14 days after first treatment and 14 days after second treatment for leaves. There were differences in severity of damage between treatments for leaf feeding, with BAS 320i, and Tick Ex demonstrating significantly higher damage than the Nontreated control (Table 13. Efficacy of several insecticides for *Popillia japonica* adults on Hibiscus (*Hibiscus syriacus*), Reding, 2007. Acelepryn and Celero 16WSG were equivalent to the Nontreated control. Flower numbers were recorded (7 and 14 days after second treatment) rather than flower damage because when Japanese beetles feed on flowers they become so severely damaged that they fall from the plant and cannot be accurately counted. Comparing the difference in number of blooms was a more accurate method to evaluate efficacy. Some treatments had a larger number of hibiscus flowers than Nontreated plants but were not statistically significant (Table 13).

No phytotoxicity was detected on any of the insecticide treated plants.

Table 13. Efficacy of several insecticides for *Popillia japonica* adults on Hibiscus (*Hibiscus syriacus*), Reding, 2007.

Treatment ^z	Rate Per 100 Gal	Percent Leaves Damaged ^y			Mean Number of Blossoms	
		7 DAT	14 DAT	28 DAT	21 DAT	28 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz	9.7	11.8a	9.5a	40.5	57.7
BAS 320i (metaflumizone)	4.5 oz	13.4	21.1c	16.8c	68.4	79.4
Celero 16 WSG (clothianidin)	4 oz	11.0	13.9ab	12.6abc	55.7	70.7
Tick Ex EC (<i>Metarhizium anisopliae</i>)	29 oz	17.4	17.4bc	14.7bc	65.8	80.6
Nontreated		11.7	12.5a	11.5ab	47.8	58.5

^zTreatments were applied as foliar sprays on 7/10/2007 and 7/30/2007.

^yMeans within columns followed by the same letter are not significantly different ANOVA ($P = 0.05$), means separated by LSD ($\alpha = 0.05$).

Schultz 2007

In this experiment, Schultz examined Acelepryn, BAS 320i, bifenthrin, Celero, *Metarhizium anisopliae*, Safari SG, and tolfenpyrad for their ability to control Japanese beetle adults on rose. Applications were made either Jun 25 or 28 as foliar sprays or drenches (Table 14). After foliage had dried (and one week after the Safari drench), 10 Japanese beetle adults were introduced into a mesh cage on a single branch. Mortality was assessed weekly (7, 14, and 21 DAT). After the 21 DAT counts, dead and remaining live beetles were removed, mesh bags were relocated on the plant, and new adults introduced. Mortality of the newly introduced beetles was taken for 7 and 14 days after introduction (28 and 35 DAT).

At 7 DAT all treatments, except Safari applied as a drench and Tick Ex, had significantly higher adult mortality for caged beetles than the Nontreated plants. At 7 DAT, Acelepryn and BAS320i had 100% mortality 7 DAT. By 14 DAT, Bifenthrin and Celero also exhibited 100% mortality. At 21 DAT, Safari and tolfenpyrad reached 100% mortality.

There was a high background population of beetles, and observations were taken on their feeding. Throughout the experiment roses treated with Acelepryn, bifenthrin, BAS 320i, Celero, and Safari SG sustained no damage to the foliage regardless of adult beetle mortality. Foliage in the other treatments (*Metarhizium*, tolfenpyrad, and Nontreated check) did exhibit foliar damage.

Addesso 2016

In this experiment, Addesso examined BotaniGard ES, IKI-3106, Tristar SL, and Xpectro OD applied as foliar sprays for their ability to control Japanese beetle adults on crape myrtle. Treatments were applied on Jul 15 and all, except BotaniGard, reapplied on Jul 29. Adult Japanese beetles were released onto plants and a PEG lure was deployed at the center of the container pad to draw beetles into the plot from nearby fields. Beetles were free to migrate into and out of the plot at will.

Adult live counts were too low to analyze by day and were pooled across all observations for statistical analysis. No significant differences were observed in either live or dead beetles across treatments. Percent defoliation was recorded at 3, 7, 14, 21 or 28 DAT. No differences in defoliation were observed due to large variation between treatment ratings.

No phytotoxicity was observed on any of the plants in any of the treatments.

Addesso 2017

In 2017, Addesso examined BotaniGard ES, IKI-3106, Tristar SL, and VST- 006350 applied as foliar sprays for their ability to control Japanese beetle adults on rose. The experiment was carried out within a 4x10x8 (WxLxH) cage with no thrips screening atop a ground cloth. IKI-3106, Tristar were applied once on Jul 12, while

Botanigard and VST- 006350 were applied twice on Jul 12 and 19. One hundred Japanese beetle adults were added to the cage at the beginning of the experiment and again on Day 7 after reapplication of the Botanigard and VST treatments. The lowest amount of feeding damage was observed consistently in the low and high rates of IKI-3106 followed by the Tristar standard (Table 16). BotaniGard and VST- 006350 did not significantly reduce feeding damage.

Addesso 2018

In 2018, Addesso compared efficacy of several products applied as foliar sprays for their ability to control Japanese beetle adults on two crape myrtle varieties. The experiment was carried out within two large screened field cages, one variety in each cage, to guarantee feeding pressure on plants. Each cage received a known number of Japanese beetle adults, as follows: D0 (Jul 9) = 100, D2 = 100, D4 = 50, D7 = 50, D10 = 75, D11 = 25. All treatments were applied on Jul 9 (D0), and reapplied on Jul 16 for BotaniGard MAXX and V-10433. The Tristar standard, followed by Pyridium and IKA-3106, resulted in the lowest number of live beetles and amount of feeding damage (Table 17). Tristar and Pyridium also provided significant increase in plant growth.

Braman 2017

In 2017, Braman compared eight treatments for efficacy on Japanese beetle adults on rose. Treatments were applied from Jul 3 to Jul 18 on a schedule shown in Table 18. Plants were placed in 2ft. x 2ft. x 2ft. mesh cages after applications were made. Japanese beetles were counted and 10 beetles were added to each of the cages. Dead beetles were counted and damage ratings were taken at each of the evaluation days. Ten additional beetles per cage were added after each evaluation.

IKI-3106 at both rates and the standard treatment Marathon II showed excellent control and remained significantly different from other treatments, but not from each other during each evaluation. BotaniGuard remained significantly different from the Nontreated up until the last evaluation, whereas Expectro showed differences after both applications. Number of dead beetles was significantly different at Days 7 and 28. Mortality at Day 7 consistently reflects damage to foliage. By day 28, the Nontreated plants had very little foliage left to support beetle feeding and likely contributed to beetle mortality. No phytotoxicity was observed.

Persad 2017

In 2017, Persad compared efficacy of several products applied foliar sprays for their ability to control Japanese beetle adults on container grown Knockout rose. All treatments were applied on Jul 17 and 24, and all, except Botanigard, reapplied on Jul 31. Talstar and IKI-3106 provided effective control of a high adult infestation; Xpectro and BotaniGard were less effective (Table 19).

Potter 2019

In 2019, Potter examined efficacy of several products applied foliar sprays for efficacy on Japanese beetles adults on littleleaf linden. Treatments were applied from Jul 9 to Jul 24 on a schedule shown in less effective (Table 20. Efficacy of several insecticides for *Popillia japonica* adults feeding on littleleaf linden (*Tilia cordata*), Potter, 2019. Table 20). Treatments were evaluated at 3, 7, 14, 21 and 28 DAT from the 9 July date. On the assay day, undamaged linden foliage was collected from each branch and placed in individual clear 8 oz deli containers. Five female JB were then added per dish. Dishes were held at 26° C; 14:10 light-dark cycle for 24 h; then the JB were removed and each cohort was transferred to a separate tray that was opened in direct sunlight. This "fly-off" assay rapidly distinguishes healthy JB, which invariably take flight, from dead or moribund individuals. Photocopies of the leaves were scanned, with missing area determined with the software program Paint.NET v4.2.1. For a second trial, ten female JB beetles were directly sprayed until wet and then transferred to a clean dish and provided with Nontreated linden foliage. After 24h, beetles were subjected to fly-off assay and foliage feeding was assessed.

The results of both experiments are shown in Table 20. In the first trial, Carbaryl, IKI-3106 (cyclaniliprole) and BCS-507 significantly reduced both feeding and beetle survival. Acelepryn also greatly reduced beetles' feeding on all dates. Acelepryn also caused beetle mortality, but not quite as consistently as the above mentioned treatments. For the feeding reduction assays, the treatments Acelepryn, BCS-507, IKI-3106 and carbaryl were consistently better than the Nontreated check. BotaniGard MAXX, V-10433 and AzaGuard provided some feeding reduction compared to the Nontreated control, but not consistently on all dates and not to the same level

as the above mentioned treatments. In the second trial, Acelepryn, BCS-507, IKI-3106, and carbaryl reduced survival by 95%. The treatments Acelepryn, BCS-507, BontiGard MAXX, IKI-3106, KOC22018-8, and carbaryl reduced feeding by 90%.

Table 14. Efficacy of several insecticides for *Popillia japonica* adults on ‘Julia Child TM. Butter Gold’ Rose, Schultz, 2007.

Treatment	Rate Per 100 Gal	Mean Number Dead Beetles per Cage (Corrected Percent Control)					
		First Challenge (after foliar applications had dried)				Second Challenge	
		0 DAT	7 DAT	14 DAT	21 DAT	28 DAT	35 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	Foliar – 10 fl oz	3.6 b (34%)	10.0 a (100%)	10.0 a (100%)	10.0 a (100%)	8.3 b (81%)	10.0 a (100%)
BAS 320i (metaflumizone)	Foliar – 16 oz	1.1 c (8%)	10.0 a (100%)	10.0 a (100%)	10.0 a (100%)	1.5 c (6%)	10.0 a (100%)
Celero 16WSG (clothianidin)	Foliar – 4 oz	5.8 a (57%)	9.8 a (98%)	10.0 a (100%)	10.0 a (100%)	10.0 a (100%)	10.0 a (100%)
Onyx (bifenthrin)	Foliar – **	5.6 a (55%)	6.9 b (68%)	10.0 a (100%)	10.0 a (100%)	10.0 a (100%)	10.0 a (100%)
Safari 20SG (dinotefuran)	Drench – 24 oz	--	1.3 c (10%)	9.5 a (94%)	10.0 a (100%)	10.0 a (100%)	10.0 a (100%)
Tick Ex EC (<i>Metarhizium anisopliae</i>)	Foliar – 29 oz	0.4 c (1%)	0.5 c (2%)	2.5 b (10%)	9.1 a (74%)	9.9 a (99%)	9.9 a (99%)
Tolfenpyrad	Foliar – 14 oz	2.1 bc (19%)	7.4 b (73%)	9.6 a (95%)	10.0 a (100%)	9.0 ab (89%)	9.6 a (95%)
Nontreated		0.3 c (0%)	0.3 c (0%)	1.7 c (0%)	6.6 b (0%)	1.0 c (0%)	1.4 b (0%)

** no rate provided in report so the high label rate of 12.8 fl oz per 100 gal was assumed.

Table 15. Efficacy of several insecticides for *Popillia japonica* adults on ‘Pink Velour’ crape myrtle (*Lagerstroemia indica*), Adesso, 2016.

Treatment	Rate Per 100 Gal	Number of Beetles		Percent Leaf Damage				
		Live	Dead	Day 3	Day 7	Day 14	Day 21	Day 28
BotaniGard ES (<i>Beauveria bassiana</i>)	32 fl oz	1	5	2.82	5.68	4.01	3.28	1.23
IKI-3106 (cyclaniliprole)	22 fl oz	0	7	1.38	2.57	2.73	2.39	2.58
IKI-3106 (cyclaniliprole)	27 fl oz	1	11	2.20	8.58	2.35	3.36	3.40
Tristar 8.5 SL (acetamiprid)	32 fl oz	0	15	1.84	3.30	1.13	0.56	1.07
Xpectro OD (Pythrerins + <i>Beauveria bassiana</i>)	25.3 fl oz	0	8	1.46	3.95	3.61	4.07	3.35
Nontreated	-	0	5	2.53	2.69	2.58	2.58	1.20
		ns	ns	ns	ns	ns	ns	ns

Table 16. Efficacy of several insecticides for *Popillia japonica* adults on Rose (*Rosa* sp.) 'Louis Phillippe', Addesso, 2017.

Treatment	Rate Per 100 Gal	Number of Beetles ^x		Percent Leaf Damage ^y		
		Live	Dead	Day 3	Day 7	Day 14
BotaniGard ES (<i>Beauveria bassiana</i>)	32 fl oz	6 ab	2 a	6.4a	10.8a	11.5ab
IKI-3106 (cyclaniliprole)	22 fl oz	0 b	4 a	0.6b	0.3b	1.2c
IKI-3106 (cyclaniliprole)	27 fl oz	0 b	1 a	0.8b	0.9b	2.4c
Tristar 8.5 SL (acetamiprid)	32 fl oz	1 b	5 a	0.8b	0.6b	5.6bc
VST-006350 + 0.1% LI700	1 L/30 gal	16 a	6 a	2.0b	11.5a	17.4a
Nontreated	-	2 b	2 a	2.2b	5.1ab	13.7a

^x Count data was analyzed using a generalized linear model with a log link and a negative binomial distribution.

^y Percent data was analyzed untransformed with a generalized linear model under a normal distribution (PROC Genmod, SAS Institute 2016).

Table 17. Efficacy of several insecticides for *Popillia japonica* adults on 'Hopi' and 'Pink Velour' crape myrtle (*Lagerstroemia indica*), Addesso, 2018.

Treatment	Rate Per 100 Gal	Applic. Dates	Percent Damage ^x			No. Beetles		Old Leaves Eaten	New Leaves Eaten	Plant Growth (CSI)
			Day 3	Day 7	Day 14	Live	Dead			
Botaniguard MAXX (pyrethrins + <i>Beauveria bassiana</i> strain GHA)	32 fl oz	7/9, 16	5.6 ab	13.1 ab	13.7 bc	3.0 ab	0.1 a	19.1 ab	5.2 c	57.5 bc
IKI-3106 (cyclaniliprole)	22 fl oz	7/9	0.7 bcd	1.7 b	10.3 bc	0.8 cd	0.3 a	2.9 d	6.3 ab	40.5 c
IKI-3106 (cyclaniliprole)	27 fl oz	7/9	1.3 bcd	2.4 b	9.8 bc	0.9 dc	0.4 a	7.8 c	4.2 d	87.6 abc
Pyridium (beta-cyfluthrin)	2.1 fl oz	7/9	0.4 cd	1.6 b	11.7 bc	0.4 de	0.2 a	3.3 d	3.0 d	113.8 a
Tristar 8.5 SL (acetamiprid)	25.3 fl oz	7/9	0.3 d	1.1 b	4.0 c	0.1 e	0.4 a	2.3 d	1.9 d	128.3 a
V-10433 (V-10433)	11 fl oz	7/9, 16	2.2 bcd	7.9 ab	12.8 bc	1.5 bc	0.2 a	11.8 bc	5.8 bc	95.9 ab
V-10433 (V-10433)	22 fl oz	7/9, 16	5.6 abc	19.8 a	18.8 b	4.6 a	0.0 a	23.9 a	6.7 a	55.8 b
Nontreated	-	-	9.0 a	19.7 a	36.4 a	4.4 a	0.3 a	33.7 a	7.2 a	54.5 bc

^x Foliar feeding damage.

Growth measurement and percent defoliation were analyzed with a general linear model fit to a normal distribution (PROC GLM, SAS Institute). Count data including live and dead JB and new and old leaves damaged were analyzed with a generalized linear model fit to a negative binomial distribution. Significant treatment values were separated in pair-wise tests using least significant differences.

Table 18. Efficacy of several insecticides for *Popillia japonica* adults feeding on rose (*Rosa* sp.), Braman, 2017.

Product	Rate Per 100 Gal	Applic. Dates	% Leaf Damage ^x					No. Dead Beetles				
			Day 3	Day 7	Day 14	Day 21	Day 28	Day 3	Day 7	Day 14	Day 21	Day 28
BotaniGard ES (<i>Beauveria bassiana</i>)	32 fl oz	Jun 12, 19	12.50 b	35.00 b	62.50 b	74.17 b	81.67 a	2.83 a	5.67 c	9.00 a	8.50 a	6.83 b
Xpectro OD (Pythrerins + <i>Beauveria bassiana</i>)	32 fl oz	Jun 12, 19	11.80 b	55.83 ab	71.67 b	79.50 ab	79.67 a	2.66 a	6.67 BC	8.67 a	7.33 a	8.67 ab
IKI-3106 (cyclaniliprole)	22 fl oz	Jun 12	0.33 c	0.67 c	1.83 c	6.33 c	10.00 b	4.00 a	9.67 BA	8.5 a	7.17 a	7.33 b
IKI-3106 (cyclaniliprole)	27 fl oz	Jun 12	0.67 c	0.33 c	0.33 c	4.17 c	5.50 b	4.50 a	11.00 A	9.67 a	7.33 a	8.83 ab
Marathon II (imidacloprid)	1.7 fl oz	Jun 12	0.50 c	0.50 c	1.00 c	7.83 c	7.33 b	3.33 a	10.83 A	8.83 a	7.50 a	7.67 ab
Nontreated	-	-	27.50 a	77.50 a	95.83 a	95.5 a	98.00 a	2.66 a	6.33 C	6.50 a	10.75 a	9.75 a

^x Means followed by the same letter are not significantly different, (LSD test).

Table 19. Efficacy of several insecticides for *Popillia japonica* adults on 'Radrazz' Knockout Rose (*Rosa* sp.), Persad, 2017.

Treatment	Rate Per 100 Gal	Percent Mortality ^x				Percent Herbivory				Percent Defoliation			
		3 DAT	7 DAT	14 DAT	43 DAT	3 DAT	7 DAT	14 DAT	43 DAT	3 DAT	7 DAT	14 DAT	43 DAT
BotaniGard ES	32 fl oz	51 c	57 c	66 b	74 b	27 b	30 b	29 b	30 b	33 b	33 b	31 b	17 c
IKI-3106	22 fl oz	80 a	90 a	100 a	100 a	20 c	20 c	20 c	20 c	15 c	15 c	15 c	10 d
IKI-3106	27 fl oz	85 a	90 a	100 a	100 a	20 c	20 c	20 c	20 c	15 c	15 c	15 c	10 d
Talstar	10 fl oz	91 a	94 a	100 a	100 a	18 c	18 c	16 c	19 c	14 c	14 c	13 c	10 d
Xpectro OD	32 fl oz	64 b	73 b	83 a	83 b	16 c	20 c	20 c	20 c	27 b	31 b	32 b	22 b
Nontreated		0 d	0 d	0 c	0 c	60 a	71 a	79 a	85 a	55 a	65 a	76 a	80 a

^x Means followed by the same letter are not significantly different, (Student-Newman-Keuls, P=.05)

Table 20. Efficacy of several insecticides for *Popillia japonica* adults feeding on littleleaf linden (*Tilia cordata*), Potter, 2019.

Treatment	Rate Per 100 Gal	Applic. Dates	First Experiment								Second Experiment	
			No. surviving JB adults (out of 5) ^x				Leaf Area Eaten ^y				JB adults (out of 10)	Leaf Area Eaten ^z
			Day 3	Day 7	Day 14	Day 28	Day 3	Day 7	Day 14	Day 28	1 DAT	1 DAT
Acelepryn	8 fl oz	7/9, 24	0.5±0.3 c	1.2±0.5 b	1.5±0.5 b	1.8±0.5 b	0.3±0.1 d	0.4±0.1 c	0.4±0.1 d	0.3±0.1 d	0.6±0.4 ef	0.04±0.02 d
AzaGuard (azadirachtin)	16 fl oz	7/9, 16, 24	4.8±0.2 a	5.0 a	5.0 a	4.2±0.6 a	9.5±0.6 bc	13.4±1.1 b	10.5±1.5 abc	8.9±1.7 ab	6.6±0.7 b	2.9±0.6 d
BCS-507	16 oz	7/9, 24	0.5±0.3 c	0.5±0.3 c	0.5±0.3 c	0.3±0.3 c	0.4±0.1 d	1.1±0.5 c	0.4±0.1 d	0.3±0.03 d	0.0 f	0.04±0.04 d
BotaniGard MAXX	32 fl oz	7/9, 16	4.5±0.2 a	5.0 a	4.7±0.2 a	4.3±0.4 a	9.4±2.0 bc	11.3±1.1 b	7.6±1.3 c	6.9±0.8 ab	1.0±0.5 ef	0.0 d
IKI-3106 (cyclaniliprole)	27 fl oz	7/9, 24	0.2±0.2 c	0.0 c	0.7±0.3 c	1.5±1.0 bc	0.2±0.1 d	0.3±0.04 c	0.6±0.2 d	1.7±0.9 d	0.0 f	0.1±0.1 d
KOC22018-8 (botanical oil blend)	128 fl oz	7/9, 16, 24	4.7±0.2 a	4.7±0.2 a	5.0 a	3.5±0.7 a	11.1±0.8 ab	12.3±2.5 b	9.8±1.9 bc	5.7±1.4 bc	2.6±0.7 de	1.4±0.3 d
Sevin SL	150 fl oz	7/9, 24	0.0 c	0.0 c	0.0 c	0.5±0.5 bc	0.2±0.2 d	0.1±0.03 c	0.5±0.3 d	2.8±2.1 cd	0.0 f	0.0 d
TetraCURB Conc. (rosemary oil)	128 fl oz	7/9, 16, 24	4.7±0.3 a	5.0 a	5.0 a	4.3±0.4 a	11.3±1.3 ab	14.1±1.6 ab	11.1±1.0 ab	9.7±1.5 a	5.0±1.0 bc	7.9±1.9 c
TetraCURB Org (rosemary oil)	128 fl oz	7/9, 16, 24	4.8±0.2 a	4.8±0.2 a	4.8±0.2 a	4.5±0.2 a	9.7±1.3 bc	12.1±0.6 b	11.4±1.0 ab	8.0±0.8 ab	4.4±1.3 cd	8.2±1.5 c
V-10433	22 fl oz	7/9, 16, 24	4.0±0.6 ab	4.8±0.2 a	5.0 a	4.3±0.3 a	6.7±1.8 c	13.2±1.6 b	8.8±0.8 bc	10.5±0.6 a	7.0±0.9 b	11.5±1.5 b
Nontreated	-	-	3.3±0.8 b	5.0 a	4.7±0.2 a	4.5±0.2 a	13.2±1.9 a	16.8±2.0 a	13.2±1.2 a	9.7±2.4 a	9.8±0.3 a	17.2±1.9 a

^x Means followed by the same letter are not significantly different, ($P > 0.05$; LSD All-Pairwise Comparisons Test).

^y Leaf area eaten in cm² by 5 female JB in a 24 hour period.

^z Leaf area eaten in cm² by 5 female JB in a 24 hour period.

Table 21. Efficacy of several insecticides for *Popillia japonica* adults feeding on littleleaf linden (*Tilia cordata*), Potter, 2020.

Treatment	Mean surviving JB adults (out of 5)				Mean sq cm of leaf tissue eaten ^a			
	3 DAT	7 DAT	14 DAT	28 DAT	3 DAT	7 DAT	14 DAT	28 DAT
BW238NWP	5.0 A	5.0 A	5.0 A	4.8±0.2 A	18.5±2.1 AB	14.8±3.3 CD	15.5±1.4 ABCDE	14.3±1.1 AB
BW238NES	5.0 A	4.8±0.2 A	4.5±0.3 A	4.5±0.3 A	15.7±2.6 BC	18.3±3.5 ABCD	11.8±1.7 DEF	12.5±2.5 ABC
BW133	5.0 A	5.0 A	4.8±0.2 A	5.0 A	14.4±2.0 BC	16.0±1.2 BCD	18.9±2.6 AB	15.1±1.7 AB
ISM-555 + Caspsil 7 day app	1.3±0.4 B	1.2±0.8 B	1.5±0.5 C	4.5±0.5 A	2.8±0.5 D	4.4±0.9 E	5.0±0.7 GH	2.9±0.7 DE
ISM-555+ Caspsil 14 day app	0.3±0.2 C	1.5±0.5 B	2.5±0.6 B	3.3±0.5 B	4.2±0.2 D	5.0±0.5 E	7.2±0.9 FG	7.4±0.7 CD
MBI-203 SC2	5.0 A	5.0 A	4.8±0.2 A	5.0 A	16.7±2.6 B	17.4±1.6 BCD	14.9±1.1 BCDE	12.4±1.6 ABC
MBI-306	5.0 A	5.0 A	4.7±0.2 A	4.8±0.2 A	18.5±1.0 AB	20.2±1.3 AB	17.7±3.4 ABC	13.0±2.0 AB
RSTA 721 Basal Trunk Spray	4.8±0.2 A	4.5±0.5 A	4.7±0.2 A	5.0 A	17.6±2.9 AB	19.4±1.9 ABC	13.7±3.3 CDE	17.4±4.4 A
RTSA 721 Soil Drench	4.8±0.2 A	4.8±0.2 A	4.5±0.2 A	4.7±0.2 A	22.8±2.1 A	23.1±3.4 A	20.2±2.3 A	15.4±0.7 AB
RTSA 721 Foliar Spray	0.2±0.2 C	1.3±0.9 B	0.0 D	0.7±0.3 C	0.5±0.2 D	0.6±0.1 E	0.5±0.1 H	1.3±0.3 E
SP3014+Capsil	5.0 A	4.8±0.2 A	4.7±0.2 A	5.0 A	16.0±2.6 BC	19.6±1.9 ABC	16.9±1.1 ABCD	10.9±3.1 BC
V-10433	5.0 A	5.0 A	4.8±0.2 A	4.5±0.3 A	11.2±1.1 C	13.5±0.9 D	11.6±1.7 EF	12.6±2.1 ABC
Carbaryl	0.0 C	0.0 C	0.0 D	0.5±0.5 C	0.3±0.2 D	0.1±0.03 E	0.4±0.2 H	2.5±1.4 DE
Nontreated	4.8±0.2 A	5.0 A	4.7±0.2 A	4.5±0.3 A	17.5±1.9 AB	19.8±1.7 ABC	17.0±0.9 ABC	16.7±2.0 A

^a Leaf area eaten in cm² by 5 female JB in a 24 hour period.

For statistical analysis of surviving JB adults (Fly-off assay): 3DAT – $F_{13,65}=191.87$, $p=0.00$; 7 DAT – $F_{13,65}=25.96$, $p=0.00$; 14 DAT – $F_{13,65}=56.10$, $p=0.00$; 28 DAT – $F_{13,65}=28.08$, $p=0.00$. Within column, means followed by the same letter are not significantly different ($P > 0.05$; LSD All-Pairwise Comparisons Test). For statistical analysis of sq cm of leaf tissue eaten: 3DAT – $F_{13,65}=15.67$, $p=0.00$; 7 DAT – $F_{13,65}=16.67$, $p=0.00$; 14 DAT – $F_{13,65}=12.98$, $p=0.00$; 28 DAT – $F_{13,65}=8.27$, $p=0.00$. Within column, means followed by the same letter are not significantly different ($P > 0.05$; LSD All-Pairwise Comparisons Test).

Adesso 2020

In 2020, Adesso assessed efficacy of several products applied as foliar sprays to manage Japanese beetle adults on crape myrtle. Pre infestation RTSA 721 trunk sprays or drenches were applied on May 25; the other products were applied as foliar sprays starting on June 15 with different schedules based on the protocol. Adult JB were collected using baited traps deployed 24 hours prior to introduction to the test cages. Due to low catch numbers on Day 0, an equal number of beetles were added to the cages each day for 4 days until 100 beetles per cage were reached. Another 100 beetles were added to each cage immediately following the applications of the Day 7 foliar treatments. The applications with the fewest live adults were ISM-555 and Onyx (Table 22), with BW238 ES and RTSA 721 foliar applications were also low but not statistically different from nontreated plants. Some treatments had higher numbers of live adults than the nontreated. Similar trends were observed with other assessments including number of damaged leaves and percent feeding damage (data not shown). Damage on nontreated plants was unexpectedly lower than many of the plants treated with other products, suggesting that either some aspect of the products were attractive to the beetles or that early damage on plants with ineffective products resulted in later aggregation of beetles on the plants resulting in higher levels of damage.

Table 22. Efficacy of several insecticides for *Popillia japonica* adults feeding on crapemyrtle (*Lagerstroemia sp.*) ‘Hopi’, Adesso, 2020

Treatment - Rate	Application Method and Date(s)	Live Adults Feeding (DAT)				Total Dead Adults
		3 DAT	7 DAT	14 DAT	Total Live	
BW 133	Foliar - 6/15/20; 6/24/20	3	3	5	11 abc	0
BW 238 ES	Foliar - 6/15/20; 6/24/20	1	2	2	5 bcd	0
BW 238 WP	Foliar - 6/15/20; 6/24/20	0	5	7	12 abc	0
ISM-555	Foliar - 6/15/20	0	1	1	2 cd	1
	Foliar - 6/15/20; 6/24/20	0	0	0	0 d	1
MBI-203	Foliar - 6/15/20; 6/24/20	2	5	11	18 ab	1
MBI-306	Foliar - 6/15/20; 6/24/20	1	3	7	11 abc	0
Onyx	Foliar - 6/15/20	0	0	0	0 d	0
RTSA 721	Drench - 5/25/20	0	1	8	9 abcd	0
	Foliar - 6/15/20	0	2	4	6 bcd	5
	Basal Trunk Spray - 5/25/20	0	1	10	11 abc	2
SP3014	Foliar - 6/15/20; 6/24/20	2	10	14	26 a	3
V-10433	Foliar - 6/15/20; 6/18/20; 6/24/20; 6/27/20	1	3	10	14 ab	1
Nontreated		0	2	10	12 abc	0

Live beetle counts were pooled across dates and analyzed with a generalized linear model using a negative binomial distribution with a log link as is appropriate for count data. Differences in the number of live beetles across treatments were observed ($\chi^2_{(13)} = 30.47$, $P = 0.004$). Pairwise comparisons were made using LSD ($\alpha = 0.05$).

Adesso 2021

During 2021, Adesso continued efficacy assessments for Japanese beetle adults on crape myrtle. Pre infestation RTSA 721 trunk sprays or drenches were applied on May 26; the other products were applied as foliar sprays starting on June 16 with different schedules based on the protocol. Adult JB were collected using baited traps deployed 24 hours prior to introduction to the test cages. Due to low feeding damage from high wind from the cooling system, the experiment was moved to screened walk-in cages. Japanese beetle adults were introduced to the greenhouse and walk-in cages throughout the experiment. The applications with the fewest live adults were ISM-555 and Onyx (Table 23), with MBI-306 at 5 fl oz, RTSA 721 foliar applications, and V-10433 were also low but not statistically different from nontreated plants. Treatments that had the least percentage of leaves with feeding damage by 28 DAT included Onyx, RTSA 721 foliar applications, MBI-306 high rate, and V-10433. The RTSA 721 drench application caused yellowing of leaf margins; transient injury was observed with MBI203 and ISM-555. No differences were observed in canopy growth among the treatments.

Table 23. Efficacy of several insecticides for *Popillia japonica* adults feeding on crapemyrtle (*Lagerstroemia sp.*) ‘Hopi’, Adesso, 2020

Treatment	PRNumbers	Number of Live JB (DAT)	
-----------	-----------	-------------------------	--

Rate per 100 gal		3	7	14	21	28	Total Live	Total Dead
ISM555 5.76 fl oz	Foliar - 06/16/21, 06/23/21	0	0	1	0	1	2ef	11d
	Foliar - 06/16/21, 06/23/21, 06/30/21	0	2	1	0	1	4def	24bcd
MBI306 5 fl oz	Foliar - 06/16/21, 06/23/21	0	1	3	2	3	9bcdef	7d
MBI306 2.5 fl oz	Foliar - 06/16/21, 06/23/21	0	0	6	0	30	36abc	13cd
MBI203 128 fl oz	Foliar - 06/16/21, 06/23/21	0	1	4	3	73	81a	17bcd
OnyxPro 7.2 fl oz	Foliar - 06/16/21, 06/30/21	0	0	0	0	0	0f	13cd
RTSA721 100 ml/1 gal	Trunk - 05/26/21	1	2	5	3	7	18abcde	94a
RTSA721 10 ml + 120 ml water / inch caliper @ 1 inch height	Drench - 05/26/21	0	2	6	3	48	59a	36abc
RTSA721 650 ml	Foliar - 05/26/21	0	0	4	0	7	11bcdef	17bcd
SP3014 26 fl oz	Foliar - 06/16/21, 06/23/21, 06/30/21	0	0	3	0	46	49ab	21bcd
SP3014 13 fl oz	Foliar - 06/16/21, 06/23/21, 06/30/21	1	1	4	0	52	58ab	14bcd
V-10433 13 fl oz	Foliar - 06/16/21, 06/19/21, 06/23/21, 06/26/21, 06/30/21	1	0	3	1	1	6cdef	11d
Nontreated	-	0	0	11	5	5	21abcd	46ab

Live beetle counts were pooled across dates and analyzed with a generalized linear model using a negative binomial distribution with a log link as is appropriate for count data. Differences in the number of live beetles across treatments were observed ($\chi^2_{(13)} = 30.47$, $P = 0.004$). Pairwise comparisons were made using LSD ($\alpha = 0.05$).

Potter 2021

During 2021, Potter implemented two experiments studying new insecticides for impact on Japanese Beetle adults on linden trees. In the first experiment, RTSA-721 was applied as foliar spray, trunk injection or soil injection (drench) and compared to the consumer formulation of Sevin containing zeta-cypermethrin. Trunk and soil injections were applied May 7, 2021, while foliar applications were applied on July 5, 2021; evaluations started July 5. The second experiment contained foliar applications only of seven different insecticides which were applied on July 5, the same date as evaluations began. For both experiments, adult beetles were live trapped and starved for 24 hours before being placed into containers with leaves from treated plants. After 24 hours of exposure, beetles were transfer to separate containers and their ability to fly was assessed to indicate health. Leaves were scanned electronically and feeding damage (missing area) was determined. In the first experiment, the RTSA 721 foliar application and Sevin (zeta-cypermethrin) reduced the number of viable adult beetles and feeding damage, while the trunk and soil injections of RTSA-721 were not effective. In the second experiment, zeta-cypermethrin generally reduced adult viability and feeding damage. ISM-555 did not reduce the number of viable beetles based on flight assessments but feeding damage was reduce in comparison to the nontreated controls. Acelepryn reduced viability on some dates, but feeding damage was greatly reduced through 14DAT. MBI-306 at 5 fl oz significantly reduced adult viability, but there was no impact on feeding. MBI-203, SP3014, and V-10433 did not impact adult viability or feeding in this experiment.

Table 24. Efficacy of several insecticides for *Popillia japonica* adults feeding on Linden (*Tilia cordata*), Potter, 2021a

Treatment Rate per 100 gal	Mean (\pm SE) number of viable beetles (out of 5) ¹					Mean (\pm SE) cm ² of leaf tissue eaten ^a				
	0 DAT	3 DAT	7 DAT	14 DAT	28 DAT	0 DAT	3 DAT	7 DAT	14 DAT	28 DAT
RSTA 721 Basal Trunk Injection*	5.0 A	5.0 A	5.0 A	5.0 A	4.6 A	31.7 A	40.3 A	22.2 B	31.3 A	17.8 A
RTSA 721 Soil Injection*	4.0 A	4.8 A	4.8 A	4.8 A	5.0 A	27.2 A	33.3 B	30.9 A	32.6 A	20.3 A
RTSA 721 Foliar Spray	1.8 B	0.2 B	0.8 B	1.0 B	0.8 B	11.2 BC	0.7 C	1.9 C	1.6 C	1.4 B
Zeta-cypermethrin (Sevin®)	0.0 C	0.2 B	3.8 A	0.6 B	0.2 B	6.0 C	0.4 C	0.8 C	0.2 C	0.2 B
Untreated	4.2 A	5.0 A	5.0 A	4.8 A	4.8 A	24.4 AB	30.8 B	24.2 AB	23.2 B	18.2 A

Table 25. Efficacy of several insecticides for *Popillia japonica* adults feeding on Linden (*Tilia cordata*), Potter, 2021b

Treatment Rate per 100 gal	Mean (\pm SE) number of viable beetles (out of 5) ¹					Mean (\pm SE) cm ² of leaf tissue eaten ^a				
	0 DAT	3 DAT	7 DAT	14 DAT	28 DAT	0 DAT	3 DAT	7 DAT	14 DAT	28 DAT
ISM-555 +Capsil@0.05% 5.76 fl oz applied weekly	4.8 AB	5.0 A	4.8 A	5.0 A	5.0 A	13.2 BC	13.1 CD	9.5 BC	13.3 CD	5.7 D
ISM-555 +Capsil@0.05% 5.76 fl oz applied biweekly	4.8 AB	5.0 A	5.0 A	4.8 A	4.8 A	7.4 C	8.2 DE	5.0 CD	12.8 D	6.8 D
MBI-203 SC2 128 fl oz	3.8 BC	4.6 A	5.0 A	4.4 A	4.6 A	14.8 B	22.8 B	13.9 AB	19.5 BCD	20.3 AB
MBI-306 5 fl oz	2.8 CD	3.8 B	3.0 B	2.6 B	3.4 B	25.0 A	20.3 BC	18.4 A	22.6 AB	20.7 A
MBI-306 2.5 fl oz	5.0 A	5.0 A	5.0 A	5.0 A	4.8 A	22.4 A	28.8 AB	17.4 A	24.3 AB	17.6 ABC
SP3014 + Capsil@0.05% 26 fl oz	5.0 A	5.0 A	4.8 A	5.0 A	5.0 A	26.5 A	24.9 AB	16.1 A	19.7 BC	18.0 ABC
V-10433 11 fl oz	5.0 A	5.0 A	5.0 A	5.0 A	5.0 A	21.5 A	25.8 AB	18.3 A	21.4 AB	15.3 BC
Acelepryn 6 fl oz	1.8 DE	3.2 C	4.8 A	4.8 A	3.2 B	0.6 D	1.1 E	0.5 D	1.0 E	4.7 DE
Zeta-cypermethrin (Sevin®) 4 fl oz	0.8 E	0.4 D	2.0 B	1.0 C	0.0 C	0.0 D	0.1 E	0.1 D	0.0 E	0.1 E
Untreated	5.0 A	5.0 A	4.8 A	4.0 AB	4.4 A	25.4 A	32.1 A	17.2 A	27.5 A	13.8 C

Comparative Efficacy on Viburnum Leaf Beetle (Pyrrhalta viburni)

Viburnum leaf beetle is native to Europe and Asia and was first detected in North America in 1947 in Ontario, Canada. Since 1978 when breeding populations were discovered in the Ottawa/Hull region of Canada, viburnum leaf beetle has slowly spread south and was found in Maine in 1994 and in New York in 1996. Currently, it has been found as far south as Pennsylvania and Ohio. Viburnum leaf beetle feeds exclusively on viburnum species. The most susceptible include arrowwood viburnum (*V. dentatum*), European cranberry bush viburnum (*Viburnum opulus*), Rafinesque viburnum (*V. rafinesquianum*), and Sargent viburnum (*V. sargentii*). It will also feed on wayfaring tree viburnum (*V. lantana*), nannyberry viburnum (*V. lentago*), blackhaw viburnum (*V. prunifolium*) and several other species (http://creatures.ifas.ufl.edu/orn/beetles/viburnum_leaf_beetle.htm).

In the series of experiments presented here, most tested products reduced viburnum leaf beetle populations and feeding damage. The best consistently performing products in these three experiments were BAS 320i, Celero, and Safari. See the individual reports below for more information.

Costa 2006

In this experiment, Costa examined 5 products for efficacy of viburnum leaf beetle larvae. Treatments were applied on May 16 to runoff using a handheld, pump sprayer with a second TriStar treatment made on Jun 13. The extent (percentage of affected leaves) and severity (percentage area affected on damaged leaves) of leaf feeding by larvae were assessed at 2, 7, 14, and 28 DAT by visually inspecting each plant and rating damage on a scale ranging from 1 – 10.

By 2 DAT, no live larvae were detected outside of plants in the water control. By 14 DAT all insecticide treatments had significantly less feeding damage than the water-treated control (Table 26, Table 27). None of the treatments varied significantly from each other. The second application of Tristar was made according an industry-prescribed protocol and was not necessary for larval control. All chemical insecticide treatments provided effective control of VLB larvae.

No phytotoxicity was observed.

Costa 2007

In this experiment, Costa examined 8 products for efficacy of viburnum leaf beetle larvae. All products except Safari 2G were applied as foliar sprays. Safari 2G was broadcast by hand around the plant base.

By 7 DAT, viburnum plants treated with Acelepryn, BAS 320i, Celero, Permethrin, Safari, and Tolfenpyrad exhibited significantly less feeding damage than the Nontreated plants (Table 28). Throughout this experiment Met 52 was equivalent to the Nontreated. Safari 2G did reduce feeding damage at 14 DAT. While most products reduced severity of feeding, the extent of defoliation was only reduced by BAS 320i and Safari through 14 DAT (Table 29). By 28 DAT, these two products plus Celero, Permethrin and Tolfenpyrad reduced defoliation.

Table 26. Efficacy of several insecticides for Viburnum Leaf Beetle (*Pyrrhalta viburni*) larval management on Arrowwood viburnum (*Viburnum dentatum*) – Defoliation Severity, Costa, 2006.

Treatment	Rate Per 100 Gal	Defoliation Severity Rating Relative Area ^z			
		2 DAT	7 DAT	14 DAT ^y	28 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	16 fl oz	2.2	3.0	3.8*	2.8*
Celero 16WSG (clothianidin)	4 oz	2.6	3.4	3.4*	3.0*
Permethrin 2.5EC	128 fl oz	2.4	3.0	3.8*	3.2*
Safari 20SG (dinotefuran)	8 oz	2.8	3.4	3.8*	3.4*
TriStar 70WSP (acetamiprid) + Cohere	96 g + 0.125%	2.8	3.4	4.2*	3.4*
Nontreated		2.8	5.6	6.2	7.0

^zThe severity of larval feeding post treatment as determined by qualitative rating of relative area affected on damaged leaves. Scale 1-10 is for 0 to 100% (1=0, 2=1-5, 3=6-15, 4=16-30, 5=31-50, 6=51-70, 7=71-85, 8=86-95, 9=96-99, 10=100% affected).

^yAn ‘*’ indicates a significant difference between insecticide treatments and the water treated control (alpha = 0.05; one sided Dunnett’s after GLM). There were no significant differences among insecticide treatments (P>0.05, GLM-ANOVA).

Table 27. Efficacy of several insecticides for Viburnum Leaf Beetle (*Pyrrhalta viburni*) larval management on Arrowwood viburnum (*Viburnum dentatum*) – Defoliation Extent, Costa, 2006.

Treatment	Rate Per 100 Gal	Defoliation Extent Rating ^z			
		2 DAT	7 DAT	14 DAT ^y	28 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	16 fl oz	2.8	3.2	4.2*	2.6*
Celero 16WSG (clothianidin)	4 oz	3.0	3.4	3.8*	2.8*
Permethrin 2.5EC	128 fl oz	3.0	3.8	4.4*	3.2*
Safari 20SG (dinotefuran)	8 oz	3.0	3.6	4.2*	3.2*
TriStar 70WSP (acetamiprid) + Cohere	96 g + 0.125%	3.2	3.8	5.0*	3.4*
Nontreated		3.6	4.4	6.6	7.0

^zThe

extent of larval feeding post treatment as determined by qualitative rating of percentage of affected leaves. Scale 1-10 is for 0 to 100% (1=0, 2=1-5, 3=6-15, 4=16-30, 5=31-50, 6=51-70, 7=71-85, 8=86-95, 9=96-99, 10=100% affected).

^yAn ‘*’ indicates a significant difference between insecticide treatments and the water treated control (alpha = 0.05; one sided Dunnett’s after GLM). There were no significant differences among insecticide treatments (P>0.05, GLM-ANOVA).

Table 28. Efficacy of several insecticides for Viburnum Leaf Beetle (*Pyrrhalta viburni*) larval management on Arrowwood viburnum (*Viburnum dentatum*) – Defoliation Severity, Costa, 2007.

Treatment	Rate Per 100 Gal	Defoliation Severity Rating (±SE) Relative Area ^z			
		Pre-Trt ^y	Week 1 ^x	Week 2	Week 4
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz	5.2 (0.2)	4.6 (0.2)*	4.6 (0.2)*	4.4 (0.2)*
BAS 320i (metaflumizone)	16 oz	3.6 (0.4)	4.2 (0.5)*	4.4 (0.2)*	4.2 (0.2)*
Celero 16WSG (clothianidin)	4 oz	4.2 (0.4)	4.2 (0.2)*	4.4 (0.2)*	4.4 (0.2)*
<i>Metarhizium anisopliae</i> (Strain F52)	29 oz	4.6 (0.2)	6.6 (0.5)	6.4 (0.5)	5.8 (0.4)
Permethrin 2.5EC	128 fl oz	4.0 (0.5)	4.2 (0.2)*	4.4 (0.2)*	4.0 (0.3)*
Safari 2G (dinotefuran)	2.2 g/gal potting media	4.8 (0.6)	5.5 (0.9)	5.0 (0.4)*	5.0 (0.6)
Safari 20SG (dinotefuran)	8 oz	4.0 (0.0)	4.5 (0.3)*	4.5 (0.3)*	4.3 (0.5)*
Tolfenpyrad EC	21 fl oz	3.5 (0.3)	4.0 (0.0)*	4.3 (0.3)*	4.5 (0.5)
Nontreated		4.8 (0.6)	6.2 (0.0)	6.4 (0.5)	5.8 (0.5)

^z The severity of larval feeding post treatment as determined by qualitative rating of relative area affected on damaged leaves. Scale 1-10 is for 0 to 100% (1=0, 2=1-5, 3=6-15, 4=16-30, 5=31-50, 6=51-70, 7=71-85, 8=86-95, 9=96-99, 10=100% affected).

^y Pre-treatment (Pre-Trt) ratings were taken the day applications were made.

^x An ‘*’ indicates a significant difference between insecticide treatments and the water treated control (alpha = 0.05; one sided Dunnett’s after GLM). There were no significant differences among insecticide treatments (P>0.05, GLM-ANOVA).

Table 29. Efficacy of several insecticides for Viburnum Leaf Beetle (*Pyrrhalta viburni*) larval management on Arrowwood viburnum (*Viburnum dentatum*) – Defoliation Extent, Costa, 2007.

Treatment	Rate Per 100 Gal	Defoliation Extent Rating (\pm SE) ^z				^z The extent of larval
		Pre-Trt ^y	Week 1 ^x	Week 2	Week 4	
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz	5.8 (0.2)	7.0 (0.6)	6.6 (0.6)	5.8 (0.6)	
BAS 320i (metaflumizone)	16 oz	5.0 (0.6)	4.8 (0.7)*	4.8 (0.4)*	4.2 (0.4)*	
Celero 16WSG (clothianidin)	4 oz	5.0 (0.5)	5.4 (0.7)	6.0 (0.7)	4.8 (0.6)*	
<i>Metarhizium anisopliae</i> (Strain F52)	29 oz	5.4 (0.2)	7.2 (0.6)	7.4 (0.2)	7.2 (0.4)	
Permethrin	128 fl oz	5.4 (0.5)	5.2 (0.7)	5.6 (0.7)	4.6 (0.4)*	
Safari 2G (dinotefuran)	2.2 g/gal potting media	5.5 (0.7)	6.3 (1.4)	6.0 (0.9)	5.8 (0.8)	
Safari 20SG (dinotefuran)	8 oz	4.8 (0.5)	5.0 (0.7)*	4.8 (0.9)*	4.3 (0.8)*	
Tolfenpyrad EC	21 fl oz	4.5 (0.3)	4.5 (0.7)	5.3 (0.9)	4.5 (0.7)*	
Nontreated		6.0 (0.6)	7.4 (1.1)	7.6 (0.8)	7.0 (0.8)	

feeding post treatment as determined by qualitative rating of percentage of affected leaves. Scale 1-10 is for 0 to 100% (1=0, 2=1-5, 3=6-15, 4=16-30, 5=31-50, 6=51-70, 7=71-85, 8=86-95, 9=96-99, 10=100% affected).

^y Pre-treatment (Pre-Trt) ratings were taken the day applications were made.

^x An ‘*’ indicates a significant difference between insecticide treatments and the water treated control ($\alpha = 0.05$; one sided Dunnett’s after GLM). There were no significant differences among insecticide treatments ($P > 0.05$, GLM-ANOVA).

Holmes 2007

During 2007, Holmes generated viburnum leaf beetle efficacy data for chlorantraniliprole and thia-methoxam. Several rate range experiments were conducted with Actara (the Canadian trade name for Flagship 25WG) and Rynoxapyr (the food use trade name for Acelepryn). The trade names within the US are used to summarize the information and in the data tables (Table 30- Table 33).

Four experiments were conducted simultaneously; two each for each product. Prior to application, each plant was assessed for a minimum level of hatched larvae. Crop tolerance and efficacy ratings were done at 7, 12, 35 and 61 days after treatment on June 13, 2007. In the first experiment with Flagship (Table 30), all of the treatments had significantly less foliar damage than the Nontreated check through 35 DAT. At the final rating date, Conserve-treated plants exhibited the same amount of foliar feeding damage as the Nontreated plants, while all rates of Flagship were significantly lower than the Nontreated but not Conserve. In other words, control was beginning to break approximately 2 months after treatment. However in the second experiment (Table 31), Conserve and all three rates of Flagship provided excellent control through 2 months, albeit with a slightly lower infestation pressure.

In both experiments testing Acelepryn (Table 32 and Table 33), Acelepryn at all three tested rates provided great efficacy, equivalent to Conserve through 2 months.

None of the treatments resulted in any phytotoxicity symptoms.

Table 30. Foliar damage due to Viburnum Leaf Beetle on *Viburnum trilobum* treated with thiamethoxam (Actara 25WG) Experiment 1, Holmes, 2007 a

Treatment	Rate	Foliar Damage (Percent)			
		7 DAT	12 DAT	35 DAT	61 DAT
Conserve	50 ml /1000L	5.8 b	7.5 b	5.0 b	10.0 ab
Flagship (Actara) 25 WG (thiamethoxam)	0.28 kg/ha	6.3 b	6.3 b	21.3 a	8.0 b
	0.56 kg/ha	6.3 b	5.5 b	5.0 b	5.5 b
	1.12 kg/ha	4.5 b	5.0 b	5.0 b	4.3 b
Nontreated		16.3 a	18.8 a	23.8 a	21.3 a

Initial application occurred on June 13, 2007.

Means followed by same letter do not significantly differ ($p = 0.05$, Tukey.s HSD).

Table 31. Foliar damage due to Viburnum Leaf Beetle on *Viburnum trilobum* treated with thiamethoxam (Actara 25WG) Experiment 2, Holmes, 2007 b

Treatment	Rate	Foliar Damage (Percent)			
		7 DAT	12 DAT	35 DAT	61 DAT
Conserve	50 ml /1000L	1.5 b	1.3 b	2.8 b	0.5 b
Flagship (Actara) 25 WG (thiamethoxam)	0.28 kg/ha	1.0 b	1.3 b	2.0 b	0.5 b
	0.56 kg/ha	2.0 b	1.5 b	2.8 b	0.5 b
	1.12 kg/ha	1.0 b	1.0 b	2.0 b	0.5 b
Nontreated		8.8 a	10.0 a	12.5 a	6.8 a

Initial application occurred on June 13, 2007.

Means followed by same letter do not significantly differ (p = 0.05, Tukey's HSD).

Table 32. Foliar damage due to Viburnum Leaf Beetle on *Viburnum trilobum* treated with Rynaxapyr (DPX 2EY45 20SC) Experiment 1, Holmes, 2007 a

Treatment	Rate	Foliar Damage (Percent)			
		8 DAT	13 DAT	36 DAT	62 DAT
Conserve	50 ml /1000L	5.8 b	5.0 b	5.0 b	2.8 b
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.28 kg/ha	7.0 b	7.5 b	8.8 b	8.8 b
	0.56 kg/ha	5.3 b	5.5 b	5.0 b	2.0 b
	1.12 kg/ha	4.5 b	5.0 b	5.0 b	3.5 b
Nontreated		21.3 a	35.0 a	83.8 a	22.5 a

Initial application occurred on June 13, 2007.

Means followed by same letter do not significantly differ (p = 0.05, Tukey's HSD).

Table 33. Foliar damage due to Viburnum Leaf Beetle on *Viburnum trilobum* treated with Rynaxapyr (DPX 2EY45 20SC) Experiment 2, Holmes, 2007 b

Treatment	Rate	Foliar Damage (Percent)			
		8 DAT	13 DAT	36 DAT	62 DAT
Conserve	50 ml /1000L	1.5 b	1.3 b	1.5 b	3.5 b
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.28 kg/ha	1.3 b	1.3 b	2.8 b	8.0 b
	0.56 kg/ha	1.3 b	1.5 b	2.3 b	2.8 b
	1.12 kg/ha	1.3 b	1.3 b	1.3 b	1.8 b
Nontreated		3.5 a	8.8 a	21.3 a	28.8 a

Initial application occurred on June 13, 2007.

Means followed by same letter do not significantly differ (p = 0.05, Tukey's HSD).

Isaacson 2007

To generate efficacy data for chlorantraniliprole and thiamethoxam to manage viburnum leaf beetle Isaacson ran several rate range experiments with Actara (the Canadian trade name for Flagship 25WG) and Rynoxapyr (the food use trade name for Acelepryn). The trade names within the US are used to summarize the information and in the data tables (

Table 34 - Table 37).

All three rates for Flagship provided good to excellent control throughout both experiments, equivalent to Conserve SC (

Table 34 and Table 35). All three rates of Acelepryn also provided good to excellent efficacy throughout the experiments. (Table 36 and Table 37).

No phytotoxicity was observed.

**Table 34 Summary of Damage ratings (0 – 10) for *Viburnum opulus nanum* infested with VLB, treated with thiamethoxam (Actara 25WG)
Experiment 1, Isaacson, 2007 a**

Treatment	Rate	May 22	May 29	Jun 5	Jun 12	Jun 20	Jun 27	July 4	July 25	Aug 15	Aug 29
Flagship (Actara) 25 WG (thiamethoxam)	0.28 kg/ha	0.58	0.58	0.58	0.58	0.58	0.58	0.50	0.00	0.25	0.58
	0.56 kg/ha	0.42	0.42	0.42	0.42	0.42	0.25	0.08	0.00	0.00	0.20
	1.12 kg/ha	0.42	0.42	0.42	0.42	0.42	0.33	0.25	0.00	0.00	0.17
Conserve SC (spinosad)		0.75	0.75	0.75	0.75	0.75	0.75	0.33	0.00	0.00	0.33
Nontreated		0.67	2.17	2.58	2.58	2.58	2.58	2.17	1.83	2.42	3.25

**Table 35 Summary of Damage ratings (0 – 10) for *Viburnum opulus nanum* infested with VLB, treated with thiamethoxam (Actara 25WG)
Experiment 2, Isaacson, 2007 b**

Treatment	Rate	May 22	May 29	Jun 5	Jun 12	Jun 20	Jun 27	July 4	July 25	Aug 15	Aug 29
Flagship (Actara) 25 WG (thiamethoxam)	0.28 kg/ha	0.50	0.50	0.50	0.50	0.50	0.50	0.42	0.00	0.25	0.50
	0.56 kg/ha	0.67	0.67	0.67	0.67	0.67	0.42	0.08	0.00	0.00	0.25
	1.12 kg/ha	0.67	0.67	0.67	0.67	0.67	0.50	0.08	0.00	0.00	0.17
Conserve SC (spinosad)		0.67	0.67	0.67	0.67	0.67	0.58	0.25	0.00	0.00	0.33
Nontreated		0.67	2.17	2.58	2.58	2.58	2.58	2.17	1.83	2.42	3.25

**Table 36 Summary of Damage ratings (0 – 10) for *Viburnum opulus nanum* infested with VLB, treated with Rynaxapyr (DPX 2EY45 20SC)
Experiment 1, Isaacson, 2007 a**

Treatment	Rate	May 22	May 29	Jun 5	Jun 12	Jun 20	Jun 27	July 4	July 25	Aug 15	Aug 29
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	31.25 ml/L	0.58	0.58	0.58	0.58	0.58	0.50	0.25	0.00	0.08	0.17
	62.5 ml/L	0.50	0.50	0.50	0.50	0.50	0.50	0.17	0.00	0.08	0.33
	125 ml/L	0.67	0.67	0.67	0.67	0.67	0.67	0.17	0.00	0.00	0.50
Conserve SC (spinosad)		0.75	0.75	0.75	0.75	0.75	0.75	0.33	0.00	0.00	0.33
Nontreated		0.67	2.17	2.58	2.58	2.58	2.58	2.17	1.83	2.42	3.25

**Table 37 Summary of Damage ratings (0 – 10) for *Viburnum opulus nanum* infested with VLB, treated with Rynaxapyr (DPX 2EY45 20SC)
Experiment 2, Isaacson, 2007 b**

Treatment	Rate	May 22	May 29	Jun 5	Jun 12	Jun 20	Jun 27	July 4	July 25	Aug 15	Aug 29
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	31.25 ml/L	0.75	0.75	0.75	0.75	0.75	0.50	0.17	0.00	0.00	0.25
	62.5 ml/L	0.58	0.58	0.58	0.58	0.58	0.50	0.08	0.00	0.00	0.25
	125 ml/L	0.50	0.50	0.50	0.50	0.50	0.42	0.08	0.00	0.00	0.17
Conserve SC (spinosad)		0.67	0.67	0.67	0.67	0.67	0.58	0.25	0.00	0.00	0.33
Nontreated		0.58	1.83	2.83	2.83	2.83	2.67	2.08	1.58	2.00	3.17

Weston 2007

In this experiment, Weston tested seven products for their efficacy on viburnum leaf beetle infesting established arrowwood viburnum (*Viburnum dentatum*) in field plots at the Bluegrass Lane Turf and Ornamentals Research Farm in Ithaca, NY. The shrubs, which had been growing under field conditions for 7 years, were approximately 6' tall and were naturally infested by viburnum leaf beetle in previous years. Products were applied as foliar sprays on May 22, 2007 when viburnum leaf beetle was in its first larval instar (egg hatch had begun on May 9). Five plants (replicates) were used for each treatment, and larval feeding damage was assessed 1 and 2 weeks after treatment. Data were analyzed with randomized complete block ANOVA, and treatments were compared with LSD.

The range of feeding damage was dramatic, ranging from 55% on the Nontreated control to near zero for the most effective treatments (Table 38). *Metarhizium anisopliae*, a fungus effective against many immature insects, had no effect on larvae (defoliation was virtually identical to that of the Nontreated control). The remaining products provided good to excellent control. Most effective were Celero and Safari, which were slightly more efficacious than Acelepryn, BAS 320i, Merit and Tolfenpyrad through 14 DAT.

[NOTE: Earlier field trials by Weston have shown that soil drenches with Merit 75 WP have resulted in nearly complete protection from viburnum leaf beetle for several years. In the current trial, Merit was applied as a foliar spray, like all of the other test products.]

Table 38. Efficacy of several insecticides for Viburnum Leaf Beetle (*Pyrrhalta viburni*) management on Arrowwood viburnum (*Viburnum dentatum*), Weston, 2007.

Treatment	Rate Per 100 Gal	Defoliation	
		1 WAT	2 WAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz	3.3 ab	4.3 b
BAS 320i (metaflumizone)	16 oz	2.5 b	2.2 bc
Celero (clothianidin)	4 oz	0.7 b	0.7 d
Merit (imidacloprid)	10 tsp	3.6 ab	3.6 b
<i>Metarhizium anisopliae</i> (Strain F52)	29 oz	22.2 ab	50.2 a
Safari 20SG (dinotefuran)	8 oz	2.7 b	1.7 cd
Tolfenpyrad EC	21 oz	4.4 ab	3.9 b
Nontreated		24.4 a	55.4 a

Comparative Efficacy on Red headed Flea Beetle (Systema frontalis)

Redheaded flea beetle (also called cranberry flea beetle), *Systema frontalis*, has become a serious pest of nursery stock and has been considered the most damaging flea beetle in container nurseries in recent years. It has a wide range of ornamental hosts including Itea, hydrangea, forsythia, roses, holly, azalea, hibiscus, asters, chrysanthemum and zinnia, where the adults typically chew leaves causing small holes and skeletonized leaves. They also feed on growing tips causing deeply notched leaves.

In the series of 10 experiments presented here, most tested products reduced redheaded flea beetle populations and feeding damage. The best consistently performing products in these experiments were Aloft, IKI-3106 and Tristar. See the individual reports below for more information.

Braman 2012 and 2013

In 2012 and 2013, Braman compared several products applied foliar for ability to protect nursery grown *Itea* and *Hydrangea* from chewing damage by red headed flea beetle. In 2012, all treatments were applied on May 11; a second application was made for MBI-203 on May 18. Plants treated at the nursery were evaluated on-site for foliar damage at 1, 2, 4, and 11 weeks-post application. In both years, detached leaves from treated plants were bagged and returned to the laboratory and evaluated for beetle survival and % leaf damage (chewing injury) at various times after treatment application in a petri dish exposure trial. Data were subjected to ANOVA using the GLM procedure in SAS and mean separation was accomplished using the LSD procedure.

In the detached-leaf study in 2012, Aloft and Flagship consistently reduced beetle survival and reduced damage in both *Itea* and *hydrangea* (Table 39, Table 40). In the nursery, Marathon provided the most consistent damage reduction from 1 to 11 weeks post application on both *Itea* and *hydrangea*. At 1 week post application, Hachi-Hachi also reduced damage on *Itea*; on *Hydrangea*, Marathon, Hachi-Hachi, Aloft and Safari provided the greatest protection. At 11 weeks post application, all products showed a reduction in damage on *Itea*, with the least damage observed on plants treated with Marathon, Aloft, Safari, Hachi-Hachi and MBI-203; on *hydrangea*, only Marathon treated plants displayed less damage than the Nontreated.

In the 2013 detached leaf study, Only Aloft provided a significant reduction in beetle survival relative to the UTC at 1 and 10 days after treatment (Table 41). Beetle survival at 21, 24 and 30 DAT did not vary significantly with treatment. Leaf injury, however, was affected on all sample dates, suggesting antifeedant behavior in response to residual application even when mortality was not inflicted by treatment. Damage was reduced at 1 DAT by Aloft, Safari, GF-2860, MBI 203, Onyx and Discus. Damage was reduced at 10 DAT by Aloft, Safari- foliar and drench, GF-2860, MBI 203, Onyx and Discus. At 21 DAT, feeding was significantly reduced by the Safari drench, GF-2860, A20520, Aloft and Discus. Subsequent exposure by that same set of beetles resulted in continued suppression of damage on leaves from plants treated with Safari (drench), Aloft and Discus.

Frank 2013

In 2013, Frank compared efficacy of several products applied foliar to protect nursery grown Virginia sweetspire from chewing damage by cranberry flea beetle, *Systema frontalis*. All products were sprayed once on Jul 10 except Discus which was sprayed on Jul 10 and 24, and MBI-203, sprayed on Jul 10, 17, 24 and 31. All applications were made at least 4 hours prior to rain; however, it rained nearly every day during the experiment. Flea beetle abundance was not a good assessment of flea beetle damage because flea beetles were jumping from plants as observers approached plots. Thus, no significant differences were detected in number of flea beetles per plant (Table 42); however, assessment of shot-hole leaf damage showed that all products significantly reduced feeding injury.

Kunkel 2014

In this experiment, Kunkel compared several products (GF-2860, Hachi-Hachi, Mainspring, MBI-203, Safari and Scimitar) for ability to protect Virginia sweetspire (*Itea virginica*) sage (*Salvia nemorosa*), and stonecrop (*Sedum telephium*) from chewing damage by red headed flea beetle (Table). All products were applied as foliar sprays,

except Safari applied as drench in one treatment. Treatments were applied on Jun 16, and all treatments, except Hachi-Hachi and Safari, reapplied on Jul 7. A minimum of six (caged) or eight replicates (natural) were requested for this trial; however, nurseries did not have enough plants of one species or cultivar available to use in the experiments. The trial had three replicates of *Salvia* and two replicates of *Itea* and *Sedum* placed side-by-side in the greenhouse. Previous work with redheaded flea beetles found all three species to be suitable hosts. Adult feeding damage was assessed at various times after treatment.

The three different species of plants resulted in a complicated analysis because there was a three-way interaction between products, plants, and amount of damage (Table 43). Scimitar and Safari provided the most consistent control across the duration of the experiment. Adult *S. frontalis* prefer to feed on *Itea* and *Salvia* when compared to *Sedum*, and they fed on either *Itea* or *Salvia* similarly most of the observation periods. Additional research is needed to develop management tactics for both larva and adult *S. frontalis*.

In summary, Scimitar, Safari and most frequently the higher rate of GF-2860 significantly reduced redheaded flea beetle feeding damage on these ornamental plants tested; Mainspring and MBI-203 also provided some damage reduction. *Sedum* frequently had significantly less damage than either *Itea* or *Salvia*. The experiment found greater than acceptable foliage damage occur from 14 to 21 DAT after initial treatment. Although populations were low preceding 14 DAT, this beetle exhibited the capability of increasing populations rapidly. This increase in population could be from within the nursery crops or potentially peripheral areas since it has a wide host range that includes many weed species. This aspect of the pest's biology warrants further investigation.

No phytotoxicity and no significant differences in plant height and width between treatments were observed.

Cloyd 2016

In 2016, Cloyd compared efficacy of several products applied foliar to protect container grown Virginia sweetspire from chewing damage by cranberry flea beetle. All products were sprayed on Jun 23, Jul 1, 8, 15, 22, Aug 12 and Sep 2. By the end of the sampling season the insecticides BeetleGONE® and Hachi-Hachi® appeared to limit the frequency of foliar damage caused by redheaded flea beetle adults when compared to the Nontreated check; however, the insecticides Preferal®, Safari®, and TriStar® never exceeded 1 to 10% damage (Graph 44). Redheaded flea beetle adults were observed throughout the sampling period; however, there was a noticeable peak of activity in October. In addition, redheaded flea beetle adults were abundant in mid-to late-October with a noticeable increase in feeding damage on the test plants. Therefore, applications of the insecticides should have been continued into October.

Gilrein 2016

In 2016, Gilrein compared efficacy of several products applied foliar to protect container grown hydrangea from chewing damage by redheaded flea beetle (RFB). All products were sprayed on Jul 2; BeetleGONE, and XXpire treatments were repeated on Jul 9. Approximately ten RFB were introduced to screen cages on two terminals per plant just prior to first treatment application. A single application of Hachi-Hachi, IKI-3106, XXpire, and Scimitar provided the most effective control of adult RFB (Table 45). BeetleGONE and Preferal applications provided very little control and efficacy of these products were not statistically different from water-treated plants. Foliar feeding damage by RFB was significantly higher on plants treated with BeetleGONE and Preferal and not statistically different from damage on control plants. No noticeable insecticide residue, injury or phytotoxicity on plants was associated with any treatment.

Chong 2017

In 2017, Chong compared efficacy of several products applied foliar to protect container grown hydrangea from a natural infestation of redheaded flea beetle. Fungicides were sprayed at various times from Jul 26 to Aug 11. No treatments provided consistent reduction in the adult flea beetles densities throughout the study (Table 46). Results suggested that, although some products such as Scimitar was effectively in killing adult flea beetles at the time of the application, no treatment provided sufficient repellency or residue toxicity to reduce the numbers of adult flea beetles on the hydrangea plants for more than 2 days after application and throughout the experiment. As a result, the % defoliation and crop quality worsened and were not different among the insecticide-treated and

water-treated plants. This study has illustrated the difficulty of managing the redheaded flea beetles. No product tested in this study provide quick knockdown of the beetles and provide sufficient repellency or residual control. No phytotoxicity was associated with any treatment.

Kunkel 2017

In 2017, Kunkel compared efficacy of several products applied foliar to protect container grown forsythia from chewing damage by red headed flea beetle (RHFB). Treatments were applied as foliar sprays from May 31 to Jun 14 on a schedule shown in Table 47 and Table 48. Product efficacy was evaluated multiple ways to determine if products were able to manage flea beetle populations and their damage. There were significantly more live adult beetles found on plants treated with Hachi-Hachi, IKI, and Preferal during the Jun 14 evaluation period. Other time periods found no differences in the number of live beetles counted on plants. There was no significant differences in the number of damaged leaves or branches with damaged leaves due to treatments throughout the experiment; however, there was significant differences on damaged new growth. Upstar-Gold, Preferal, BeetleGONE, VST combined with BeetleGONE, and Tristar all reduced flea beetle feeding on new growth at some point during the experiment. Upstar-Gold provided the most consistent protection of new foliage followed by Preferal, BeetleGONE, Tristar, and the VST combo product. Upstar-Gold, Tristar, BeetleGONE, and the VST combination treatment with significantly lower damage ratings compared to the Nontreated control forsythia plants. Damaged leaves removed from treated plants were brought into the lab and the amount of leaf area damaged from flea beetle feeding was evaluated using the program Image-J. Treatments did not significantly reduce feeding damage when foliage was evaluated in this manner. No phytotoxicity or growth reduction was associated with any treatment.

Frank 2017

In 2017, Frank conducted laboratory assays to compare efficacy of several products applied foliar to protect container grown hydrangea from chewing damage by adult red headed flea beetle (RHFB). Treatments were applied on Jun 13 and 20. Beetles collected from a local nursery were placed in petri dishes with leaf disks cut from treated plants 1, 7, and 14 days after initial treatment. Each assay lasted 5 days with percent damage and beetle mortality measured on days 1, 3, 5. Throughout the experiment leaves treated with the standard imidacloprid incurred the least damage from beetles. One DAT Hachi-Hachi and IKI-3106 had similarly low damage as imidacloprid treated leaves. Seven DAT leaves treated with imidacloprid and those treated with Hachi-Hachi were damaged least by beetles. After one day of feeding on 14 DAT leaves those treated with imidacloprid, preferal, or VST 06330 had the least damage (Table 49). After 3 and 5 days of feeding on these leaves there were no significant difference among treatments. Beetle mortality was highest 1 DAT in petri dishes with leaves treated with imidacloprid. Nearly all beetles in this treatment were dead after 5 days of exposure. Mortality was low in other treatments. There were no significant differences in mortality after exposure to leaves 7 and 14 DAT. At the end of the experiment, damage to 5 leaves per plant outdoors that were marked prior to insecticide applications was low but highest on Nontreated plants and imidacloprid treated plants.

Gilrein 2018

In 2018, Gilrein compared efficacy of several products applied foliar to protect container grown hydrangea from chewing damage by redheaded flea beetle (RFB). Treatments were applied from Jul 3 to Jul 18 on a schedule shown in Table 50. Hachi-Hachi and Pyridium (applied once) and IKI- 3106 and TriStar (applied twice) provided very effective control of adult RFB. Both rates of V- 10433 provided moderate control, while Ancora provided very little control. Foliar feeding damage by RFB correlated well with treatment efficacy. No noticeable insecticide residue or phytotoxicity on plants was associated with any treatment.

Frank 2019

During 2019, Frank studied 9 new products for their impact on redheaded flea beetle (RFB) populations and feeding damage on hydrangea. Foliar applications began Jul 5 and ended Jul 30. Adult beetles were counted by visual scan for 10 seconds followed by 20 seconds of combing through foliage. Population counts on nontreated control plants were very low. BCS-507, Hachi Hachi, Sarissa, and Tristar 8.5SL provided very effective control of adult RFB at 3 and 7 DAIT (Table 51). At 3 DAIT, Ancora, TetraCURB Concentrate, and V-10433 statistically reduced populations. Very little feeding injury was observed, and treatments were not statistically different from

each other (data not shown). There was no noticeable insecticide residue or phytotoxicity associated with any treatment and no differences in plant growth observed among treatments.

Gilrein 2019

In 2019, Gilrein compared efficacy of several products applied foliar to protect container grown panicle hydrangea from chewing damage by redheaded flea beetle (RFB). Treatments were applied from Jul 2 to Jul 17 on a schedule shown in Table 50. BCS-507, IKI-3106, KOC22018-8, Mainspring GNL, TetraCURB Concentrate, V-10433, and TriStar 8.5SL provided very effective control of adult RFB for the duration of the trial (Table 52). Foliar feeding damage by RFB generally correlated well with treatment efficacy; however, plants treated with TetraCURB Concentrate and V-10433 had foliar damage ratings lower than but not significantly different from control plants despite the high beetle mortality associated with those treatments over the course of the trial. There was no noticeable insecticide residue or phytotoxicity associated with any treatment and no differences in plant growth observed among treatments (data not presented).

Gilrein 2020

During 2020, Gilrein compared efficacy of several products applied foliar to protect container grown panicle hydrangea from chewing damage by redheaded flea beetle (RFB). Adult beetles were introduced to tulle sleeve cages on hydrangea terminals on July 1. Treatments were applied from July 1 to Jul 16 with some treatments applied once or up to 5 times based on the protocol. ISM-555 and TriStar 8.5SL provided very effective, quick acting control of adult RFB and lasted for the duration of the trial (Table 53) as assessed by adult beetle mortality and feeding damage. By 2WAT, BW133 and BW238ES exhibited greater than 90% population reduction and significantly less feeding damage than the water sprayed controls. MBI-203, MBI-306, and SP3014 did not lessen populations or significantly reduce feeding damage. Minor marginal necrosis occurred 14 days after the first application with BW133 and BW238ES.

Sadof 2021

In 2021, Sadof compared efficacy of several products applied foliar to protect container grown panicle hydrangea from chewing damage by redheaded flea beetle. Foliar applications occurred from May 25 through July 14. Only the ISM 555 product at 3.84 and 5.76 fl oz provided significantly lower numbers of chewed leaves over the course of the study based on a Tukey's HSD test ($P < 0.05$ and 0.001) respectively (Table 54). Sarissa, Pradia and Xxpire provide acceptable levels of control until 28 Jun (37 Days after the first treatment) but were not statistically different. Similar results found for the subjective damage evaluation with both rates of ISM-555 being < 2 (10% damaged leaves) over the course of the study (data not shown). Numbers of beetles present on the last day of evaluation followed the same pattern.

Gilrein 2021

In 2021, Gilrein compared efficacy of several products applied foliar to protect container grown panicle hydrangea from chewing damage by redheaded flea beetle (RFB). Treatments were applied from June 29 to Jul 13 with some treatments applied once or up to 5 times based on the protocol. ISM-555 and TriStar 8.5SL provided very effective control of adult RFB for the duration of the trial (Table 55) as assessed by adult beetle mortality and feeding damage. By 5WAT, SP3014 at 26 fl oz and Velifer suppressed populations, but feeding damage was not significantly different from the nontreated control. There was no noticeable insecticide residue or phytotoxicity associated with any treatment.

Table 39. Detached Leaf Study for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Virginia Sweetspire (*Itea virginica*) 'Henry's Garnet' and Hydrangea (*Hydrangea* sp.) 'White Diamonds', Braman, 2012.

Treatment	Rate Per 100 Gal	No. of Living Beetles ^x		% Damage Itea		% Damage Hydrangea	
		Day 1*	Day 5*	Day 1*	Day 5*	Day 1*	Day 5*
Aloft (clothianidin/bifenthrin)	14.9 fl oz	0.2 c	0 b	0.4 c	0 c	0 c	0 c
Flagship 25WG (thiamethoxam)	8 fl oz	0.4 bc	0 b	1.0 bc	0 c	0 c	1.0 c
Hachi-Hachi (tolfenpyrad)	21 fl oz	1.0 a	1.0 a	7.0 abc	14.0 abc	9.0 a	15.0 ab
Hachi-Hachi (tolfenpyrad)	32 fl oz	0.8 ab	0.8 a	5.4 abc	4.0 bc	1.0 bc	13.0 abc
Marathon G (imidacloprid)	7 g/pot	0.8 ab	0.6 a	2.8 bc	2.0 bc	1.6 abc	7.4 abc
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	1 lb	1.0 a	1.0 a	11.0 a	15.0 ab	5.0 abc	20.0 a
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	2 lb	0.8 ab	0.8 a	3.0 bc	16.0 ab	8.0 ab	12.0 abc
Onyx (bifenthrin)	12.8 fl oz	1.0 a	1.0 a	3.0 bc	12.0 abc	7.0 abc	12.0 abc
Safari 20SG (dinotefuran)	8 oz	0.8 ab	0.6 a	5.4 abc	3.0 bc	1.0 bc	15.0 abc
TriStar 30SG (acetamiprid)	8 g	0.8 ab	0.6 a	3.6 bc	5.6 bc	3.0 abc	3.4 bc
Nontreated		1.0 a	1.0 a	8.0 ab	23.0 a	6.0 abc	18.0 a

Table 40. Damage Rating for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Virginia Sweetspire (*Itea virginica*) 'Henry's Garnet' and Hydrangea (*Hydrangea* sp.) 'White Diamonds', Braman, 2012.

Treatment	Rate Per 100 Gal	Itea				Hydrangea			
		1 WAT	2 WAT	4 WAT	11 WAT	1 WAT	2 WAT	4 WAT	11 WAT
Aloft (clothianidin/bifenthrin)	14.9 fl oz	5.6 a ^x	0 a	0.4 bc	0.4 de	2.0 cd	1.2 a	0 b	0.8 bc
Flagship 25WG (thiamethoxam)	8 oz	3.6 bc	0.6 a	0.8 bc	1.2 c	3.4 bc	0.8 a	0.6 b	0.6 bc
Hachi-Hachi (tolfenpyrad)	21 fl oz	1.0 d	0.2 a	0.8 bc	1.0 cd	0.4 d	0.6 a	0.2 b	1.6 a
Hachi-Hachi (tolfenpyrad)	32 fl oz	3.8 bc	0.2 a	0.2 c	1.4 bc	1.2 d	1.0 a	0.2 b	0.8 bc
Marathon G (imidacloprid)	7 g/pot	0 d	0.4 a	0.7 bc	0.4 de	1.6 d	0 a	0.4 b	0.2 c
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	1 lb per	4.4 ab	0.6 a	1.0 b	2.0 b	4.4 b	1.6 a	0.6 b	0.8 bc
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	2 lb	3.0 bc	3.0 a	2.0 a	0.2 e	4.0 bc	1.0 a	3.0 a	1.0 ab
Onyx (bifenthrin)	12.8 fl oz	3.4 bc	0.2 a	1.0 b	1.4 bc	3.6 bc	1.0 a	0.2 b	1.2 ab
Safari 20SG (dinotefuran)	8 oz	2.6 c	0.2 a	0.4 bc	1.0 cd	0.2 d	1.0 a	0.2 b	0.8 bc
TriStar 30SG (acetamiprid)	8 g	3.2 bc	0 a	0.4 bc	1.2 c	3.6 bc	2.8 a	0 b	0.6 bc
Nontreated		2.6 c	1.0 a	0.6 bc	3.0 a	8.4 a	1.0 a	1.0 b	1.2 ab

* Days post-exposure

^x Means followed by the same letter are not significantly different, P = 0.05

^z Rating: 1 = 10 % defoliation, 10 = 100 % defoliation.

Table 41. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Virginia Sweetspire (*Itea virginica*), Braman, 2013.

Treatment	Rate Per 100 Gal	No. of Beetles at Days Post Treatment ^x					% Leaf Damage at Days Post Treatment				
		1	10	21	24	30	1	10	21	24	30
Aloft (clothianidin/bifenthrin)	15 fl oz	0.2 c	0.2 b	1.0 a	0.7 a	0.7 a	0.2 c	0.2 c	0 c	2.0 cd	9.0 bc
Discus (imidacloprid + cyfluthrin)	50 oz	0.5 bc	1.0 a	0.7 a	0.7 a	0.7 a	1.6 c	1.6 c	0.3 c	0.7 d	1.0 c
Hachi-Hachi (tolfenpyrad)	32 fl oz	1.0 a	1.0 a	1.0 a	1.0 a	1.0 a	8.6 abc	8.6 abc	8.3 ab	18.7 abc	43.3 a
Mainspring/A20520A (cyantraniliprole)	16 oz	1.0 a	1.0 a	1.0 a	1.0 a	1.0 a	12.0 ab	12.0 ab	3.3 bc	16.9 a-d	21.7 abc
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	2 lb	1.0 a	1.0 a	1.0 a	1.0 a	1.0 a	3.2 c	3.2 c	11.7 a	26.7 a	45.0 a
Onyx (bifenthrin)	1.28 fl oz	0.7 ab	0.8 b	1.0 a	1.0 a	1.0 a	8.0 bc	8.0 bc	8.3 ab	18.3 a-d	33.3 ab
Safari 20SG foliar (dinotefuran)	8 oz	0.7 ab	0.8 a	1.0 a	1.0 a	1.0 a	0.2 c	0.2 c	9.0 ab	11.7 a-d	30.0 abc
Safari 20SG drench(dinotefuran)	24 oz	0.6 ab	0.6 ab	1.0 a	1.0 a	0.7 a	1.2 c	1.2 c	0 c	4.3 bcd	9.3 bc
XXpire/GF-2860 40WP (spinetoram+sulfoxaflor) + 6 oz Capsil	3.5 oz	0.8 ab	1.0 a	1.0 a	1.0 a	1.0 a	6.0 bc	6.0 bc	3.3 bc	20.0 ab	43.3 a
Nontreated		0.8 ab	1.0 a	1.0 a	1.0 a	1.0 a	8.0 a	15.0 a	15.0a	25.0 a	26.7 abc

^x Means followed by the same letter are not significantly different, P= 0.05

Table 42. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Virginia Sweetspire (*Itea virginica*), Frank, 2013.

Treatment	Rate Per 100 Gal	No. of Beetles/Plant at Days Post Initial Treatment ^x					% Leaf Area Consumed at Days Post Initial Treatment				
		-1	7	14	33	57	-1	7	14	33	57
Bifenthrin	40 fl oz	0.5 ab	0 b	0 a	0 a	0 b	13.5 a	17.50 a	7.08 b	1.92 cd	1.83 b
Cyantraniliprole	8 fl oz	0.3 ab	0 b	0 a	0 a	0 b	4.75 a	5.33 abc	2.58b	2.00 cd	9.17 ab
Cyantraniliprole	16 fl oz	0 b	0 b	0 a	0 a	0 b	3.83 a	12 abc	5.50 b	8.58 a-d	2.67 b
Discus (imidacloprid + cyfluthrin)	50 fl oz	0.3 ab	0 b	0.2 a	0 a	0 b	15.17 a	15.83 ab	7.83 b	1.67 cd	1.75 b
Hachi-Hachi (tolfenpyrad)	21 fl oz	0 b	0 b	0.2 a	0 a	0 b	7.83 a	8.50 abc	3.67 b	3.67 cd	4.58 b
Hachi-Hachi (tolfenpyrad)	32 fl oz	0 b	0 b	0 a	0 a	0.2 a	3.92 a	10.25 abc	3.92 b	5.92bcd	7.75 ab
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	1 lb	0.3 ab	0 b	0 a	0 a	0 b	5.92 a	2.17 c	3.83 b	15.25 ab	15.92 a
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	2 lb	0.2 ab	0 b	0 a	0 a	0 b	4.42 a	3.00 bc	3.00 b	0.42 d	2.58 b
Safari 20SG foliar (dinotefuran)	8 fl oz	0 b	0 b	0 a	0 a	0 b	10.92 a	6.83 abc	3.50 b	4.50 cd	3.58 b
Safari 20SG foliar (dinotefuran)	24 fl oz	0.2 ab	0 b	0 a	0 a	0 b	4.33 a	3.67 bc	6.25 b	4.33 cd	1.67 b
XXpire/GF-2860 (spinetoram + sulfoxaflor)	3.5 fl oz	0.3 ab	0 b	0 a	0 a	0 b	10.33 a	14.33 abc	0.50 b	1.83 cd	1.17 b
XXpire/GF-2860 (spinetoram + sulfoxaflor)	7 fl oz	0.8 a	0 b	0 a	0 a	0 b	4.83 a	11.17 abc	0.33 b	11.17 abc	9.33 ab
Nontreated		0.8 a	0.7 a	0 a	0 a	0 b	7.00 a	9.42 ab	18.75 a	16.83 a	9.66 ab

^x Means followed by the same letter are not significantly different, P= 0.05, Duncan's New MRT.

Table 43. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Virginia Sweetspire (*Itea virginica*) 'Little Henry', Sage (*Salvia nemorosa*) 'Blue Hill', and Stonecrop (*Sedum telephium*) 'Autumn Joy', Kunkel, 2014.

		Rate Per 100 Gal	Pretrt	7 DAT	14 DAT	21 DAT	28 DAT	35 DAT	42 DAT	49 DAT
Treatment			Percent Damaged Foliage on Virginia Sweetspire							
% Damaged Foliage on Viginia Sweetspire ^x										
Treatment	Rate (per 100 gal)	Pretrt	7 DAT	14 DAT	21 DAT	28 DAT	35 DAT	42 DAT	49 DAT	
Hachi-Hachi (tolfenpyrad)	21 fl oz	1.3 b	2.5 ab	11.9 a-e	25.0 a-d	21.3 a-i	36.3 a-h	26.3 a-g	38.8 a-g	
Hachi-Hachi (tolfenpyrad)	32 fl oz	4.4 a	4.4 ab	10.0 b-f	21.3 a-f	15.0 c-k	25.0 c-k	21.3 a-h	38.8 a-g	
Mainspring (cyantraniliprole)	8 fl oz	1.3 b	1.3 ab	10.6 a-f	23.8 a-d	36.3 a-d	36.3 a-g	30.0 a-g	47.5 a-d	
Mainspring (cyantraniliprole)	16 fl oz	0.0 b	2.5 ab	6.9 b-g	15.0 c-g	15.0 c-k	27.5 a-k	20.0 b-h	28.8 a-i	
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	16 oz	0.0 b	1.9 ab	3.1 fg	15.0 b-g	13.8d-k	26.3 b-k	17.5 b-h	25.0 b-j	
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	32 oz	0.0 b	0.0 b	5.6 b-g	15.0 b-g	12.5 f-k	31.3 a-i	20.0 b-h	21.3 b-j	
Safari 20SG foliar (dinotefuran)	8 oz	0.0 b	1.3 ab	16.3 ab	18.8 a-f	30.0 a-f	48.8 a	37.5 ab	36.3 a-g	
Safari 20SG drench (dinotefuran)	24 oz	0.0 b	0.6 ab	8.1 b-g	18.8 a-f	33.8 a-d	46.3abc	32.5 a-e	48.8 ab	
Scimitar (lambda-cyhalothrin)	5 fl oz	0.6 b	2.5 ab	3.8 efg	11.3 c-g	13.8 e-k	20.0 f-l	16.3 c-h	23.8 b-j	
XXpire/GF-2860 (spinetoram + sulfoxaflor) + Capsil	2.75 oz	0.0 b	5.0 ab	8.1 b-g	22.5 a-f	18.8 b-j	47.5 ab	32.5 a-e	48.8 abc	
XXpire/GF-2860 (spinetoram + sulfoxaflor) + Capsil	3.5 oz	0.6 b	1.3 ab	10.0 b-f	15.0 b-g	20.0 b-j	30.0 a-i	23.8 a-h	26.3 a-j	
Nontreated		1.3 b	5.6 ab	10.6 a-f	23.8 a-e	21.3 b-j	45.0 a-d	36.3 abc	43.8 a-e	
% Damaged Foliage on Sage ^x										
Hachi-Hachi (tolfenpyrad)	21 fl oz	0.8 a	5.8 ab	14.2 abc	26.7 abc	21.7 b-j	38.3 a-f	25.8 a-g	23.3 c-j	
Hachi-Hachi (tolfenpyrad)	32 fl oz	2.1 a	1.3 ab	9.2 b-g	21.7 a-f	25.8 a-g	32.5 a-h	28.3 a-f	30.0 a-h	
Mainspring (cyantraniliprole)	8 fl oz	1.7 a	4.2 ab	7.1 b-g	24.2a-d	23.3 a-g	36.7 a-g	31.3 a-f	56.7 a	
Mainspring (cyantraniliprole)	16 fl oz	0.8 a	2.1 ab	7.9 b-g	22.5 a-f	20.8 b-i	31.7 a-h	20.0 b-h	38.3 a-f	
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	16 oz	0.0 a	2.5 ab	12.9 a-d	33.3 ab	25.8 a-g	36.7 a-g	20.0 b-h	4.2 a-g	
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	32 oz	0.4 a	2.1 ab	7.1 b-g	12.9 c-g	17.5 d-j	25.4 e-k	15.8 e-h	26.7 c-j	
Safari 20SG foliar (dinotefuran)	8 oz	0.4 a	2.9 ab	10.8 b-e	20.8 a-f	20.8 b-i	39.2 a-e	21.7 b-h	22.5 d-j	
Safari 20SG drench (dinotefuran)	24 oz	1.3 a	2.1 ab	4.6 efg	9.6 e-h	11.7 j-k	25.0 e-k	13.3 fgh	16.7 hij	
Scimitar (lambda-cyhalothrin)	5 fl oz	0.0 a	2.1 ab	4.2 efg	8.8 e-h	9.6 ijk	20.8 g-l	11.7 gh	18.8 g-j	

Treatment	Rate Per 100 Gal	Pretrt	7 DAT	14 DAT	21 DAT	28 DAT	35 DAT	42 DAT	49 DAT
		<i>Percent Damaged Foliage on Virginia Sweetspire</i>							
XXpire/GF-2860 (spinetoram + sulfoxaflor) + Capsil	2.75 oz	1.7 a	8.8 a	21.7 a	33.3 a	30.0 a-e	44.2 a-d	28.3 a-f	21.7 d-j
XXpire/GF-2860 (spinetoram + sulfoxaflor) + Capsil	3.5 oz	0.8 a	2.1 ab	7.5 b-g	16.7 b-f	20.8 b-h	26.7 b-j	25.8 a-g	19.2 f-j
Nontreated		0.0 a	7.1 a	11.3 a-f	19.2 a-f	22.5 b-h	36.7 a-g	20.4 b-h	23.3b-j
<i>% Damaged Foliage on Stonecrop^x</i>									
Hachi-Hachi (tolfenpyrad)	21 fl oz	1.9 a	2.5 ab	12.5 a-d	15.6 b-g	18.8 b-j	25.0 d-k	25.0 a-h	30.0 a-j
Hachi-Hachi (tolfenpyrad)	32 fl oz	2.5 a	3.8 ab	9.4 b-g	36.3 a	47.5 a	36.3 a-g	45.0 a	21.3 c-j
Mainspring (cyantraniliprole)	8 fl oz	0.0 a	0.6 ab	9.4 b-g	18.8 a-f	20.0 b-i	22.5 e-l	22.5 a-h	28.8 a-i
Mainspring (cyantraniliprole)	16 fl oz	1.9 a	2.5 ab	8.1 b-g	14.4 b-g	21.9 b-j	20.6 f-l	15.0 d-h	27.5 a-j
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	16 oz	2.5 a	1.9 ab	10.6 a-f	18.8 a-f	16.3 b-j	14.4 i-m	15.6 d-h	18.8 f-j
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PRAA4-1T)	32 oz	0.0 a	5.6 ab	15.0 abc	23.8 a-d	35.0 abc	18.8 g-m	18.8 b-h	25.0 a-j
Safari 20SG foliar (dinotefuran)	8 oz	0.0 a	1.3 ab	8.8 b-g	9.4 e-h	17.5 c-k	16.9 h-m	13.8 e-h	20.0 f-j
Safari 20SG drench (dinotefuran)	24 oz	0.6 a	1.9 ab	4.4 d-g	6.3 gh	5.0 k	8.1 lm	6.3 h	10.0 ij
Scimitar (lambda-cyhalothrin)	5 fl oz	0.0 a	1.3 ab	2.5 g	3.8 h	8.1 jk	5.0 m	6.9 h	9.4 j
XXpire/GF-2860 (spinetoram + sulfoxaflor) + Capsil	2.75 oz	0.6 a	5.0 ab	9.4 b-g	13.8 c-g	7.5 jk	11.3 j-m	12.5 e-h	13.1 hij
XXpire/GF-2860 (spinetoram + sulfoxaflor) + Capsil	3.5 oz	0.6 a	3.1 ab	7.5 b-g	8.8 e-h	9.4 h-k	10.6 klm	10.0 gh	18.8 e-j
Nontreated		0.0 a	1.3 ab	13.1 a-d	36.3 a	38.8 ab	46.3 abc	35.0 a-d	37.5 a-g

^x Means followed by the same letter are not significantly different, P = 0.05, Tukey's HSD.

Graph 44. Foliar damage from Jul 1 to Oct 7 caused by Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Virginia Sweetspire (*Itea virginica*), Cloyd, 2016.




Table 45. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on 'Pee Gee' Hydrangea (*Hydrangea paniculata* 'Grandiflora'), Gilrein, 2016.

Treatment	Rate Per 100 Gal	% Mortality ^x				% Foliar Damage			
		7/1	7/5	7/8	7/15	7/1	7/5	7/8	7/15
BeetleGONE (<i>Bacillus thuringiensis galleriae</i>) + NuFilm P	16 lb + 0.125%	0.74 a	-	6.25 c	12.77 bc	2.08 a	-	60.83 b	79.16 a
Hachi-Hachi SC (tolfenpyrad)	21 fl oz	1.41 a	82.28 a	99.30 a	99.30 a	0.0 a	15.08 c	9.00 c	9.50 c
Hachi-Hachi SC	27 fl oz	0.69 a	81.52 ab	98.61 a	98.61 a	0.0 a	11.16 c	8.30 c	8.33 c
IKI-3106 50SL (cyclaniliprole)	22.0 fl oz	0.00 a	61.51 b	99.30 a	99.30 a	0.41 a	10.83 c	5.50 c	5.50 c
IKI-3106 50SL	27.0 fl oz	1.41 a	79.64 ab	79.64 ab	94.63 a	0.83 a	9.66 c	6.83 c	6.83 c
Preferal (<i>Isaria fumosoroseus</i>)	1 lb	0.00 a	-	4.57 c	12.70 bc	0.50 a	-	76.91 a	84.33 a
Scimitar GC (lambda-cyhalothrin)	3.2 fl oz	0.00 a	-	88.14 ab	97.65 a	0.00 a	-	12.16 c	6.41 c
XXpire 40WG (2860 (spinetoram + sulfoxaflor) + Capsil	3.5 oz + 6 fl oz	0.73 a	80.82 ab	80.92 b	91.58 a	0.83 a	18.33 c	13.16 c	5.58 c
Nontreated (water)	-	0.00 a	0.83 c	1.52 c	3.25 c	0.00 a	61.66 a	78.33 a	86.41 a

^xMeans within columns followed by the same letter are not significantly different at p=0.05, Tukey's HSD. Treatments applied on Jul 2; BeetleGone and Scimitar also applied on Jul 9.

Table 46. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Hydrangea (*Hydrangea paniculata*) 'Baby Lace', Chong, 2017.

Treatment	Rate Per 100 Gal	Applic Dates	0 DAT	2 DAT	7 DAT	14 DAT	21 DAT	28 DAT	35 DAT
<i>Numbers of adults</i>									
BeetleGone! (<i>Bacillus thuringiensis galleriae</i>) + LI700	16 lb + 0.1%	7/26, 8/2, 11	1.8 ± 0.5 a	0.7 ± 0.3 a	1.7 ± 0.5 a	0.8 ± 0.4 a	0.7 ± 0.4 a	0.3 ± 0.2 a	1.6 ± 0.4 a
Hachi-Hachi SC (tolfenpyrad)	27 fl oz	7/26	1.8 ± 0.4 a	0.2 ± 0.1 a	2.1 ± 0.5 a	0.4 ± 0.3 a	0.8 ± 0.3 a	0.1 ± 0.1 a	2.0 ± 0.7 a
IKI-3106 50SL (cyclaniliprole)	22 fl oz	7/26, 8/11	2.0 ± 0.6 a	0.1 ± 0.1 a	2.2 ± 0.6 a	1.3 ± 0.5 a	1.0 ± 0.4 a	0.1 ± 0.1 a	1.2 ± 0.5 a
Preferal (<i>Isaria fumosoroseus</i>)	1 lb	7/26	1.8 ± 0.4 a	0.1 ± 0.1 a	1.1 ± 0.3 ab	0.7 ± 0.3 a	0.1 ± 0.1 a	0.3 ± 0.2 a	0.9 ± 0.3 a
Scimitar GC (lambda-cyhalothrin)+ Capsil	5 fl oz + 6 fl oz	7/26, 8/2, 11	1.3 ± 0.5 a	0.1 ± 0.1 a	1.6 ± 0.3 a	1.3 ± 0.3 a	0.7 ± 0.2 a	0.2 ± 0.1 a	1.1 ± 0.3 a
VST-006350 + LI700	3.33 L + 0.1%	7/26, 8/2, 11	1.8 ± 0.5 a	0.7 ± 0.5 a	0.7 ± 0.4 b	0.4 ± 0.2 a	0.3 ± 0.2 a	0.2 ± 0.1 a	1.0 ± 0.4 a
VST-006350 + Beetle Gone! + LI700	3.33 L + 16 lb + 0.1%	7/26, 8/2, 11	1.8 ± 0.4 a	0.8 ± 0.3 a	1.3 ± 0.2 a	0.6 ± 0.2 a	0.3 ± 0.2 a	0.2 ± 0.1 a	1.8 ± 0.5 a
Nontreated (Water)	-	-	1.7 ± 0.4 a	0.4 ± 0.3 a	1.2 ± 0.4 ab	0.6 ± 0.2 a	1.1 ± 0.7 a	0.3 ± 0.2 a	1.8 ± 0.7 a
<i>Percent Defoliation</i>									
BeetleGone! (<i>Bacillus thuringiensis galleriae</i>) + LI700	16 lb + 0.1%	7/26, 8/2, 11	17.8 ± 2.9 ab	9.4 ± 1.3 a	9.6 ± 1.6 a	10.8 ± 1.6 a	13.7 ± 1.5 bcd	12.4 ± 0.8 ab	16.2 ± 1.8 a
Hachi-Hachi SC (tolfenpyrad)	27 fl oz	7/26	10.0 ± 1.9 c	7.8 ± 1.2 a	8.2 ± 0.7 a	11.8 ± 1.8 a	18.9 ± 2.7 ab	9.5 ± 1.7 b	13.3 ± 1.9 a
IKI-3106 50SL (cyclaniliprole)	22 fl oz	7/26, 8/11	12.2 ± 1.5 bc	10.0 ± 0.8	10.2 ± 0.7 a	12.4 ± 1.3 a	18.1 ± 2.5 abc	9.8 ± 1.1 b	12.7 ± 1.7 a
Preferal (<i>Isaria fumosoroseus</i>)	1 lb	7/26	11.7 ± 1.4 c	9.4 ± 1.0 a	9.6 ± 1.1 a	12.2 ± 1.7 a	12.0 ± 1.3 d	11.9 ± 1.6 ab	12.2 ± 1.5 a
Scimitar GC (lambda-cyhalothrin)+ Capsil	5 fl oz + 6 fl oz	7/26, 8/2, 11	13.9 ± 1.8 abc	10.6 ± 1.0 a	10.1 ± 0.5 a	14.1 ± 1.5 a	13.6 ± 1.9 bcd	11.6 ± 1.4 ab	14.3 ± 1.7 a
VST-006350 + LI700	3.33 L + 0.1%	7/26, 8/2, 11	12.8 ± 1.9 bc	10.0 ± 1.2 a	9.9 ± 0.8 a	11.9 ± 1.0 a	13.0 ± 1.0 cd	11.9 ± 1.6 ab	14.4 ± 1.7 a
VST-006350 + Beetle Gone! + LI700	3.33 L + 16 lb + 0.1%	7/26, 8/2, 11	19.4 ± 3.2 a	11.1 ± 1.4 a	11.0 ± 0.6 a	17.0 ± 1.7 a	19.6 ± 0.9 a	13.9 ± 1.4 a	17.1 ± 1.5 a
Nontreated (Water)	-	-	13.3 ± 2.4 bc	12.2 ± 1.2 a	9.6 ± 0.5 a	15.6 ± 1.9 a	23.3 ± 2.4 a	15.0 ± 1.4 a	16.3 ± 1.0 a

^x Mean number of adults counted from 4 leaves pre-treatment and after the first treatment application (DAT). Means within column followed by the same letter are not significantly different (Fisher's LSD, P=0.05).

Table 47. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Forsythia (*Forsythia* sp.), Kunkel, 2017a.

Treatment	Rate Per 100 Gal	Applic. Dates	No. Live Adults / Plant						No. Damaged Leaves / Plant		
			Jun 7	Jun 11	Jun 14	Jun 18	Jun 21	Jun 28	Jun 7	Jun 11	Jun 14
BeetleGONE (<i>Bacillus thuringiensis galleriae</i> strain SDS-502)	16 oz	5/31, 6/7, 14	10.2	3.7	11.7 bc	1.5	0.5	0.7	27.2	17.5	55.5
Hachi-Hachi (tolfenpyrad)	27 fl oz	5/31	7.8	n/a	19.7 ab	1.3	2.5	0.0	17.0	n/a	61.2
IKI-3106 (cyclaniliprole)	22 fl oz	5/31, 6/14	12.5	n/a	25.2 a	2.3	0.5	0.3	35.5	n/a	49.0
Preferal (<i>Isaria fumosoroseus</i>)	8 oz	5/31, 6/14	11.5	n/a	19.0 ab	3.5	1.2	0.3	25.0	n/a	53.8
Tristar (acetamiprid)	12 oz	5/31, 6/14	15.3	n/a	15.3 bc	2.2	1.2	1.3	19.5	n/a	49.8
Upstar-Gold (bifenthrin)	8 fl oz	5/31, 6/14	6.2	n/a	10.0 c	2.2	1.5	1.0	15.2	n/a	58.5
VST6350 (VST6350) + Li-700	3.33 l + 1% v/v	5/31, 6/7, 14	16.5	3.5	16.7 abc	3.3	1.3	0.2	25.0	18.8	64.3
VST6350 + BeetleGONE + LI-700	3.33 l + 16 oz + 1%	5/31, 6/7, 14	10.0	3.0	14.7 bc	4.7	0.8	0.2	24.8	13.2	49.5
Nontreated	-	-	6.2	2.8	8.0 c	2.0	0.5	0.2	31.0	18.8	58.0
P-value from ANOVA			ns	ns	0.0254	ns	ns	ns	ns	ns	ns

^x Means followed by the same letter are not significantly different, P = 0.05, Fisher's LSD.

^y Scale of 0-2 where: 0=no feeding damage, 1=light feeding damage, 2=heavy feeding damage.

^z Scale of 0-3 where: 0=no damage, 1=light damage, 2=medium damage, , 3=heavy damage.

Table 48. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Forsythia (*Forsythia* sp.), Kunkel, 2017a.

Treatment	Rate Per 100 Gal	Applic. Dates	% Branches with RHFB Feeding Damage			New Growth Damage Rating ^y				% Damage d Leaves Jun 28	Final Overall Damage Rating ^z
			Jun 7	Jun 11	Jun 14	Jun 14	Jun 18	Jun 21	Jun 28		
BeetleGONE (<i>Bacillus thuringiensis galleriae</i> strain SDS-502)	16 oz	5/31, 6/7, 14	75.0	90.0	98.3	1.3 bc	1.1 bc	0.9 bc	0.7 cd	5.5 ± 1.1	2.0 bc
Hachi-Hachi (tolfenpyrad)	27 fl oz	5/31	68.3	n/a	88.3	1.5 ab	1.4 ab	1.1 abc	1.2 ab	3.5 ± 0.8	2.8 a
IKI-3106 (cyclaniliprole)	22 fl oz	5/31, 6/14	76.7	n/a	100.0	1.6 a	1.4 ab	1.2 abc	0.8 bc	5.8 ± 1.8	2.3 ab
Preferal (<i>Isaria fumosoroseus</i>)	8 oz	5/31, 6/14	65.0	n/a	96.7	1.0 cd	1.2 b	1.0 bc	0.6 cd	4.5 ± 0.9	2.3 ab
Tristar (acetamiprid)	12 oz	5/31, 6/14	70.0	n/a	96.7	1.6 a	1.2 ab	0.8 cd	0.9 bc	4.1 ± 0.9	1.5 cd
Upstar-Gold (bifenthrin)	8 fl oz	5/31, 6/14	46.7	n/a	93.3	0.8 d	0.7 c	0.4 d	0.3 d	3.1 ± 0.6	1.2 d
VST6350 (VST6350) + Li-700	3.33 l + 1% v/v	5/31, 6/7, 14	71.7	85.0	98.3	1.6 ab	1.3 ab	1.5 a	1.3 ab	4.9 ± 0.8	2.7 a
VST6350 + BeetleGONE + LI-700	3.33 l + 16 oz + 1%	5/31, 6/7, 14	66.7	91.7	98.3	1.4 ab	1.2 b	0.8 c	0.5 cd	5.0 ± 0.8	1.5 cd
Nontreated	-	-	73.3	88.3	100.0	1.6 a	1.6 a	1.3 ab	1.4 a	5.6 ± 1.5	2.7 a
P-value from ANOVA			ns	ns	ns	<0.0001	0.0165	0.0008	<0.0001	ns	<0.0001

^x Means followed by the same letter are not significantly different, P = 0.05, Fisher's LSD.

^y Scale of 0-2 where: 0=no feeding damage, 1=light feeding damage, 2=heavy feeding damage.

^z Scale of 0-3 where: 0=no damage, 1=light damage, 2=medium damage, , 3=heavy damage.

Table 49. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Panicle Hydrangea (*Hydrangea paniculata*) ‘Vanilla Strawberry’, Frank, 2017.

Treatment	Rate Per 100 Gal	Percent Leaf Damage to Leaf Disks Collected at Various Intervals After Initial Treatment ^x									No. Dead Beetles ^y	% Damage on Leaves ^z
		1 Day After Treatment			7 Days After Treatment			14 Days After Treatment				
		Day* 1	Day 3	Day 5	Day 1	Day 3	Day 5	Day 1	Day 3	Day 5		
BeetleGONE + Capsil	16 lb	17.5 c	34.2 b	46.7 c	12.7 a	43.4 a	78.9 ab	18.3 bc	42.4	65.7	0.50 ± 0.22	2.8 ± 1.0
Hachi Hachi SC (tolfenpyrad)	27 fl oz	8.6 d	20.3 c	28.2 d	8.4 bc	43.0 a	63.8 bc	17.8 bc	29.1	39.1	0.17 ±0.17	2.3 ± 0.7
IKI-3106 (cyclaniliprole)	22 fl oz	9.4 d	11.5 cd	17.8 de	14.4 a	49.0 a	89.3 a	21.1 abc	36.9	53.2	1.67 ± 0.49	1.2 ± 0.8
Marathon (imidacloprid)	1.7 fl oz	5.3 d	4.0 d	4.8 b	7.6 c	25.3 b	50.3 c	14.8 c	37.2	58.8	4.83 ± 0.17	7.2 ± 0.8
Preferal (<i>Isaria fumosorosea</i> Apopka Strain 97)	1 lb	23.3 ab	49.1 a	67.4 ab	12.3 ab	43.1 a	76.2 ab	15.5 c	41.0	64.7	0.17 ± 0.17	1.6 ± 0.8
VST 06330 + Capsil	2 lb	24.3 a	38.4 b	66.3 ab	13.3 a	47.5 a	87.8 a	16.4 c	39.8	59.0	0.17 ± 0.17	3.1 ± 1.9
VST 06330 + BeetleGONE + Capsil	2 lb + 16 lb	18.3 bc	42.5 ab	54.2 bc	12.9 a	40.4 a	76.7 ab	24.9 ab	47.0	75.3	0.00 ± 0.00	2.8 ± 1.1
Nontreated	-	23.7 a	49.9 a	74.8 a	11.2 abc	38.3 a	67.3 bc	26.6 a	40.5	60.3	0.00 ± 0.00	8.8 ± 0.8
	<i>F</i>	18.92	22.81	23.68	2.71	3.26	3.56	2.29	1.11	1.45	2.29 ±	24.07
	<i>P</i>	0.001	0.001	0.001	0.021	0.008	0.005	0.046	0.377	0.214	0.046	0.001

* Days of exposure

^x Means within columns followed by the same letter are not significantly different.

^y Number of dead flea beetles (out of 5) after five days of exposure to leaves in petri dishes collected 1 day after treatment.

^z Estimated percent damage on 5 marked leaves on experimental plants outdoors.

Table 50. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Panicle Hydrangea (*Hydrangea paniculata* ‘Bokrathirteen’) ‘Sweet Summer’, Gilrein, 2018.

Treatment	Rate Per 100 Gal	Applic. Dates	% Insect Mortality ^x					% Foliage Damaged ^{x,y}				
			Jul 3	Jul 6	Jul 10	Jul 17	Jul 31	Jul 3	Jul 6	Jul 10	Jul 17	Jul 31
Ancora (<i>Isaria fumosoroseus</i>)	1.0 lb	7/3	0.0 a	3.47 e	7.84 c	15.28 d	25.71 c	1.72 a	25.00 a	26.94 a	46.94 a	63.61 a
Hachi-Hachi SC (tolfenpyrad)	27.0 fl oz	7/3	0.0 a	77.38 b	84.39 a	89.48 a	92.52 a	2.22 a	12.78 c	12.50 bc	17.50 bc	16.11 b
IKI-3106 (cyclaniliprole)	22.0 fl oz	7/3, 18	0.0 a	94.44 a	95.83 a	98.61 a	100.00 a	2.28 a	7.28 d	7.61 cd	8.11 cd	9.00 bc
Pyridium (beta-cyfluthrin)	2.1 fl oz	7/3	0.0 a	93.06 ab	95.83 a	96.53 a	96.53 a	2.44 a	4.89 d	5.22 d	6.89 d	11.44 bc
TriStar 8.5SL (acetamiprid)	25.3 fl oz	7/3, 18	0.0 a	94.44 a	96.53 a	97.92 a	100.00 a	2.06 a	5.06 d	5.83 d	5.61 d	6.89 c
V-10433 (V-10433)	11.0 fl oz	7/3, 10, 18	0.0 a	20.83 d	26.42 b	34.66 c	57.34 b	2.28 a	18.61 abc	21.11 ab	36.94 a	51.94 a
V-10433 (V-10433)	22.0 fl oz	7/3, 10, 18	0.0 a	43.87 c	38.19 b	57.24 b	68.61 b	2.56 a	22.22 ab	18.61 ab	23.89 ab	35.83 a
Nontreated	-	7/3	0.0 a	3.47 e	9.03 c	10.19 d	18.62 c	2.39 a	14.44 bc	28.61 a	42.50 a	55.00 a

^x Means within columns followed by the same letter are not significantly different at p=0.05 (LS Means Tukey's HSD).

^y Data were transformed prior to analysis using log(y)

Table 51. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Panicle Hydrangea (*Hydrangea paniculata*) ‘Vanilla Strawberry’, Frank, 2019.

Product	Rate/100 gal	Mean Flea Beetles observed on hydrangeas					Plant Growth	
		3 DAIT	7 DAIT	14 DAIT	21 DAIT	28 DAIT	Change in Height	Change in Width
Acelepryn	16 fl oz	2.0 a	1.3 a	0.3	0.0	0.2	28.8	20
Ancora	1 lb	1.0 bc	1.3 a	0.0	0.2	0.2	24.1	33
BCS-507	16 oz	0.0 d	0.2 bc	0.0	0.3	0.0	24.1	27.9
Hachi Hachi SC	27 fl oz	0.0 d	0.0 c	0.0	0.2	0.2	25.3	16.6
IKI-3106	27 fl oz	0.2 cd	0.0 c	0.6	0.3	0.0	26.3	30.8
KOC22018-8	128 fl oz	1.5 ab	0.8 abc	0.2	0.0	0.0	25.3	20.2
Mainspring GNL	16 fl oz	0.5 cd	1.2 ab	0.2	0.0	0.0	23.0	23.8
TetraCURB Conc.	128 fl oz	0.7 bc	1.8 a	0.6	0.0	0.0	27.3	23.0
TetraCURB Organic	128 fl oz	2.2 a	1.7 a	0.3	0.0	0.2	23.0	30.1
V-10433	22 fl oz	0.3 cd	0.8 abc	0.3	0.2	0.2	31.7	24.4
Tristar	25 fl oz	0.0 d	0.0 c	0.0	0.2	0.0	20.3	24.9
Nontreated	N/A	2.3 a	1.7 a	0.3	0.3	0.2	27.1	27.4

Table 52. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Panicle Hydrangea (*Hydrangea paniculata*) ‘Moonrock’, Gilrein, 2019.

Treatment	Rate Per 100 Gal	Applic. Dates	% Insect Mortality ^x						% Foliage Damaged ^{x,y}					
			Jul 2	Jul 5	Jul 9	Jul 16	Jul 23	Jul 30	Jul 2	Jul 5	Jul 9	Jul 16	Jul 23	Jul 30
Acelepryn	16 fl. oz	7/2, 17	0.0 a	14.6 c	28.3 cd	35.6 d	52.6 b	67.6 b	5.67 a	14.4 a	15.8 bc	23.8 b	37.1 ab	45.0 a
BCS-507	16 fl. oz	7/2, 17	0.0 a	89.4 a	100.0 a	100.0 a	100.0 a	100.0 a	4.25ab	3.6 c	4.7 d	5.4 d	5.0 d	5.0 d
IKI-3106 (cyclaniliprole)	22 fl. oz	7/2	0.0 a	96.4 a	100.0 a	100.0 a	100.0 a	100.0 a	4.42 ab	4.0 c	5.0 d	5.0 d	4.8 d	5.0 d
KOC22018-8 (botanical oil blend)	128 fl. oz	7/2, 9, 17	0.0 a	50.3 b	52.7 b	88.1 ab	97.9 a	95.2 a	3.3 ab	12.2 a	19.4 ab	17.9 b	18.8 bc	17.5 bc
Mainspring GNL	16 fl. oz	7/2	0.0 a	73.3 ab	90.6 a	97.9 a	98.8 a	100.0 a	3.2 ab	4.5 bc	9.8 c	9.6 c	6.9 d	13.8 cd
TetraCURB Conc. (rosemary oil)	128 fl. oz	7/2, 9, 17	0.0 a	62.1 b	55.7 b	72.0 bc	95.5 a	94.0 a	3.3 ab	10.1 ab	20.0 ab	25.0 b	16.2 c	36.7 ab
TriStar	25.3 fl oz	7/2	0.0 a	97.8 a	100.0 a	100.0 a	100.0 a	100.0 a	2.5 ab	4.0 c	4.0 d	5.0 d	4.6 d	5.0 d
V-10433	22 fl oz	7/2, 9, 17	0.0 a	60.2 b	47.6 bc	64.6 c	84.8 a	90.2 ab	4.8 ab	15.5 a	19.2 ab	27.1 ab	26.0 abc	39.2 ab
Nontreated	-	-	0.0 a	11.1 c	10.7 d	24.4 d	38.2 b	41.6 c	2.2 b	19.2 a	33.8 a	42.9 a	47.1 a	67.5 a

^x Means within columns followed by the same letter are not significantly different at p=0.05 (LS Means Tukey's HSD).

^y Data were transformed prior to analysis using log(y)

Table 53. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Panicle Hydrangea (*Hydrangea paniculata*) ‘Moonrock’, Gilrein, 2020.

Treatment	Rate Per 100 Gal	Applic. Dates	% Insect Mortality ^x						% Foliage Damaged ^{x,y}					
			Jul 2	Jul 5	Jul 9	Jul 16	Jul 23	Jul 30	Jul 2	Jul 5	Jul 9	Jul 16	Jul 23	Jul 30
BW133	5.0 lb	7/2, 7/9, 7/16	0.0ns	49.76ab	50.51ab	94.84a	99.21a	100.00a	0.0ns	5.44 bcd	6.94 cd	9.72 c	10.69 b	14.31 b
BW238 ES	2.0 qt	7/2, 7/9, 7/16	0.0ns	27.01 bc	50.32ab	90.05a	99.31a	100.00a	0.0ns	6.61 bc	10.56 bc	13.06 bc	9.44 bc	13.33 b
ISM-555 + CapSil	5.76 fl oz	7/2, 7/9, 7/16	0.0ns	100.00a	100.00a	100.00a	100.00a	100.00a	0.0ns	2.00 d	4.39 d	4.39 c	4.39 c	4.39 c
MBI-203 SC2	128.0 fl oz	7/2, 7/9	0.0ns	11.71 c	16.76 bc	37.53 b	51.51 bc	62.14 b	0.0ns	7.22 b	15.28ab	21.53ab	30.00a	36.67a
MBI-306	5.0 fl oz	7/2, 7/9	0.0ns	16.19 bc	9.87 c	19.61 b	33.41 bc	30.16 c	0.0ns	6.78 b	19.72a	27.08a	35.69a	45.42a
SP3014 + CapSil	26.0 fl oz	7/2, 7/9, 7/16	0.0ns	20.62 bc	24.60 bc	35.15 b	55.73 b	55.55 bc	0.0ns	9.08ab	11.92 bc	20.00ab	25.83a	35.56a
TriStar	25.3 fl oz	2-Jul	0.0ns	96.85a	100.00a	100.00a	100.00a	100.00a	0.0ns	2.17 cd	4.89 d	4.89 c	4.89 c	4.89 c
Nontreated		7/2, 7/9, 7/16	0.0ns	6.07 c	9.59 c	14.33 b	24.18 c	41.60 bc	0.0ns	12.22a	16.67ab	29.17a	46.53a	50.14a

^x Means within columns followed by the same letter are not significantly different at p=0.05 (LS Means Tukey's HSD).

^y Data were transformed prior to analysis using log(y)

Table 54. Efficacy of several insecticides applied as foliar sprays for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Panicle Hydrangea (*Hydrangea paniculata*) ‘Limelight’, Sadof, 2021.

Treatment – Rate per 100 gal	Mean Number of Leaves Chewed per Plant					Mean Percentage of Damaged Leaves per Plant			Mean number of Beetles per plant in 30 seconds
	29 DAT (June 22)	35 DAT (June 28)	42 DAT (July 5)	49 DAT (July 12)	58 DAT (July 21)	42 DAT (July 5)	49 DAT (July 12)	58 DAT (July 21)	58 DAT (July 21)
ISM-555 – 3.84 oz	14.5 ± 8.2	37.7 ± 13.9	32.7 ± 3.6	74.0 ± 13.9	113.7 ± 13.2	11.7 ± 3.8	30 ± 11.5	51.7 ± 6.7	1.2 ± 0.7
ISM-555 – 5.76 oz	1.0 ± 0.3	8.8 ± 3	13.3 ± 3.1	41.0 ± 6	58.2 ± 6.6	5.8 ± 1.5	10.8 ± 2	15 ± 1.8	0.8 ± 0.3
MBI-203 – 128 fl oz	8.3 ± 3.3	97.7 ± 24.6	110.8 ± 31.2	210.3 ± 40.8	195.7 ± 32.6	51.7 ± 12.8	78.3 ± 7.9	75.8 ± 4.9	2.7 ± 1.3
MBI-306 – 2.5 fl oz	23.8 ± 13	158.5 ± 42.7	141.7 ± 46.8	253.8 ± 66.1	219.8 ± 44.2	46.7 ± 16.1	90 ± 2.6	86.7 ± 4.4	5.8 ± 2
MBI -306 – 5 fl oz	15.7 ± 10.8	102.2 ± 6	117.3 ± 15.7	208.3 ± 28	185 ± 9.9	63.3 ± 9.9	93.3 ± 4.9	70.8 ± 11.4	2.8 ± 0.5
SP3014 – 26 fl oz	20.8 ± 8.7	115.2 ± 26.2	78.8 ± 10.3	149.7 ± 27.4	178.3 ± 31	46.7 ± 12.3	78.3 ± 10.8	80 ± 5.3	3.3 ± 2.2
SP3014 – 13 fl oz	26.7 ± 12.3	136.5 ± 43.5	99.2 ± 14.1	173.2 ± 42.2	200.5 ± 50	41.7 ± 13	75 ± 10.6	85 ± 4.3	6.3 ± 1.9
Pradia – 17 fl oz	18.3 ± 8.3	62 ± 11.9	64.5 ± 10.6	190.5 ± 34	198.2 ± 43.4	30 ± 5.2	76.7 ± 8	75.8 ± 7.2	3.7 ± 1.2
Sarissa – 25 fl oz	8.2 ± 2.6	40.3 ± 16.5	49.2 ± 11.7	158.3 ± 22.3	195 ± 40.8	26.7 ± 13	58.3 ± 12.5	66.7 ± 7.3	1.7 ± 0.6
Acelepryn – 8 fl oz	11.8 ± 6.2	104.7 ± 17	113.2 ± 21.6	185.3 ± 27.5	171.8 ± 19.2	60 ± 11	83.3 ± 9.2	73.3 ± 5.6	3.2 ± 1
Acelepryn – 6 fl oz	7.3 ± 3.4	64.2 ± 16	95 ± 25.7	211.7 ± 42.5	199.8 ± 26.2	36.7 ± 12.8	80 ± 6.3	75.5 ± 7	6.3 ± 2.3
Xxpire – 2.5 oz	6.2 ± 2.7	28.3 ± 6.6	109.2 ± 30.8	238.2 ± 57.4	190 ± 35.4	38.3 ± 7	68.3 ± 12.5	71.7 ± 6	4 ± 1.2
Water	10.7 ± 6.6	102.5 ± 16.8	116.7 ± 24.8	195.2 ± 40.3	235.5 ± 73.7	45.8 ± 12.7	85.8 ± 5.8	78.3 ± 6.9	7.2 ± 1.8

Table 55. Efficacy of several insecticides for Redheaded Flea Beetle (*Systema frontalis*) adults feeding on Panicle Hydrangea (*Hydrangea paniculata*) ‘Fire and Ice’, Gilrein, 2021.

Treatment	Rate/ 100gal	2021 Average Beetle Mortality (%)						Percent Foliar Damage					
		6/29	7/2*	7/6*	7/13**	7/20**	7/27**	6/29	7/2**	7/6**	7/13*	7/20**	7/27**
MBI-203 SC2	128.0 fl oz	0.0ns	13.49 b	21.85 bc	22.84 c	32.71 b	56.96 bcd	0.0ns	4.72abcd	16.67 cd	37.50 c	54.58a	68.06a
MBI-306	5.0 fl oz	0.0ns	5.90 b	21.44 bc	12.37 c	36.03 b	46.06 cd	0.0ns	9.86a	33.06ab	75.28a	47.92a	58.33a
SP3014	26.0 fl oz	0.0ns	14.78 b	31.24 b	48.91 b	58.05 b	80.46ab	0.0ns	6.63abc	26.94abc	37.64 c	39.44a	35.97a
SP3014	13.0 fl oz	0.0ns	5.70 b	15.64 bc	23.80 c	43.98 b	47.78 cd	0.0ns	8.28a	42.50a	49.58 bc	58.61a	61.81a
Velifer	13.0 fl oz	0.0ns	8.68 b	8.71 c	60.38 b	57.60 b	78.07abc	0.0ns	7.78ab	22.64 bc	47.78 c	50.83a	60.00a
V-10433	11.0 fl oz	0.0ns	2.93 b	8.10 c	8.66 c	29.81 b	40.79 d	0.0ns	7.50ab	32.50ab	70.28ab	60.56a	58.47a
ISM-555	3.84 fl oz	0.0ns	98.96a	100.00a	100.00a	100.00a	100.00a	0.0ns	3.89 bcd	10.42 de	10.28 d	10.28 b	13.75 b
ISM-555	5.76 fl oz	0.0ns	100.00a	100.00a	100.00a	100.00a	100.00a	0.0ns	2.28 d	8.75 e	8.75 d	8.75 b	10.56 b
TriStar	25.3 fl oz	0.0ns	100.00a	100.00a	100.00a	100.00a	100.00a	0.0ns	3.25 cd	8.61 e	8.61 d	8.61 b	6.94 b
Nontreated		0.0ns	10.12 b	17.49 bc	23.91 c	33.43 b	33.14 d	0.0ns	6.81abc	33.72ab	46.81 c	51.25a	67.11a

Comparative Efficacy on European Elm Flea Weevil (Orchestes alni)

The European elm flea weevil (*Orchestes alni*) is a native to Europe. It feeds many elm species but can be especially damaging to Siberian and hybrid elms. Adults overwinter and feed during the early spring mainly on the underside of leaves, producing shothole injuries to the leaf interior. Soon after bud break the adults lay eggs and larvae hatch. The larvae are leafminers and feed in a serpentine pattern moving toward the leaf margin. They drop to the ground to pupate and adults emerge and feed for the remainder of the growing season. Damage is primarily aesthetic.

In 2013, Jones compared efficacy of several products applied foliar or drench for European elm flea weevil feeding (EEFW) on elm (Table 56 - Table 58). All treatments resulted in better control of European elm flea weevil activity (% leaf area affected, % the tree canopy affected, and leafminer activity) than the Nontreated control. Additionally, all treatments provided EEFW control that consumers would find acceptable. Of the treatments, Xytect resulted in the least EEFW activity. There were no significant difference in EEFW activity between the two Hachi-Hachi rates and the two MBI-203 rates.

Table 56. Efficacy of several insecticides for European Elm Flea Weevil (*Orchestes alni*) feeding on Elm (*Ulmus* sp.) ‘Patriot’ - % leaf area affected, Jones, 2013.

% Leaf Area Affected by Weevil Feeding ^x									
Treatment ^z	Rate Per 100 Gal	5/23	5/31	6/10	6/17	7/1	7/16	7/30	8/28
Aloft (clothianidin/bifenthrin)	15 fl oz	3 a	6 b	9 bc	9 b	14 b	11 cd	8 bc	6 b
Hachi-Hachi (tolfenpyrad)	21 fl oz	3 a	7 b	9 bc	9 b	14 b	17 b	10 b	6 b
Hachi-Hachi (tolfenpyrad)	32 fl oz	3 a	7 b	9 bc	9 b	14 b	15 bc	9 b	5 b
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PR4A4-1T)	1 lb	3 a	6 b	10 b	9 b	15 b	16 b	8 bc	5 b
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PR4A4-1T)	2 lb	3 a	6 b	9 bc	9 b	13 b	14 bc	9 b	6 b
Safari 20SG (dinotefuran)	12 g / DBH inch	3 a	6 b	9 bc	8 b	13 b	9 d	5 bc	4 bc
Xytect 2F (imidacloprid)	0.2 fl oz / DBH inch	3 a	4 b	7 c	5 b	11 b	8 d	4 c	2 c
Nontreated		4 a	14 a	19 a	24 a	33 a	23 a	19 a	13 a

^x Means followed by the same letter are not significantly different, P = 0.05, Duncan's New MRT.

^yRatings were made using a 0-1 scale where 0 = no leafminer activity and 1 = leafminer activity present.

^z Aloft and Hachi-Hachi applied foliar on May 18, MBI 203 applied foliar on May 18, 23, 31 and Jun 10; Xytect and Safari applied as soil drench on Apr 5 and May 6, respectively.

Table 57. Efficacy of several insecticides for European Elm Flea Weevil (*Orchestes alni*) feeding on Elm (*Ulmus* sp.) ‘Patriot’ - % tree canopy affected, Jones, 2013.

% Tree Canopy Area Affected by Weevil Feeding ^x									
Treatment ^z	Rate Per 100 Gal	5/23	5/31	6/10	6/17	7/1	7/16	7/30	8/28
Aloft (clothianidin/bifenthrin)	15 fl oz	3 a	11 b	19 b	14 b	19 b	25 cd	29 bc	18 b
Hachi-Hachi SC (tolfenpyrad)	21 fl oz	3 a	12 b	18 b	14 b	19 b	38 b	40 b	19 b
Hachi-Hachi SC (tolfenpyrad)	32 fl oz	3 a	12 b	18 bc	13 b	19 b	33 bc	38 b	16 b
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PR4A4-1T)	1 lb	3 a	11 b	9 b	14 b	20 b	38 b	33 b	19 b
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PR4A4-1T)	2 lb	3 a	11 b	18 bc	14 b	18 b	31 bc	39 b	17 b
Safari 20SG (dinotefuran)	12 g / inch DBH	3 a	11 b	17 bc	13 b	18 b	17 d	21 cd	13 b
Xytect 2F (imidacloprid)	0.2 fl oz / inch DBH	3 a	8 b	14 c	10 b	18 b	15 d	12 d	6 c
Nontreated		4 a	24 a	29 a	29 a	38 a	61 a	76 a	68 a

^x Means followed by the same letter are not significantly different, P = 0.05, Duncan's New MRT.

^yRatings were made using a 0-1 scale where 0 = no leafminer activity and 1 = leafminer activity present.

^z Aloft and Hachi-Hachi applied foliar on May 18, MBI 203 applied foliar on May 18, 23, 31 and Jun 10; Xytect and Safari applied as soil drench on Apr 5 and May 6, respectively.

Table 58. Efficacy of several insecticides for European Elm Flea Weevil (*Orchestes alni*) feeding on Elm (*Ulmus sp.*) ‘Patriot’ - presence of leafmine activity, Jones, 2013.

<i>Presence of Weevil Leafminer Activity^{x,y}</i>				
Treatment^z	Rate Per 100 Gal	5/31	6/10	6/17
Aloft (clothianidin/bifenthrin)	15 fl oz	0.1 b	0.1 b	0.1 b
Hachi-Hachi SC (tolfenpyrad)	21 fl oz	0.1 b	0.1 b	0.1 b
Hachi-Hachi SC (tolfenpyrad)	32 fl oz	0.1 b	0.0 b	0.0 b
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PR4A4-1T)	1 lb	0.0 b	0.0 b	0.0 b
MBI-203 DF (<i>Chromobacterium subtsugae</i> strain PR4A4-1T)	2 lb	0.1 b	0.0 b	0.1 b
Safari 20SG (dinotefuran)	12 g / inch DBH	0.1 b	0.1 b	0.1 b
Xytect 2F (imidacloprid)	0.2 fl oz / inch DBH	0.0 b	0.0 b	0.0 b
Nontreated		1.0 a	0.9 a	0.9 a

^x Means followed by the same letter are not significantly different, P = 0.05, Duncan's New MRT.

^yRatings were made using a 0-1 scale where 0 = no leafminer activity and 1 = leafminer activity present.

^z Aloft and Hachi-Hachi applied foliar on May 18, MBI 203 applied foliar on May 18, 23, 31 and Jun 10; Xytect and Safari applied as soil drench on Apr 5 and May 6, respectively.

Comparative Efficacy on Sri Lankan Weevil (Mylocerus undatus)

The Sri Lankan weevil (*Mylocerus undatus*) is a native to southern India, Sri Lanka and Pakistan. The species has been reported in Florida on numerous ornamental plants and fruit crops. Adults feed on leaves potentially reducing the quality and quantity of ornamentals and fruit production. Damage can range from notching on the leaf margins in an irregular pattern to much more extensive feeding along the leaf veins. Small plants and young trees usually need protection. Oviposition occurs in soil close to roots. A single female lays on an average of 360 eggs over a period of 24 days. Larvae feed on roots, however, extent of root damage is unknown. Pupation occurs in soil inside the earthen cells and take about one week. Life cycle is usually completed in 6-8 weeks. This is considered a pest of quarantine significance.

In 2016, Dale compared efficacy of several products applied foliar on Sri Lankan weevil feeding on hibiscus (Table 59 - Table 61). Three days after treatment, the industry standard TriStar, as well as BeetleGONE and MBI-203 had significantly fewer beetles per plant than the control. At 7 DAT, Tristar and BeetleGONE had four times fewer beetles per plant than the Nontreated control. Although not statistically different, MBI-203 and TriStar maintained the lowest beetle abundance per plant through 28 DAT. Although not statistically different, by the conclusion of the study, average percent plant chewing damage was less than the control for all treatments except BotaniGard and Hachi-Hachi 3.28. Among treated plants, those treated with Xpectro and BotaniGard had the highest percentage of leaves with greater than 1% damage. Those treated with Acelepryn and MBI-203 had the fewest leaves with at least 1% damage ($P=0.0008$). At 14 DAT MBI-203 and Acelepryn still had significantly fewer leaves with at least 1% damage.

In general, Acelepryn and MBI- 203 provided the greatest efficacy through the duration of the study. TriStar, the industry standard, was also among the most efficacious products tested. Plants treated with these products had the lowest beetle abundance through 14 DAT, the fewest leaves with greater than 1% herbivory through 14 DAT, and the lowest leaf-level % chewing damage through 14 DAT. No treatments showed phytotoxic effects or had any association with plant growth.

Table 59. Efficacy of several insecticides for Sri Lankan Weevil (*Myloccerus undatus*) feeding on Hibiscus (*Hibiscus rosa-sinensis*) ‘Double Peach’ - % leaf area affected, Dale, 2016.

% Leaf Area Affected by Weevil Feeding ^x							
Treatment	Rate Per 100 Gal	Applic Dates	0 DAT	3 DAT	7 DAT	14 DAT	28 DAT
Acelepryn (chlorantraniliprole)	16 fl oz	5/31	0	2.7 ± 0.71ab	1.8 ± 0.75ab	1.3 ± 0.49	1.2 ± 0.65
AzaGuard (azadirachtin)	16 fl oz	5/31, 6/8, 6/14	0	1.3 ± 0.42abc	2.0 ± 0.77ab	2.0 ± 0.89	0.8 ± 0.48
BeetleGONE (<i>Bacillus thuringiensis galleriae</i> str) + NuFilm 17	16 lb	5/31, 6/8, 6/14	0	0.8 ± 0.31bc	1.3 ± 0.76b	1.5 ± 0.62	2.0 ± 0.68
BotaniGard (<i>Beauveria bassiana</i> GHA)	32 fl oz	5/31	0	2.3 ± 0.33ab	2.5 ± 0.43ab	1.5 ± 0.56	1.2 ± 0.79
Hachi-Hachi (tolfenpyrad)	21 fl oz	5/31	0	3.0 ± 1.20ab	2.5 ± 1.0ab	3.2 ± 1.1	1.8 ± 0.70
Hachi-Hachi (tolfenpyrad)	27 fl oz	5/31	0	2.3 ± 0.56abc	2.5 ± 0.89ab	2.2 ± 0.83	1.5 ± 0.50
IKI-3106 (cyclaniliprole)	22 fl oz	5/31	0	3.2 ± 1.10a	1.7 ± 0.67ab	1.5 ± 1.0	1.5 ± 0.72
IKI-3106 (cyclaniliprole)	27 fl oz	5/31	0	1.5 ± 0.5abc	2.8 ± 0.60ab	2.8 ± 0.65	1.0 ± 0.52
Lynx (pyrethrins)	16 fl oz	5/31, 6/8	0	1.7 ± 0.33abc	4.0 ± 0.37a	1.7 ± 0.42	1.5 ± 0.96
MBI-203 (<i>Chromobacterium subsugae</i>) + NuFilm 17	3 lb	5/31, 6/8, 6/14	0	1 ± 0.36bc	2.5 ± 0.85ab	1.8 ± 0.60	0.5 ± 0.34
Preferal (<i>Isaria fumosoroseus</i>)	1 lb	5/31	0	2.5 ± 0.67ab	2.3 ± 0.76ab	2.2 ± 0.75	1.3 ± 0.42
TriStar (acetamiprid)	8 oz	5/31	0	0.7 ± 0.21c	0.7 ± 0.21b	1.3 ± 0.49	0.7 ± 0.33
Venerate (<i>Burkholderia rinojensis</i> A396) + NuFilm 17	1 gallon	5/31, 6/8, 6/14	0	1.2 ± 0.54abc	3.0 ± 0.86ab	1.8 ± 0.70	0.6 ± 0.49
Xpectro (Pyrethrins + <i>Beauveria bassiana</i> GHA)	32 oz	5/31, 6/8	0	2.0 ± 0.89abc	2.8 ± 1.05ab	2.0 ± 0.52	2.2 ± 0.79
Nontreated			0	3.0 ± 0.77a	4.0 ± 0.97a	3.3 ± 1.2	1.8 ± 0.48
		F14,70	-	2.08	2.08	0.65	1.50
		P	-	0.0233	0.0440	0.7629	0.1676
^x Means followed by the same letter are not significantly different based on Duncan-Waller's test. Data were log (x+1) transformed and analyzed with one-way ANOVA.							

Table 60. Efficacy of several insecticides for Sri Lankan Weevil (*Myloccerus undatus*) feeding on Hibiscus (*Hibiscus rosa-sinensis*) ‘Double Peach’ - % herbivory per 20 leaves, Dale, 2016.

Mean percent herbivory per 20 leaves ^x							
Treatment	Rate Per 100 Gal	Applic Dates	0 DAT	3 DAT	7 DAT	14 DAT	28 DAT
Acelepryn (chlorantraniliprole)	16 fl oz	5/31	0.0 ± 0.04	0.5 ± 0.35	0.3 ± 0.28	0.7 ± 0.22	1.1 ± 0.28
AzaGuard (azadirachtin)	16 fl oz	5/31, 6/8, 6/14	0.0 ± 0.02	0.37 ± 0.12	2.0 ± 0.68	2.1 ± 0.75	1.8 ± 0.34
BeetleGONE (<i>Bacillus thuringiensis galleriae</i> str) + NuFilm 17	16 lb	5/31, 6/8, 6/14	0.0 ± 0.02	1.1 ± 0.46	0.2 ± 0.08	1.1 ± 0.48	1.3 ± 0.24
BotaniGard (<i>Beauveria bassiana</i> GHA)	32 fl oz	5/31	0.7 ± 0.69	1.8 ± 1.06	1.6 ± 0.79	1.3 ± 0.33	1.5 ± 0.45
Hachi-Hachi (tolfenpyrad)	21 fl oz	5/31	0.2 ± 0.2	0.8 ± 0.46	1.2 ± 0.65	1.6 ± 0.52	1.9 ± 0.44
Hachi-Hachi (tolfenpyrad)	27 fl oz	5/31	0.1 ± 0.04	0.3 ± 0.17	1.3 ± 0.62	1.9 ± 0.91	1.5 ± 0.29
IKI-3106 (cyclaniliprole)	22 fl oz	5/31	0.2 ± 0.10	0.6 ± 0.26	0.7 ± 0.25	0.9 ± 0.33	2.1 ± 0.44
IKI-3106 (cyclaniliprole)	27 fl oz	5/31	0.1 ± 0.12	1.3 ± 0.73	1.1 ± 0.39	2.6 ± 0.83	2.4 ± 0.76
Lynx (pyrethrins)	16 fl oz	5/31, 6/8	0.7 ± 0.59	0.5 ± 0.22	1.0 ± 0.48	1.3 ± 0.38	3.1 ± 0.92
MBI-203 (<i>Chromobacterium subsugae</i>) + NuFilm 17	3 lb	5/31, 6/8, 6/14	0 ± 0.01	0.7 ± 0.36	0.1 ± 0.07	0.8 ± 0.23	1.7 ± 0.41
Preferal (<i>Isaria fumosoroseus</i>)	1 lb	5/31	0.0 ± 0.01	0.68 ± 0.26	0.6 ± 0.19	2.5 ± 0.92	2.0 ± 0.61
TriStar (acetamiprid)	8 oz	5/31	0.3 ± 0.23	0.5 ± 0.34	0.8 ± 0.74	1.8 ± 0.64	2.1 ± 0.46
Venerate (<i>Burkholderia rinojensis</i> A396) + NuFilm 17	1 gallon	5/31, 6/8, 6/14	0.1 ± 0.04	1.3 ± 0.39	0.6 ± 0.28	1.9 ± 0.96	1.6 ± 0.25
Xpectro (Pyrethrins + <i>Beauveria bassiana</i> GHA)	32 oz	5/31, 6/8	0.0 ± 0.02	0.54 ± 0.34	1.1 ± 0.44	1.3 ± 0.46	2.6 ± 0.27
Nontreated			0.6 ± 0.62	1.7 ± 0.86	1.8 ± 0.63	1.95 ± 0.95	1.9 ± 0.44
		F14,70	13.05	12.92	27.36	9.68	12.46
		P	0.5230	0.5329	0.0173	0.7852	0.5693
^x Due to non-normal distributions, data were analyzed using the non-parametric Wilcoxon test.							

Table 61. Efficacy of several insecticides for Sri Lankan Weevil (*Mylocerus undatus*) feeding on Hibiscus (*Hibiscus rosa-sinensis*) ‘Double Peach’ – proportion of leaves with damage, Dale, 2016.

Proportion of leaves sampled with chewing damage (>1%) ^x							
Treatment	Rate Per 100 Gal	Applic Dates	0 DAT	3 DAT	7 DAT	14 DAT	28 DAT
Acelepryn (chlorantraniliprole)	16 fl oz	5/31	0.03 ± 0.03	0.12 ± 0.07	0.03 ± 0.03c	0.10 ± 0.04d	0.17 ± 0.06
AzaGuard (azadirachtin)	16 fl oz	5/31, 6/8, 6/14	0.0 ± 0.0	0.13 ± 0.04	0.17 ± 0.03bc	0.14 ± 0.04cd	0.26 ± 0.04
BeetleGONE (<i>Bacillus thuringiensis galleriae</i> str) + NuFilm 17	16 lb	5/31, 6/8, 6/14	0.02 ± 0.03	0.18 ± 0.07	0.08 ± 0.04bc	0.14 ± 0.04cd	0.29 ± 0.05
BotaniGard (<i>Beauveria bassiana</i> GHA)	32 fl oz	5/31	0.07 ± 0.06	0.23 ± 0.11	0.20 ± 0.04b	0.28 ± 0.03ab	0.30 ± 0.05
Hachi-Hachi (tolfenpyrad)	21 fl oz	5/31	0.02 ± 0.01	0.08 ± 0.04	0.12 ± 0.03bc	0.22 ± 0.04a-d	0.33 ± 0.05
Hachi-Hachi (tolfenpyrad)	27 fl oz	5/31	0.03 ± 0.02	0.10 ± 0.05	0.12 ± 0.05bc	0.26 ± 0.05abc	0.26 ± 0.03
IKI-3106 (cyclaniliprole)	22 fl oz	5/31	0.05 ± 0.01	0.18 ± 0.08	0.10 ± 0.04bc	0.18 ± 0.04bcd	0.37 ± 0.05
IKI-3106 (cyclaniliprole)	27 fl oz	5/31	0.03 ± 0.02	0.22 ± 0.11	0.15 ± 0.06bc	0.33 ± 0.06a	0.32 ± 0.06
Lynx (pyrethrins)	16 fl oz	5/31, 6/8	0.1 ± 0.08	0.10 ± 0.04	0.18 ± 0.05bc	0.25 ± 0.04abc	0.37 ± 0.06
MBI-203 (<i>Chromobacterium subsugae</i>) + NuFilm 17	3 lb	5/31, 6/8, 6/14	0.0 ± 0.0	0.17 ± 0.08	0.07 ± 0.03bc	0.12 ± 0.02d	0.19 ± 0.04
Preferal (<i>Isaria fumosoroseus</i>)	1 lb	5/31	0.0 ± 0.0	0.12 ± 0.04	0.18 ± 0.05bc	0.28 ± 0.03ab	0.28 ± 0.06
TriStar (acetamiprid)	8 oz	5/31	0.03 ± 0.02	0.13 ± 0.08	0.08 ± 0.05bc	0.14 ± 0.05cd	0.23 ± 0.04
Venerate (<i>Burkholderia rinojensis</i> A396) + NuFilm 17	1 gallon	5/31, 6/8, 6/14	0.02 ± 0.03	0.25 ± 0.08	0.10 ± 0.04bc	0.15 ± 0.03cd	0.27 ± 0.02
Xpectro (Pyrethrins + <i>Beauveria bassiana</i> GHA)	32 oz	5/31, 6/8	0.02 ± 0.01	0.10 ± 0.08	0.20 ± 0.06b	0.18 ± 0.03bcd	0.32 ± 0.03
Nontreated			0.05 ± .01	0.20 ± .08	0.38 ± 0.08a	0.28 ± 0.06abc	0.33 ± 0.05
			F14,70	0.65	0.89	3.13	3.60
			P	0.8970	0.5740	0.0008	0.0002
^x Means followed by the same letter are not significantly different based on Duncan-Waller's test. Data were log(x+1) transformed and analyzed with one-way ANOVA.							

Results: Soil Dwelling Immatures

Comparative Efficacy on Black Vine Weevil (Otiorhynchus sulcatus)

Black vine weevil (*Otiorhynchus sulcatus*) is a serious pest of ornamental nursery crops (field and container-grown), vineyards, strawberries and hops. Even though, it is suspected the black vine weevil (BVW) originated in northern Europe, it was first identified in North America in 1835 and became a notable pest in Missouri by 1871. It is found predominantly in the northern portions of the United States, but its range extends into Virginia and out to the Pacific Northwest.

Throughout Asia, Europe, and North America, black vine weevil adults feed on the foliage and larvae feed on the roots damaging a tremendous variety of species, including azalea, strawberry, begonia, blackberry, blueberry, and cranberry, cyclamen, euonymus, forsythia, fuchsia, hemlock, impatiens, primrose, rhododendron, sedum and yew (http://www.mortonarb.org/res/CLINIC_pests_BlackVineWeevil.pdf; <http://www.entomology.umn.edu/cues/blackvw/blackvh.html>).

IR-4 sponsored nine experiments studies on BVW larvae and a single experiment on adult BVW [See Comparative Efficacy on Black Vine Weevil Adults (*Otiorhynchus sulcatus*)].

In general, preventative applications provided better control than curative rescue treatments (Table 62). The preventative applications were incorporated into the soil prior to transplanting or were drenched prior to infestation. In at least one experiment, Acelepryn, BAS 320i, Celero, Safari, and Talstar provided excellent (greater than 95%) control of black vine weevil. Curative drenches can reduce populations also, but in this series of experiments only BAS 320i provided excellent control when used in this manner.

Table 62. Summary of Black Vine Weevil (*Otiorhynchus sulcatus*) Efficacy.

Treatment	Timing	Gilrein	Reding	Reding	Reding	Cowles	Cowles	Reding	Alm	Nielsen
		Astilbe	Rhodo.	Sedum	Sedum	Strawberry	Strawberry	Yew	Yew	Yew
		2005*	2004	2004	2006	2007	2008	2004	2006	2006
		47 DAT	16 DAT	138 DAT	149 DAT	70 DAT	11 WAT	138 DAT	121 DAT	119 DAT
		45 larvae	40 larvae	4 adults	5 adults	45 eggs	0.83 larvae	4 adults	4,263 eggs	eggs
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	Pre-infestation drench				++				-	
	Pre-plant incorporation					++	-			
	Curative drench					+/-	-			+
Arena 50WDG (clothianidin)	Curative drench					-				
	Pre-plant incorporation						++			
BAS 320i EC (metaflumizone)	Pre-plant incorporation					++	++			
	Curative drench					++	++			
BotaniGard ES (Beauveria bassiana)	Curative drench						-			
	Pre-plant incorporation						-			
BotaniGard WP (Beauveria bassiana)	Curative drench						-			
	Pre-plant incorporation						-			
Celero 16WSG (clothianidin)	Pre-infestation drench				++				-	
	Curative drench									+
Discus (imidacloprid + cyfluthrin)	Curative drench					-				
Flagship 25WDG (thiamethoxam)	Curative drench					-				+
Mach 2 SC	Curative drench									-
Marathon 1G (imidacloprid)	Soil incorporation									-
Marathon II 2F	Curative drench	-								
<i>Metarhizium anisopliae</i>	Soil incorporation					-	-			-
	Curative drench	+/-					-			
Precise G & N (acephate)	Soil incorporation									-
Safari 2G (dinotefuran)	Pre-plant incorporation						-			

Treatment	Timing	Gilrein	Reding	Reding	Reding	Cowles	Cowles	Reding	Alm	Nielsen
		Astilbe	Rhodo.	Sedum	Sedum	Strawberry	Strawberry	Yew	Yew	Yew
		2005*	2004	2004	2006	2007	2008	2004	2006	2006
		47 DAT	16 DAT	138 DAT	149 DAT	70 DAT	11 WAT	138 DAT	121 DAT	119 DAT
		45 larvae	40 larvae	4 adults	5 adults	45 eggs	0.83 larvae	4 adults	4,263 eggs	eggs
Safari 20SG (dinotefuran)	Pre-infestation drench			++	++				-	
	Curative drench	-	-			-				+
	Foliar spray							-		
	Pre-plant incorporation						++			
Talstar 0.2G (bifenthrin)	Pre-plant incorporation					++				-
Tolfenpyrad	Pre-plant incorporation						-			

* Not an IR-4-sponsored experiment.

^z Data not conclusive.

¹ Rating Scale: ++ = clearly statistically better than Nontreated and greater than 95% control; + = statistically better than Nontreated and between 85 and 95% control; +/- statistically better than Nontreated with control between 70 and 85%; - = statistically equivalent to Nontreated and/or efficacy less than 70%.

² Where more than one rate or application type for a product was included in the experiment and each performed statistically different, the better rating is provided in this table.

Alm 2006

In 2006, Alm examined the efficacy of three products to control BVW larvae on yew. This experiment also tested these products on oriental beetle; see Comparative Efficacy on Oriental Beetle (*Anomala orientalis*) for information on this pest. Drenches of Acelepryn, Celero, and Safari were applied on Jul 11. Alm collected black vine weevil eggs placed them into the pots on twelve dates: Jul 13, Jul 20, Jul 24, Aug 1, Aug 7, Aug 9, Aug 17, Aug 23, Aug 29, Sep 6, Sep 15, and Sep 26. A total of 4,263 eggs were added to each pot. Pots were destructively sampled on Nov 9 and the number of live larvae were counted.

None of the treatments provided significant control of BVW larvae (Table 63). This could have been due to the constant, but realistic, repeat infestations after application. Another factor could be the length of residual control for these three products and that no repeat drench applications were made.

Table 63. Efficacy of several insecticides for Black Vine Weevil on Yew (*Taxus media*) ‘Nigra’, Alm, 2006.

Treatment	Rate	Mean Number Larvae per pot ¹²¹ DAT (Nov 9) ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	6.47 fl oz	91.8 ± 13.8 a
Celero 16WSG (clothianidin)	1.2 oz	164.6 ± 28.0 a
Safari 20SG (dinotefuran)	24 oz	132.4 ± 23.8 a
Nontreated		145.6 ± 27.5 a

^zMeans in the same column followed by the same letter are not significantly different, ($P = 0.05$, LSD test).

Cowles 2007

This experiment used potted strawberries as a model system to determine which insecticides would provide protection against BVW larvae. Special attention was paid to determine whether there was a dose-response for Acelepryn and BAS 320, to compare neonicotinoids, and to compare preventive preplant incorporation into potting media with curative drenches targeting 3rd instars. Strawberry daughter plants were taken from research plots at the Valley Laboratory on Nov 16, 2006. Cultivars used were 'Allstar', 'Annapolis', 'Darselect', 'Idea' (two replicates), and 'Jewel'. Black vine weevil eggs were obtained from a colony at the Valley Laboratory. About 200 adults were enclosed in a 20 L plastic bucket with yew foliage, and eggs were collected by shaking foliage on a weekly basis. Pots were repeatedly infested in January and early February until there were 45 eggs added per pot. Eggs were placed 1 – 2 cm deep in the soil, close to the crown of the plant. Larvae were counted on Apr 16 and 17 by sifting through the potting media shaken from the strawberry root system.

All preplant potting mix incorporation treatments, including the positive control standard Talstar 0.2G, were completely effective (Table 64). All tested concentrations of BAS 320i caused complete mortality. Although the Acelepryn pretreatment was completely effective, only the higher dosage of this product applied as an early curative drench suppressed the BVW population. Neonicotinoid products performed poorly against BVW larvae as early curative drench treatments. Results with *Metarhizium anisopliae* were disappointing. Cool temperatures and poor quality inoculum may have contributed to its ineffectiveness in this experiment.

Table 64. Efficacy of several insecticides for Black Vine Weevil on Strawberry (*Fragaria* sp.), Cowles, 2007.

Treatment	Application Method	Application Date	Rate	Active Ingredient (mg/pot)^z	Number of Larvae 10 weeks after last infestation^{y, x}	Percent Control
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	Curative drench	Mar 15	0.8 fl oz/100 gal	0.873	4.8 a	0
			6.5 fl oz/100 gal	7.07	0.8 cd	75
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	Pre-plant incorporation	Nov 10	5 ppm	3.02	0.0 d	100
			10 ppm	6.03	0.0 d	100
			20 ppm	12.1	0.0 d	100
Arena 50WDG (clothianidin)	Curative drench	Mar 15	1.28 oz/100 gal	4.79	5.4 a	0
BAS 320i SC (metaflumizone)	Curative drench	Mar 15	50 ppm	30.2	0.0 d	100
			100 ppm	60.4	0.0 d	100
BAS 320i EC (metaflumizone)	Pre-plant incorporation	Nov 10	25 ppm	15.1	0.0 d	100
			50 ppm	30.2	0.0 d	100
			100 ppm	60.4	0.0 d	100
			200 ppm	121	0.0 d	100
Discus (imidacloprid + cyfluthrin)	Curative drench	Mar 15	13 fl oz/100 gal	1.0 + 0.239	4.2 ab	0
Flagship 25WDG (thiamethoxam)	Curative drench	Mar 15	8 oz/100 gal	0.5	3.8 ab	0
<i>Metarhizium anisopliae</i>	Soil incorporation	Jan 9	4.5×10^8 spores/L	28.3	4.8 a	0
Safari 20SG (dinotefuran)	Curative drench	Mar 15	24 oz/100 gal	27.0	1.0 bcd	69
Talstar 0.2G (bifenthrin)	Pre-plant incorporation	Nov 10	10 ppm	6.03	0.0 d	100
Nontreated					3.2 abc	0

^zExpressed in µl per pot. This amounts to 1.00 mg of imidacloprid and 0.239 mg cyfluthrin per pot.

^y Pots were infested repeatedly from Jan 9 to Feb 6 until there were 45 eggs added per pot.

^x Means followed by the same letter do not significantly differ (Fisher's protected LSD test, $P < 0.05$).

Cowles 2008

Cowles conducted a second experiment with potted strawberries to continue investigating dose-response of Acelepryn and BAS 320i, to compare neonicotinoids (Safari vs. Arena), and to compare preventive preplant incorporation into potting media with curative drenches.

Black vine weevil eggs were repeatedly infested to potted strawberries on June 5, 12 and 19, for a total of 123 eggs per pot. Eggs were placed 1 – 2 cm deep in the soil, close to the crown of the plant. Larvae were counted on September 4, 5 and 8 by sifting through the potting media shaken from the strawberry root system. Root systems were rated on a 0-4 scale, where 0 indicated total destruction, always involving crown feeding and imminent plant death; 1 was poor roots, in which the living roots were found within a 50 ml volume of medium; 2 was fair, with most of the soil falling away from the roots; 3 was good, with most of the soil being held together by the root system; 4 was excellent, with roots encircling the bottom of the pot and great difficulty in removing soil from the extensive root system.

Recovery of BVW larvae was unusually low, with the Nontreated Check having one of the lowest numbers and the most extensive root system (Table 65). Two factors other than insecticide treatment probably contributed to poor recovery of larvae: ant predation (maximized by running the trial during the summer) and over-exploitation of food resources by BVW larvae in some treatments. Therefore, the results from this experiment have to be interpreted from the combination of the number of larvae and the root ratings. Based on these data, BAS 320i, applied as preventive preplant incorporation into potting media or curative drenches, should be considered an outstanding material to target control of BVW. Acelepryn looked ineffective at the lower rates used this year compared to those in the 2007 study. Arena was effective applied as a preplant incorporation; in 2007, it was ineffective when applied as a curative drench. Safari 20SG was effective in this year's trial with preplant incorporation while Safari 2G was ineffective. The poorer control from the Safari 2G granules suggests that the product distribution within the mix was inadequate.

BotaniGard ES and WP formulations require further testing before concluding that they have poor potential for control of BVW. It is possible that the dosage applied was inadequate. *Metarhizium*- and Tolfenpyrad-treated pots had uniformly healthy plants and few larvae, however the low counts in the Nontreated check preclude being able to determine whether these treatments provided significant benefits.

Table 65. Efficacy of several insecticides for Black Vine Weevil on Strawberry (*Fragaria* sp.), Cowles, 2008.

Treatment	Application Method	Application Date	Rate	Active Ingredient (mg/pot)	Number of Larvae 11 weeks after last infestation^{z, y} (% control)	Root Rating^{z, x} 11 WAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	Curative drench	Jul 15	0.8 fl oz/100 gal	0.44	4.5 bc (0)	2.8 ab
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	Pre-plant incorporation	Apr 24	1 ppm	0.22	3.8 bcd (0)	2.7 bc
			2 ppm	0.43	10.8 a (0)	1.0 d
			4 ppm	0.86	0.67 e (20)	4.0 a
Arena 50WDG (clothianidin)	Pre-plant incorporation	Apr 24	49 mg/pot	24.7	0 e (100)	4.0 a
BAS 320i SC (metaflumizone)	Curative drench	Jul 15, 29, Aug 13, 28	16 fl oz/100 gal	240	0 e (100)	4.0 a
BAS 320i EC (metaflumizone)	Pre-plant incorporation	Apr 24	1 ppm	0.22	0.50 e (40)	4.0 a
			2 ppm	0.43	0.33 e (60)	4.0 a
			4 ppm	0.86	0 e (100)	4.0 a
BotaniGard ES (<i>Beauveria bassiana</i>)	Curative drench	Jul 15, 29, Aug 13, 28	39 µL/pot		4.2 bc (0)	1.3 d
BotaniGard WP (<i>Beauveria bassiana</i>)	Curative drench	Jul 15, 29, Aug 13, 28	18.7 mg/pot		1.2 de (0)	0.5 d
BotaniGard ES (<i>Beauveria bassiana</i>)	Pre-plant incorporation	Apr 24	39 µL/pot		2.0 cde (0)	1.3 d
BotaniGard WP (<i>Beauveria bassiana</i>)	Pre-plant incorporation	Apr 24	18.7 mg/pot		5.8 b (0)	1.5 d
<i>Metarhizium anisopliae</i>	Curative drench	Jul 15	2.9 g/pot		0.67 e (20)	4.0 a
<i>Metarhizium anisopliae</i>	Pre-plant incorporation	Apr 24	2.9 g/pot		0.33 e (60)	4.0 a
Safari 2G (dinotefuran)	Pre-plant incorporation	Apr 24	1.23 g/pot	24.7	1.7 cde (0)	2.7 bc
Safari 20SG (dinotefuran)	Pre-plant incorporation	Apr 24	24 oz/100 gal	24.7	0 e (100)	4.0 a
Tolfenpyrad	Pre-plant incorporation	Apr 24	10 ppm	2.2	0.83 e (0)	3.8 ab
Nontreated					0.83 e (0)	4.0 a

^z Pots were infested repeatedly from Jun 5, 12 and 19 until there were 123 eggs added per pot.

^y Means followed by the same letter do not significantly differ (Fisher's protected LSD test, P < 0.05).

^x Root rating 0-4, where 0 = all roots destroyed, 4 = extensive root system.

Gilrein 2005

In 2005, Gilrein tested whether Marathon II, *Metarhizium anisopliae* and Safari as curative applications would control BVW on astilbe 'Pink Sprite'. Laboratory reared black vine weevil larvae were placed into pots on Sept 10, Oct 5, and Oct 26. Treatments were drenched on Nov 5 using 120 ml solution per pot. ANOVA and multiple comparisons among treatment means were performed using a statistical multiple comparison procedure (SuperAnova v. 1.1, Abacus Concepts).

Only *Metarhizium anisopliae* provided both statistically and biologically significant levels of control 47 days after treatment (Table 66). Applications to early instars may have improved control so early curative or preventative applications could have reduced populations to a greater extent.

Table 66. Efficacy of several insecticides for curative control of Black Vine Weevil on Astilbe (*Astilbe simplicifolia*) 'Pink Sprite', Gilrein, 2005*.

Treatment	Rate	Live Grubs per pot 47 DAT (% control) ^z
Marathon II 2F	20 g/650 pots	9.6 bc (18.6%)
<i>Metarhizium anisopliae</i> (Strain F52)	14.04 cfu/pot	2.0 a (83.1%)
	28.08 cfu/pot	2.6 a (78.0%)
	56.16 cfu/pot	3.5 a (70.3%)
Safari 20SG (dinotefuran)	12 oz / 100 gal	9.0 b (23.7%)
	24 oz / 100 gal	8.7 b (26.3%)
Nontreated	--	11.8 c (0.0%)

* Not an IR-4 experiment.

^zMeans within columns followed by the same letter are not significantly different at p=0.05 (Fisher's LSD).

Nielsen 2006

Nielsen tested nine products for their control of BVW larvae on yew. Treatments consisted of preventative soil mixtures or curative top dressings and drenches approximately 3 weeks before or 2 weeks after infestations, respectively. Incubated, BVW eggs were obtained from weevils collected from a taxus nursery and maintained in a rearing room on taxus foliage. Plants were maintained outdoors under standard nursery management conditions throughout the growing season. Phytotoxicity was assessed 7 & 14 days after treatment, periodically after that and at the time of efficacy evaluation, when roots were evaluated for phytotoxicity.

Larval establishment in Nontreated check containers was marginal but sufficient to compare treatment effects. Marathon and Talstar, the industry standards, *Metarhizium*, Mach 2 and Precise failed in this trial (Table 67). Acelepryn, Celero, and Safari drenches prevented establishment of black vine weevil larvae: No larvae were found in any containers that received these treatments. The Flagship drench was nearly as effective.

No phytotoxicity was observed.

Table 67. Efficacy of several insecticides for Black Vine Weevil on Yew (*Taxus media densiformis*), Nielsen, 2006.

Treatment	Rate	Application Method	Number of larvae per pot 12/6 17 WAT ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	16 oz/100 gal	8 oz drench per pot	0.0 a
Celero 16WSG (clothianidin)	20 oz/100 gal	8 oz drench per pot	0.0 a
Flagship 25WG (thiamethoxam)	8 oz/100 gal	8 oz drench per pot	0.2 a
Mach 2 SC	2 lb ai/Acre	8 oz drench per pot	8.2 b
Marathon 1G (imidacloprid)	0.1 g ai/gal	Premixed into soil media	3.0 ab
<i>Metarhizium</i>	6.25 grams/pot	Premixed into soil media	9.4 b
Precise G & N (acephate)	6 g product/can	Premixed into soil media	6.8 ab
Safari 20 SG (dinotefuran)	24 oz/100 gal	8 oz drench per pot	0.0 a
Talstar 0.2 G (bifenthrin)	25 ppm	Premixed into soil media	4.8 b
Nontreated			4.2 ab

^z Means within columns followed by the same letter are not significantly different at p=0.05 (Fisher-Hayter LSD).

Reding 2004

In 2004, Reding conducted a series of experiments testing Safari as a tool to manage black vine weevil.

For the first experiment, Rhododendron plants were purchased as 6" plants and transplanted into 2 gallon pots on Apr 23. The containers were then infested with BVW larvae (40 larvae per pot). Because the larvae were mature and many had begun to pupate by the time the drench was applied, the treatments were considered curative, rescue applications. The containers were put outdoors in a container nursery and irrigated when needed with a drip irrigation system. On Apr 28, pots were drenched with 240 ml of solution per each 2 gallon pot. On May 14 (16 DAT), the containers were dumped and the soil sifted for black vine weevils. There was no biologically significant impact on BVW larvae with drenches on late instars (Table 68).

For the second experiment, sedum plants were purchased as 4" bare root plants and transplanted into 2 gallon pots on Apr 15. The drench treatment was applied on Jun 2 and four hours later each container was infested with 4 adult black vine weevils and caged to prevent the insects from leaving. The containers were put outdoors in a container nursery and irrigated as needed with a drip irrigation system. Plants were evaluated for feeding damage on three dates (Table 69). To determine efficacy, the containers were dumped on Oct 18 and the roots and soil carefully examined for BVW larvae. Many of the larvae found were within the roots. The plants were destroyed in the process of examination. All Safari drench applications significantly reduced feeding damage and the number of larvae found in Sedum roots.

For the third experiment, yew plants were purchased as 6" bare root plants and transplanted into 2 gallon pots on May 7. The foliar spray was applied on Jun 2 and four hours later each container was infested with 4 adult black vine weevils and caged to prevent the insects from leaving. The containers were put outdoors in a container nursery and irrigated as needed with a drip irrigation system. Plants were evaluated for feeding damage on three dates. To determine efficacy, the containers were dumped on Oct 18 and the roots and soil carefully examined for BVW larvae. Many of the larvae found were within the roots. The plants were destroyed in the process of examination. There were no significant differences in the number of larvae or in feeding damage (Table 70).

There was no phytotoxicity on any of the treated plants.

Table 68. Efficacy of Safari 20SG for Black Vine Weevil on Rhododendron (*Rhododendron* sp.) ‘Nova Zembla’, Reding, 2004.

Treatment	Rate Per 100 Gal ^z	Insect Counts 16 DAT ^y
<i>Late Instar Larvae</i>		
Safari 20SG (dinotefuran)	12 oz	3.8
Safari 20SG (dinotefuran)	24 oz	3.8
Safari 20SG (dinotefuran)	48 oz	2.8
Nontreated		3.8
<i>Pupae</i>		
Safari 20SG (dinotefuran)	12 oz	14.3
Safari 20SG (dinotefuran)	24 oz	10.6
Safari 20SG (dinotefuran)	48 oz	7.0
Nontreated		14.3
<i>Adults</i>		
Safari 20SG (dinotefuran)	12 oz	1.5
Safari 20SG (dinotefuran)	24 oz	2.0
Safari 20SG (dinotefuran)	48 oz	5.2
Nontreated		3.8
<i>Total</i>		
Safari 20SG (dinotefuran)	12 oz	19.7
Safari 20SG (dinotefuran)	24 oz	15.5
Safari 20SG (dinotefuran)	48 oz	15.5
Nontreated		22.0

^z Treatments were applied as 240 ml of solution to each 10" (2 gallon) pot on April 23, 2004.

^y There were no statistical differences according to ANOVA and Fisher-Hayter LSD.

Table 69. Efficacy of Safari 20SG drenches for Black Vine Weevil on Sedum (*Sedum* sp.) ‘Vera Jameson’, Reding, 2004.

Treatment	Rate Per 100 Gal ^z	Feeding damage ^y			Number of live larvae 10/18 138 DAT
		6/22/04 20 DAT	7/27/04 57 DAT	8/20/04 79 DAT	
Safari 20SG (dinotefuran)	12 oz	1.5 a	1.0 a	1.3 a	0.0 a
Safari 20SG (dinotefuran)	24 oz	0.6 a	0.5 a	0.3 a	0.0 a
Safari 20SG (dinotefuran)	48 oz	0.6 a	0.3 a	0.1 a	0.5 a
Nontreated		8.8 b	10.0 b	6.8 b	8.3 b

^z Treatments were applied as 240 ml of solution to each 10" (2 gallon) pot on June 2, 2004.

^y Means within columns followed by the same letter are not significantly different at p=0.05 (Fisher-Hayter LSD).

Table 70. Efficacy of Safari 20SG drenches for Black Vine Weevil on Yew (*Taxus* sp.) ‘Brownii’, Reding, 2004.

Treatment	Rate Per 100 Gal ^z	Feeding damage			Number of live larvae 10/18 138 DAT
		20 DAT ^y	57 DAT	79 DAT	
Safari 20SG (dinotefuran)	12 oz	2.3	3.0	2.0	4.0
Safari 20SG (dinotefuran)	24 oz	0.5	0.4	0.3	0.5
Safari 20SG (dinotefuran)	48 oz	1.0	1.3	1.0	3.8
Nontreated		2.5	2.8	1.5	5.5

^z Treatments were applied as 240 ml of solution to each 10" (2 gallon) pot on June 2, 2004.

^y There were no statistical differences according to ANOVA and Fisher-Hayter LSD.

Reding 2006

Sedum spurium 'Vera Jameson' plants, grown in one gallon containers, were used to test the efficacy of soil drench treatments of various products against BVW. All treatment rates were applied in 200 ml of solution as a surface drench poured over the potting media, on Jun 5. On Jun 6, four adult BVW were placed on each caged plant. One additional BVW adult was added to each cage 2 weeks later.

To determine any differences in insecticidal activity on adults and larvae, efficacy was measured by rating feeding damage on foliage by adults on Aug 8 (% of total damaged leaves and number of remaining blossoms per plant) and number of larvae found in the roots, on Nov 2. There was a significant difference in percentage of leaves fed on and number of flower blossoms on Nontreated and Acelepryn treated plants compared to Celero and Safari treated plants (Table 71). There was also a significant difference in numbers of larvae recovered from Nontreated plants compared to all 3 insecticide treatments

Table 71. Efficacy of several insecticides for Black Vine Weevil on Sedum (*Sedum spurium*) ‘Vera Jameson’, Reding, 2006.

Treatment	Rate Per 100 Gal ^z	Percentage of leaves with feeding damage ^y	Mean # of Sedum Blooms	Mean number of larvae per plant 149 DAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	47.9 oz	93.2 a	2.5 a	1.0 b
Celero 16WSG (clothianidin)	4 oz	21.5b	11.8b	0.0b
Safari 20SG (dinotefuran)	24 oz	22.0b	9.8b	0.0b
Nontreated		93.2a	1.8a	12.2a

^z Each treatment was drenched onto soil using 200 ml per plant.

^y Means within columns followed by the same letter are not significantly different ANOVA ($P = 0.05$), means separated by LSD ($\alpha = 0.05$).

Comparative Efficacy on Japanese Beetle Grubs (*Popillia japonica*)

The Japanese beetle (*Popillia japonica*) is a widespread and destructive exotic pest of turf, landscape, and ornamental plants in the United States. Outside of its native Japan, it is also found in China, Russia, Portugal, and Canada. Since the first detection in the US in a nursery near Riverton, New Jersey in 1916, it has spread to many states east of the Mississippi River, as well as parts of Wisconsin, Minnesota, Iowa, Missouri, Nebraska, Kansas, Arkansas and Oklahoma. Despite regulatory efforts, it has become established in at least 30 states by 2002. Occasional introductions are made into western states such as California and Oregon when the adult beetles or larvae are shipped in commerce.

The Japanese beetle (JB) has a total host range of more than 400 plant species, including turf, ornamentals, fruits, and vegetables. Currently the Japanese beetle is the most widespread pest of turf and costs the turf and ornamental industry approximately \$450 million each year in management alone (<http://ohioline.osu.edu/hyg-fact/2000/2504.html>, <http://edis.ifas.ufl.edu/IN630>).

IR-4 sponsored a couple experiments on JB larvae and several experiments on the adults [See Comparative Efficacy on Japanese Beetle Adults (*Popillia japonica*)].

Braman 2006

In 2006, Braman tested seven products as soil applications for efficacy on JB grubs infesting Caldwell Pink roses. Pots were treated on Jun 13 with drenches or surface application of granular products. On Aug 2 (50 DAT), five replications of the 8 treatments were infested with 5 second instar grubs recovered from turf plot cages used to enable adult beetles to lay eggs and larvae to grow. Pots were destructively sampled on Aug 16 (64 DAT) and evaluated for surviving larvae. Data were subjected to ANOVA using the GLM procedure of SAS.

Recovery in control pots was 92% of the original larvae used to infest the pots. Although mortality ranged from 8 to 60%, differences among treatments were not significant (Table 72). Several factors could have contributed to the lack of mortality observed including evaluating populations at 2 weeks instead of 4 weeks post infestation and infesting 5 weeks after application date

Braman 2007

In the experiment conducted in 2007, Braman examined 5 products for efficacy on JB grubs infesting rose. Caldwell Pink roses, grown in 3 gallon pots, were treated (drenched) Jul 17. Design was a randomized complete block with 5 replications. Ten, first instar grubs were inserted into pots by inserting a pencil into the pots and dropping in the larvae on Jul 19. Pots were destructively sampled on Oct 5 (80 DAT) and the number of grubs were recorded. Data were analyzed using the GLM procedure of SAS and mean separation was by LSD.

Very few grubs survived in the pots. Differences were only significant at the 10% level and all treatments were equivalent to the UTC (Table 73).

Table 72. Efficacy of several insecticides for *Popillia japonica* on rose (*Rosa* sp.) ‘Caldwell Pink’, Braman, 2006.

Treatment	Rate Per 100 Gal	Number of Grubs per Pot 2 weeks after infestation 64 DAT ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 oz, Apply 1 pint per 2.25 sq ft	3.8 a
Celero 16WSG (clothianidin)	0.5 oz, Apply 1 pint per 2.25 sq ft	3.4 a
Flagship 25WG (thiamethoxam)	5 oz	3.6 a
Precise (acephate)	3 tsp/pot	2.8 a
Safari 20 SG (dinotefuran)	24 oz	3.6 a
Scimitar (lambda-cyhalothrin)	60 oz	2.0 a
Tristar 70 WSP (acetamipid)	8 oz	2.8 a
Nontreated		4.6 a

^z Means followed by the same letter are not significantly different, P> 0.05

Table 73. Efficacy of several insecticides for *Popillia japonica* on rose (*Rosa* sp.) ‘Caldwell Pink’, Braman, 2007.

Treatment	Rate Per 100 Gal	Drench Volume per pot	Number of Grubs per Pot 80 DAT ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 oz	100 ml	0.80 a
BAS 320i (metaflumizone)	16 oz	400 ml	1.60 a
Celero 16/WSG (clothianidin)	16 oz	500 ml	0.80 a
Met 52 (<i>Metarhizium anisopliae</i>)	58 oz	400 ml	1.00 a
Talstar (bifenthrin)	80 oz	400 ml	0.80 a
Nontreated			1.00 a

^z Means followed by the same letter are not significantly different, P> 0.05

Comparative Efficacy on May/June Beetle Grubs (Phyllophaga spp.)

True white grubs are the larvae of May beetles (also called June Beetles) found in the genus *Phyllophaga*, of which there are over 100 different species. *Phyllophaga* spp. and related insects are distributed throughout the United States and Canada. Larvae can be very damaging to the roots of ornamental plants grown in nurseries.

In 2008, Buss evaluated 3 products as soil applications on live oak trees that were severely infested with white grubs (primarily *Phyllophaga* spp.). The 40-inch diameter root ball of each tree served as a “plot”. Roots had been cut several months before the test, and the balls had been placed in burlap-wrapped wire baskets with frequent irrigation. The treatments were applied on May 19. Liquid treatments were applied to the surface area beginning at the outside edge of the ball to the trunk, then, applying similarly on four sides of the tree (90° perpendicular to the one previously administered). Trees that received the CoreTect treatment had 12 tablets each, inserted 4 inches deep, spaced 5 inches apart within the root ball. About 0.17 inch of post-treatment irrigation was immediately applied. Destructive sampling evaluation was conducted (Aug 29, Sep 2-4). The entire basket of each tree was lifted out of the ground, the burlap was removed from the root ball, and the entire root ball was sifted by hand for all grubs.

The two higher rates of Acelepryn reduced the number of May/June beetle and other grubs compared to the Nontreated Control, and were comparable to the standard Merit 75WP in efficacy (Table 74). Although CoreTect tablet provided the best efficacy, it is not a good formulation for nursery work, given the labor-intensive method of placing the tablets into the soil and the large number of trees that would need to be treated.

Table 74. Efficacy of several insecticides for May/June Beetle Grubs (*Phyllophaga* spp.) on Live Oak (*Quercus virginiana*) ‘Highrise’, Buss, 2008*.

Treatment	Rate Per Acre^z	Mean #. of <i>Phyllophaga</i> grubs per tree^y 15 WAT	Mean # of all grub species^x per tree 15 WAT
Acelepryn 1.67SC (chlorantraniliprole)	8 fl oz	181.0 b	203.2 bc
Acelepryn 1.67SC (chlorantraniliprole)	12 fl oz	87.8 ab	113.4 ab
Acelepryn 1.67SC (chlorantraniliprole)	16 fl oz	132.0 ab	159.4 ab
CoreTect (imidacloprid)	3 tablets per inch trunk diam.	55.8 a	76.2 a
Merit 75WP (imidacloprid)	6.4 oz	119.0 ab	139.8 ab
Nontreated		293.8 c	315.4 c

* Not an IR-4 experiment.

^z Each treatment was applied onto soil on May 19.

^y Means within columns followed by the same letter are not significantly different (Tukey’s HSD).

^x Species identified were May/June beetles (*Phyllophaga* spp.), flower beetle (*Euphoria sepulcralis*) and ox beetle (*Strategus antaeus*).

Comparative Efficacy on Oriental Beetle (*Anomala orientalis*)

The oriental beetle is one of the important soil pests of nursery ornamental plants and turf grasses. Four researchers examined the efficacy of 14 products and unregistered materials for managing oriental beetle grubs (Table 75). The products tested included Acelepryn, BAS 320i, Celero 16 WSG, Discus, Dylox, Flagship, Marathon, Safari 20SG, Talstar and Tick-EX. In general, treatments were more effective when applied to less mature grubs. In tests targeted to newly-hatched and 2nd instar larvae, Acelepryn, BAS 320i, Flagship and Safari generally provided good control but were ineffective in two trials targeted to 3rd instar larvae. In a Freiburger 2005 trial on natural infestation of oriental beetle, products provided more reduction of grubs when applied in summer compared to fall application. These results suggest that growers and landscapers should target applications at or immediately after peak oriental beetle mating flights.

Table 75. Summary of Oriental Beetle Grub (*Anomala orientalis*) Efficacy.

Treatment	Alm 2008	Alm 2006	Freiberger 2005		Freiberger 2009	Gilrein 2005	Gilrein 2007	Reding 2009	
	Rhododendron	Yew	Arborvitae	Holly	Arborvitae	Lawn Grass	Lawn Grass	Lilac	White Oak
	17 WAT	20 WAT				53 DAT	30 DAT	14 WAT	22 WAT
	20 larvae, 2 nd instar	20 larvae, 2 nd instar	Natural infestation	Natural infestation	Natural infestation	10 larvae, 3 rd instar	10 larvae, 3 rd instar	20 eggs	20 eggs
	Container	Container	In Ground	In Ground	In Ground	Container	Container	Container	Container
	Preventative	Preventative	Preventative	Preventative	Preventative	Curative	Curative	Early Curative	Early Curative
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	+	++	-	+/-	+	-	-	++	++
BAS 320i (metaflumizone)	++				-		-		
Cal-Agri 50 1 % (potassium phosphate)						-			
Celero 16WSG (clothianidin)		+/-	-	-		+			
Discus (imidacloprid + cyfluthrin)			-	+/-					
Dylox 80S (trichlorfon)							+		
Flagship (thiamethoxam)			-	+/-	+		-	++	++
Hexacide (rosemary oil)							-		
Marathon II 2F (imidacloprid)					+	+			
Safari 20SG (dinotefuran)	++	++			-		-	++	++
Talstar G (bifenthrin)	++								
Met52 (<i>Metarhizium anisopliae</i>)	-					-			
Sevin XLR 4F (carbaryl)						-			
Tick-EX (<i>Metarhizium anisopliae</i>)							-		

¹ Rating Scale: ++ = clearly statistically better than Nontreated and greater than 95% control; + = statistically better than Nontreated and between 85 and 95% control; +/- statistically better than Nontreated with control between 70 and 85%; - = statistically equivalent to Nontreated and/or efficacy less than 70%.

² Where more than one rate or application type for a product was included in the experiment and each performed statistically different, the better rating is provided in this table.

Alm 2006

As part of a combined study examining Acelepryn, Celero, and Safari for black vine weevil and oriental beetle control, Alm collected 500 second instar oriental beetle larvae from turf and placed 20 larvae on the surface of each pot containing yew 'Nigra' on Aug 31. Drenches of Acelepryn, Celero, and Safari were applied as preventative treatments on Jul 11. See Comparative Efficacy on Black Vine Weevil (*Otiorhynchus sulcatus*) for the results on black vine weevil.

Pots were destructively harvested 20 weeks after application. All three products provided significant control of oriental beetle larvae (Table 76).

No phytotoxicity was observed.

Table 76. Efficacy of several insecticides for Oriental Beetle grubs on Yew (*Taxus media*) 'Nigra', Alm, 2006.

Treatment	Rate Per 100 Gal ^z	Mean Number Larvae per pot 20 WAT (Nov 9) ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	6.47 fl oz	0.0 ± 0.0 a
Celero 16WSG (clothianidin)	1.2 oz	1.2 ± 0.6 a
Safari 20SG (dinotefuran)	24 oz	0.0 ± 0.0 a
Nontreated		7.6 ± 1.1 b

^z Means in the same column followed by the same letter are not significantly different, ($P = 0.05$, LSD test).

Alm 2008

In 2008 Alm evaluated several insecticides applied as preventative treatments for oriental beetle control.

Acelepryn, BAS 320i, and Safari were applied as drenches while Met52G and Talstar were incorporated with the potting mix on Jul 25 prior to planting. Alm collected 720 second instar oriental beetle larvae from turf and placed 20 larvae on the surface of each pot containing rhododendron 'Scintillation' on Aug 25.

BAS 320i, Safari 20 SG and Talstar Nursery Granular treatments provided 100% control of oriental beetle larvae (Table 77); Acelepryn also provided excellent control of larvae and was not significantly different from these treatments. Met 52 G was not significantly different from the Nontreated pots.

Table 77. Efficacy of several insecticides for Oriental Beetle grubs on Rhododendron (*Rhododendron* sp.) 'Scintillation', Alm, 2008.

Treatment	Rate Per 100 Gal	Mean Number Larvae per pot 17 WAT (Nov 5) ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 fl oz	1.7 ± 0.3b
BAS 320i (metaflumizone)	16 fl oz	0.0 ± 0.0 b
Met52G (<i>Metarhizium anisopliae</i> strain F52)	20 g/4 gal media	10.2 ± 0.9a
Safari 20SG (dinotefuran)	24 oz	0.0 ± 0.0 a
Talstar Nursery Granular (bifenthrin)	33.7 g/4 gal media	0.0 ± 0.0 b
Nontreated		11.7 ± 1.4a

^z Means in the same column followed by the same letter are not significantly different, ($P = 0.05$, LSD test).

Freiberger 2005

Arborvitae 'Emerald' and Holly 'Blue Girl' were planted into field soil at Rutgers University Tree Fruit Research & Extension Center in Cream Ridge, NJ. The field where the arborvitae and holly were planted had formerly been planted with strawberries heavily infested with oriental beetle. In 2005, four products were drenched once either during summer (Aug 3) or fall (Nov 1) with 1 pint of diluted product per plant – Acelepryn (chlorantraniliprole), Celero (clothianadin), Discus (bifenthrin +imidacloprid), and Flagship (thiamethoxam). Starting Apr 24, 2006, arborvitae and holly plants were destructively harvested and the number of oriental beetle larvae in and around each root ball was counted.

Timing of the applications was critical. When the drenches occurred in the summer all products significantly reduced larvae on both crops (Table 78). However, when the drench applications occurred in the fall, only Acelepryn exhibited a statistically significant reduction in population on arborvitae; and for holly, Acelepryn, Celero, and Discus reduced oriental beetle grubs.

No phytotoxicity was observed.

Table 78. Efficacy of several insecticides for Oriental Beetle on Arborvitae (*Thuja* sp.) ‘Emerald Giant’ and holly (*Ilex* sp.) ‘Blue Girl’ – Number of Grubs, Freiburger, 2005.

Treatment ^z	Rate Per 100 Gal	Mean number of larvae per plant ^y			
		Arborvitae		Holly	
		Summer Drench	Fall Drench	Summer Drench	Fall Drench
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 fl oz	2.2 a	4.5 a	0.8 a	1.2 a
Celero 16WSG (clothianidin)	0.5 oz	2.6 ab	4.6 ab	1.7 a	1.7 a
Discus (imidacloprid + cyfluthrin)	10.0 oz	3.7 ab	5.0 ab	0.7 a	0.8 a
Flagship (thiamethoxam)	0.4 oz	4.2 b	4.9 ab	0.9 a	2.2 ab
Nontreated		6.3 c	6.3 b	3.0 b	4.0 b

^z Each plant was drenched with 1 pint of solution.

^y Means within columns followed by the same letter are not significantly different according to Fisher-Hayton (P<0.05).

Table 79. Efficacy of several insecticides for Oriental Beetle on Arborvitae (*Thuja* sp.) ‘Emerald Giant’ and holly (*Ilex* sp.) ‘Blue Girl’ – Plant Height, Freiburger, 2005.

Treatment ^z	Rate Per 100 Gal	Plant Height ^y			
		Arborvitae		Holly	
		Summer Drench	Fall Drench	Summer Drench	Fall Drench
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 fl oz	41.3 a	43.7 a	16.3 b	15.5 b
Celero 16WSG (clothianidin)	0.5 oz	43.8 a	44.0 a	17.4 b	18.0 c
Discus (imidacloprid + cyfluthrin)	10.0 oz	41.9 a	42.4 a	13.9 a	13.2 a
Flagship (thiamethoxam)	0.4 oz	43.4 a	42.6 a	14.4 a	15.4 ab
Nontreated		42.4 a	43.5 a	15.8 ab	16.2 bc

^z Each plant was drenched with 1 pint of solution.

^y Means within columns followed by the same letter are not significantly different according to Fisher-Hayton (P<0.05).

Freiburger 2009

A field known to be infested with oriental beetle at Rutgers University Tree Fruit Research & Extension Center in Cream Ridge, NJ was planted with arborvitae (*Thuja* sp.) in the spring of 2007. The established saplings were treated with drenches or granular broadcast applications during the peak adult flight in Aug 2009. The products tested included BAS 320i (metaflumizone), Acelepryn (chlorantraniliprole), Flagship 0.22G (thiamethoxam), Flagship 25WG (thiamethoxam), Marathon II (imidacloprid), Safari 2G (dinotefuran), and Safari 20SG (dinotefuran). Starting Apr 5, 2010, arborvitae were destructively harvested and the number of oriental beetle larvae in and around each root ball was counted.

The outcome of this experiment was intriguing. While some products did decrease the average number of grubs on arborvitae, the average height did not appear to be shorter for those treatments with higher grub infestations nor did the average width correlate with the average number of grubs. In other words, the variability in arborvitae height and width observed at this site was not related the number of grubs attacking the roots in this experiment. The products with the least number of oriental beetle larvae included Acelepryn, Flagship 0.22G, Flagship 25WG, and Marathon II.

No phytotoxicity was observed.

Table 80. Efficacy of Seven Products to Reduce Oriental Beetle Populations on Arborvitae, Freiburger, 2009.

Treatment	Rate	Average Number of Grubs per Tree	Average Tree Height	Average Tree Width
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 fl oz	1.1 a	56.2 a	31.7 b
BAS 320i (metaflumizone)	16 oz per 100 gal	6.6 bc	52.9 a	31.2 ab
Flagship 0.22G (thiamethoxam)	11.25 g product per 9 sq ft	2.2 a	53.1 a	30.2 ab
Flagship 25WG (thiamethoxam)	0.1 g product per 9 sq ft in ½ gal water	1.1 a	53.3 a	30.2 ab
Marathon II (imidacloprid)	0.6 fl oz per 1,000 sq ft in 10 gal water	1.2 a	53.4 a	31.5 ab
Safari 2G (dinotefuran)	120 grams per inch dbh for trees	6.2 b	55.7 a	32.5 b
Safari 20SG (dinotefuran)	12 grams per inch dbh for trees	6.9 bc	53.5 a	31.8 b
Nontreated		8.7 c	52.3 a	28.3 a

^z Mean number of grubs counted in approximately 3.5 cu ft of soil per tree.

^y Means within column followed by the same letter are not significantly different (P>0.05, Fishers LSD).

Gilrein 2005

In 2005, Gilrein examined drenches of five products to manage oriental beetle larvae in lawn-type grass grown in pots. After grass was well established in 1 gal pots, third instar larvae were collected and 10 placed in each pot on Oct 21 and Oct 24. Treatments were applied on Nov 4 and pots were destructively harvested on Dec 27. ANOVA and multiple comparisons among treatment means were performed using a statistical multiple comparison procedure (SuperAnova v. 1.1, Abacus Concepts).

Only Celero and Marathon provided good control of third instar grubs (Table 81). Acelepryn, Cal-Agri, *Metarhizium anisopliae* and Sevin did not provide acceptable control in this test. First or second instar larvae may be more susceptible to these products.

Table 81. Efficacy of several insecticides for Oriental Beetle (*Anomala orientalis*) on lawn type grass, Gilrein, 2005*.

Treatment	Rate	Live Grubs per pot (% control) 53 DAT ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	11.5 fl oz/A	8.2 cd (4.7%)
	23 fl oz/A	7.6 bcd (11.6%)
	46 fl oz/A	7.1 bc (17.4%)
Cal-Agri 50 1% (potassium phosphate)	128 fl oz/100 gal	8.6 d (0.0%)
Celero 16WSG (clothianidin)	4 oz/1320 pots	0.6 a (93.0%)
Marathon II 2F (imidacloprid)	20 g/650 pots	1.1 a (87.2%)
	20 g/244 pots	0.6 a (93.0%)
<i>Metarhizium anisopliae</i> (Strain F52)	14.04 cfu/pot	7.1 bc (17.4%)
	28.08 cfu/pot	6.3 b (26.7%)
Sevin XLR 4F (carbaryl)	6 fl oz/1000 sq ft	7.7 bcd (10.4%)
Nontreated		8.6 d (0.0%)

* Not an IR-4 experiment.

^z Means within columns followed by the same letter are not significantly different at p=0.05 (Fisher's LSD).

Gilrein 2007

In 2007, Gilrein compared several insecticide drenches to control oriental beetle larvae in lawn-type grass grown in pots. After grass was well established in 1 gal pots, third instar larvae were collected and 10 placed in each pot on Nov 5. Treatments were applied on Nov 7 and pots were destructively harvested on Dec 7. ANOVA and multiple comparisons among treatment means were performed using a statistical multiple comparison procedure (SuperAnova v. 1.1, Abacus Concepts).

Only Dylox provided good control of third instar grubs (Table 82). Acelepryn, BAS 320i, Flagship, Hexacide, Safari and Tick-EX did not provide acceptable control in this test.

Table 82. Efficacy of several insecticides for Oriental Beetle (*Anomala orientalis*) on lawn type grass, Gilrein, 2007.

Treatment	Rate	Live Grubs per pot (% control) 30 DAT ^z
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 fl oz/100 gal	4.4 bc (37)
	3.2 fl oz/100 gal	5.2 bcd (26)
BAS 320i 240SC	16 fl oz/100 gal	8.3 ef (0)
Dylox 80S (trichlorfon)	3.75 oz/100 gal	0.5a (93)
Flagship 25WG (thiamethoxam)	8 oz/100 gal	3.8 b (46)
	17 oz/100 gal	4.2 bc (40)
Flagship 0.22G (thiamethoxam)	6 g/pot	3.7 b (47)
Hexacide (rosemary oil)	1.5 qt/100 gal	4.8 bc (31)
Safari 20SG (dinotefuran)	24 oz/100 gal	6.2 cde (11)
Tick-EX EC 11% (<i>Metarhizium anisopliae</i> Strain F52)	21 fl oz/100 gal	8.4 f (0)
	29 fl oz/100 gal	7.8 ef (0)
Nontreated		7.0 def (0)

^z Means within columns followed by the same letter are not significantly different at p=0.05 (Fisher's LSD).

Reding 2006

Lilac (*Syringa vulgaris*) plants were used to determine efficacy of soil drenches of various products against oriental beetle (OB) larvae. Bare root lilacs were planted in one gallon pots on Apr 17 for the study. All treatment rates were applied in 200 ml of solution as a surface drench poured over the potting media, on Jun 5. Each lilac plant was then placed in a separate cage. On Jun 12 three pairs (male/female) of OB's were placed on each caged plant.

To determine efficacy, plant roots and potting media were searched for OB larvae on Sep 19. No larvae were found in any of the treatments, including the Nontreated controls, therefore no efficacy data was acquired (Table 83). Foliar feeding damage by adult OB's could not be accessed on lilac because feeding by adults is minimal. Adult OB's were observed within the cages, on the lilac plants and potting media surface for several weeks after infestation, subsequently the lack of larvae in all pots was unexpected.

Table 83. Efficacy of several insecticides for Oriental Beetle on Lilac (*Syringa vulgaris*), Reding, 2006.

Treatment	Rate Per 100 Gal ^z	Mean number of larvae per plant ^y
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	47.9 oz	0.0 a
Celero 16WSG (clothianidin)	4 oz	0.0 a
Safari 20SG (dinotefuran)	24 oz	0.0 a
Nontreated		0.0 a

^z Each treatment was drenched onto soil using 200 ml per plant.

^y Means within columns followed by the same letter are not significantly different ANOVA ($P = 0.05$), means separated by LSD ($\alpha = 0.05$).

Reding 2009

In 2009 Reding evaluated the efficacy of soil drench applications of Acelepryn, Flagship and Safari against oriental beetle larvae. Bare root lilacs (*Syringa x chinensis*) and white oaks (*Quercus alba*) were planted in two gallon pots on Apr 23 for the study. Each plant was infested with 20 eggs on Jul 16. Insecticides were applied in 300 ml of solution as a surface drench poured over the potting media, on Jul 21. To determine efficacy, oaks were evaluated on Oct 29 and lilacs on Nov 19. Plant roots and potting media were searched for OB larvae.

All insecticides prevented infestation by oriental beetle on both species of plants (Table 84).

Table 84. Efficacy of several insecticides for Oriental Beetle on Lilac (*Syringa x chinensis*) and White Oak (*Quercus alba*), Reding, 2009.

Treatment	Rate Per 100 Gal ^z	Mean number of larvae per plant ^y	
		Lilac 14 WAT	Oak 17 WAT
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	32 fl oz	0.0 b	0.0 b
Flagship 25WG (thiamethoxam)	8 oz	0.0 b	0.0 b
Safari 20SG (dinotefuran)	24 oz	0.0 b	0.0 b
Nontreated		2.6 a	3.0 a

^z Each treatment was drenched onto soil using 300 ml per plant.

^y Means within columns followed by the same letter are not significantly different ANOVA ($P = 0.05$), means separated by LSD ($\alpha = 0.05$).


Comparative Efficacy on Strawberry Rootworm (*Paria fragariae*)

The strawberry rootworm, *Paria fragariae* Wilcox, is a beetle that is a pest of strawberries, blueberries, and even greenhouse roses. It is becoming a major insect pest in container azalea production.

Two experiments were conducted for efficacy on strawberry rootworm infesting azalea. In 2005, Hesselein examined six products: Flagship, Mach 2, Marathon, Ornazin, Orthene, and Talstar. At 21 days after inoculation plants were treated with one of eight treatments: water-treated control, Flagship (thiamethoxam) at 0.18 oz/1000 sq ft, Mach 2 (halofenozide) at 2.9 fl oz/1000 sq ft, Marathon 60 WP (imidacloprid) at 20 gm/ 3000 sq ft, Ornazin (azadirachtin) at 10 oz/ 100 gal, Orthene 97 (acephate) at 12 oz/ 100 gal, and Talstar N (bifenthrin) at 25 fl oz/ 100 gal. Each treatment was replicated 5 times. Drench volume was calculated based on 20% of the container volume. Of these treatments, the best control was achieved with Orthene and Talstar (Graph 85).

In 2007 the strawberry rootworm populations were not sufficiently high to provide a good test of the six treatments planned for that year (Table 86).

Graph 85. Efficacy of several insecticides for Strawberry Rootworm larvae (*Paria fragariae*) on Azalea (*Rhododendron sp.*), Hesselein, 2005.



Treatment columns topped by the same letter are not different, $\alpha=0.05$.

Table 86. Efficacy of several insecticides for Strawberry Rootworm (*Paria fragariae*) on Azalea (*Rhododendron* sp.), Hesselein, 2007.

Treatment	Rate per 100 gal	Adults collected on sticky cards	
		9/25 & 26	11/8
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	0.8 fl oz	0.29 a	0
BAS320i (metaflumizone)	12 fl oz	0.57 a	0
Celero (clothianidin)	12 oz	0.29 a	0
DuraGuard	50 fl oz	0.29 a	0
Met52 (<i>Metarhizium anisopliae</i> Strain F52)	58 fl oz	0.29 a	0
Safari (dinotefuran)	24 oz	0.29 a	0
Nontreated Control		0.43 a ^z	0

^z Means separation tests performed using Tukey's Studentized Range (HSD) Test, $\alpha=0.05$.

Treatments applied on Aug 23 using 375 ml per pot for all but Met52 which used 750 ml per pot.

Results: Trunk and Stem Borers

Bark beetles species in general are often difficult to control due to their behavior or physiology. These are often the insects that can impact and kill large trees in the landscape as well as providing a challenge for growers. Many of bark beetles and stem borers are introduced species and have become problematic because native trees and bushes have not co-evolved and are thus more vulnerable. Because the larvae feed deep within plant tissues, timing of control measures is critical to achieving a successful reduction in insect populations. The registration of effective insecticides for the nursery industry to manage this group of pests is imperative.

Comparative Efficacy on Asian Ambrosia Beetle (Xylosandrus crassiusculus)

The Asian ambrosia beetle or granulate ambrosia beetle, *Xylosandrus crassiusculus*, was introduced into the U.S. from Asia or Africa. In the United States, it is found in Hawaii, Delaware, Maryland, Ohio, and it is widely established in the Southeast. Unlike most ambrosia beetles, this one is known to attack apparently healthy plants. It is reported to attack over 200 species of plants in 41 families, although it prefers hardwoods. Mated females construct galleries, inoculate the wood with the ambrosia fungus, and lay their eggs in excavated galleries of susceptible trees and shrubs. They have become a very important nursery and landscape pest from Texas to New Jersey. In nurseries, seedling and small diameter trees and shrubs are attacked causing girdling, stunting, and death. Asian ambrosia beetle can also attack and kill larger trees, particularly if trees are stressed. Between 5 and 10 attacks on trees less than 10 -15 cm in diameter usually kill this sized tree. The trend in the nursery industry is to grow larger trees in larger pots for instant landscapes. As a result there are higher numbers of stressed trees available.

Ludwig 2004

In 2004, Ludwig conducted an experiment at the Tram-Tex Nursery in Tyler, TX examining the efficacy of Onyx and Talstar Flowable preventative trunk spray applications against ambrosia beetles on container produced redbuds and Bradford pears. Onyx and Talstar were applied to the tree trunks until run-off. Repeat applications were made once for Onyx and twice for Talstar. Tree trunks were visually inspected once a month for signs of beetle attack. Trees that were attacked in one sample period were not checked in following sample periods.

In the first block of redbud trees, both Onyx and Talstar reduced the overall number of ambrosia beetle attacks throughout the experiment from 72% in the Nontreated control to as low as 41% (Table 87). However, various levels of control were achieved at each reading date. This was probably a result of a beetle flight and the pesticide residues declining over time. Both Onyx and Talstar were applied after the assessment in June. The count on July 16 revealed only beetle attacks in the control plants. The results from this block suggest a shorter spray period should be evaluated.

In the second block of redbud trees, the results were mixed (Table 88). The Nontreated trees had an attack rate of 40%, but Onyx at 6.4 oz had more attacks. The other two rates of Onyx did reduce attacks with 12.8 oz having no evidence of ambrosia beetle damage. Talstar reduced the attacks down to 5% of the treated trees.

In the Bradford pear block, the Nontreated trees only had one tree attacked so no conclusions on efficacy can be drawn (Table 89).

Table 87. Efficacy of Onyx and Talstar for *Xylosandrus crassiusculus* on Redbud (*Cercis canadensis*), Ludwig, 2004a.

Treatment ^z	Rate per 100 gal	Number New Attacks			Total Trees	Percent with Attacks
		5/10	6/10	7/16		
Onyx (bifenthrin)	6.4 oz	1	11	0	24	50%
Onyx (bifenthrin)	12.8	0	7	0	17	41%
Onyx (bifenthrin)	16 oz	2	8	0	23	43%
Talstar (bifenthrin)	40 oz	3	9	0	22	55%
Nontreated		4	13	4	29	72%

^zThe Onyx treatments were applied on Apr 9 and Jun 15. The Talstar treatment was applied on Apr 9, May 10, and Jun 15.

Table 88. Efficacy of Onyx and Talstar for *Xylosandrus crassiusculus* on Redbud (*Cercis canadensis*), Ludwig, 2004b.

Treatment ^z	Rate per 100 gal	Number New Attacks		Total Trees	Percent with Attacks
		5/10	6/10		
Onyx (bifenthrin)	6.4 oz	1	6	12	58%
Onyx (bifenthrin)	12.8	0	0	20	0%
Onyx (bifenthrin)	16 oz	2	1	11	27%
Talstar (bifenthrin)	40 oz	0	1	22	5%
Nontreated		2	4	15	40%

^zThe Onyx treatments were applied on Apr 9 and Jun 15. The Talstar treatment was applied on Apr 9, May 10, and Jun 15.

Table 89. Efficacy of Onyx and Talstar for *Xylosandrus crassiusculus* on Bradford Pear (*Pyrus sp*), Ludwig, 2004.

Treatment ^z	Rate per 100 gal	Number New Attacks			Total Trees	Percent with Attacks
		5/10	6/10	7/16		
Onyx (bifenthrin)	6.4 oz	0	0	0	26	0%
Onyx(bifenthrin)	12.8	5	3	1	24	37%
Onyx(bifenthrin)	16 oz	0	0	0	27	0%
Talstar(bifenthrin)	40 oz	1	3	1	20	25%
Nontreated		1	0	0	20	5%

^zThe Onyx treatments were applied on Apr 9 and Jun 15. The Talstar treatment was applied on Apr 9, May 10, and Jun 15.

Mizell 2005

During 2005, Mizell conducted a series of 5 experiments testing various products for their efficacy in controlling Asian ambrosia beetle (AAB). For each experiment, bolts from mimosa (*Albizia julibrissin*) were cut to approximately 46 cm long x 2-7 cm in diameter. Bolts were treated immediately with insecticide by dipping the bolts for 5 sec into the insecticide solution. After the bolts dried a few minutes, they were placed in the field and subjected to a proprietary induction technique and exposed to attack by AAB. Each test consisted of six replicate bolts per insecticide treatment along with six control bolts. Blocks were separated by 20-50 m in a location with known populations and/or active infestations of AAB. AAB attacks observed on each bolt were counted and recorded daily for 7-12 days. The tests were terminated when the number of AAB attacks on the control bolts exceeded an average of 20 per bolt or the time after treatment reached 21 days. Seven to 12 days of field exposure was usually required to attain the number of attacks on the test bolts. The mean number of successful attacks per bolt per treatment was used to evaluate the efficacy of the insecticides. Analysis of variance using the Proc Mixed procedure from SAS 8.1 (Littell et al. 1996) was conducted to analyze the results.

In living nursery trees of less than 10 cm in diameter, the formation of 5-10 successful galleries usually kills the attacked trees and statistics often do not tell the true story. However, because cut bolts were used, these experiments present a distinctly more challenging situation than a typically growing environment. Any product

demonstrating significantly lower attacks than the Nontreated bolts may provide higher levels of control in a landscape or nursery setting.

In this series of experiments, Onyx routinely reduced AAB attacks. Discus performed well in the two experiments where it was tested. Ammo, Asana, Celero, chlorpyrifos and Talstar were variable in performance. In single experiments, Acelepryn and Thiodan were not statistically different than the Nontreated controls.

Table 90. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 1, Mizell, 2005.

Treatment	Rate ai per 100 gal	Mean Attacks per Bolt ^z
Ammo EC (cyfluthrin)	66.3 g	6.3 a
Dursban Turf (chlorpyrifos)	92.8 g	8.0
Onyx (bifenthrin)	212.3 g	6.5 a
Talstar F (bifenthrin)	41.0 g	8.7
Thiodan EC (endosulfan)	83.0	10.7
Nontreated		12.5

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Table 91. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 2, Mizell, 2005.

Treatment	Rate ai per 100 gal	Mean Attacks per Bolt ^z
Asana XL (esfenvalerate)	8.0 g	27.2
Bifenthrin 8% ME (bifenthrin)	42.5 g	10.2
Bifenthrin 8% ME (bifenthrin)	85.0 g	13.5
Onyx (bifenthrin)	212.3 g	0.2 a
Nontreated		48.2

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Table 92. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 3, Mizell, 2005.

Treatment	Rate ai per 100 gal	Mean Attacks per Bolt ^z
Asana XL (esfenvalerate)	76.5 g	5.2 a
Asana XL (esfenvalerate)	153.5 g	3.8 a
Bifenthrin 8% ME (bifenthrin)	170.0 g	0.7 a
Lorsban 4E (chlorpyrifos)	767.5 g	2.7 a
Onyx (bifenthrin)	212.3	0.0 a
Talstar F (bifenthrin)	135.3	0.5 a
Nontreated		15.0

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Table 93. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 4, Mizell, 2005.

Treatment	Rate ai per 100 gal	Mean Attacks per Bolt ^z
Ammo 2.5EC (cyfluthrin)	144.9 g	10.5
Asana XL (esfenvalerate)	95.9 g	11.3
Celero 16WSG (clothianidin)	170.0 g	8.2 a

Discus (cyfluthrin + imidacloprid)	19.8 + 83.3 g	6.3 a
Lorsban 4E (chlorpyrifos)	474.0 g	6.7 a
Onyx (bifenthrin)	212.3 g	4.2 a
Talstar F (bifenthrin)	214.0 g	5.0 a
Nontreated		13.8

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Table 94. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 5, Mizell, 2005.

Treatment	Rate ai per 100 gal	Monticello Mean Attacks per Bolt	Quincy Mean Attacks per Bolt ^z
Acelepryn/DPX-E2Y45 (chlorantraniliprole)	47.0 g	5.3	6.2
Acelepryn/DPX-E2Y45 (chlorantraniliprole)	189.4 g	6.3	6.3
Acelepryn/DPX-E2Y45 (chlorantraniliprole)	754.4 g	3.5	6.3
Celero 16WSG (clothianidin)	38.8 g	2.3	2.8
Celero 16WSG (clothianidin)	77.6 g	2.2	4.7
Discus (cyfluthrin + imidacloprid)	19.8 + 83.3 g	2.0	1.8 a
Discus (cyfluthrin + imidacloprid)	39.6 + 169.6 g	1.5	3.8
Onyx (bifenthrin)	212.3 g	2.3	2.5
Nontreated		3.0	6.3

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Mizell 2006

In 2006, Mizell further explored insecticides along with the common repellent, DEET to prevent Asian ambrosia beetle infestations on mimosa bolts. Methods of treatment and infestation were as in 2005.

In experiment 1, DEET, Discus, Discus + DEET, Onyx, Onyx + Celero, Onyx + DEET, and Onyx + Dursban significantly reduced AAB attacks (Table 51). Although the number of attacks was lower for Azatin, Celero, Dusban and Onyx + Azatin than the Nontreated controls, they were not statistically different 12 DAT.

In experiment 2, while there was a trend for the combination treatments with DEET to have reduced attacks, there were no statistical differences for either location.

Table 95. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 1, Mizell, 2006.

Treatment	Rate per 100 gal	Mean Attacks per Bolt ^z
Azatin XL	256 fl oz	21.3
Celero 16WSG (clothianidin)	8 oz	14.7
DEET	40%	0.0 a
DEET	90%	0.0 a
Discus (cyfluthrin + imidacloprid)	100 fl oz	8.3 a
Discus + DEET	100 fl oz + 40%	0.0 a
Dursban 2E (chlorpyrifos)	32 fl oz	10.3
Onyx (bifenthrin)	32 fl oz	2.7 a
Onyz + Azatin XL	32 fl oz + 2%	10.3
Onyx + Celero 16WSG	32 fl oz + 8 oz	0.0 a
Onyx + DEET	32 fl oz + 40%	3.5 a
Onyx + Dursban 2E	32 fl oz + 32 oz	0.0 a
Nontreated		37.2

^z Number of attacks assessed 12 DAT. Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Table 96. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 2, Mizell, 2006.

Treatment	Rate per 100 gal	Location 1 Mean Attacks per Bolt ^z	Location 2 Mean Attacks per Bolt
DEET	40%	5.5	13.5
Discus(cyfluthrin + imidacloprid)	100 fl oz	20.0	15.5
Discus + DEET	100 fl oz + 40%	2.3	8.3
Onyx (bifenthrin)	32 fl oz	19.7	11.2
Onyx + DEET	32 fl oz + 40%	3.5	9.0
Proclaim (emamectin benzoate)	256 fl oz	9.8	12.3
Talstar L&T (bifenthrin)	32 fl oz	20.8	27.5
Talstar L&T + DEET	32 fl oz + 40%	3.3	16.2
Tempo 2 (cyfluthrin)	32 fl oz	7.3	11.0
Tempo 2 + DEET	32 fl oz + 40%	1.2	11.7
Nontreated		17.5	15.7

^z Number of attacks assessed 12 DAT. Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Mizell 2007

In 2007, Mizell conducted three experiments examining Acelepryn, BAS 320i, *Metarhizium*, and Onyx. Only in experiment 1 were there any statistical differences between treated and Nontreated bolts (Tables 53 – 55). In this experiment, Onyx alone and in combination with two different surfactants provided acceptable reduction in AAB attacks. In experiment 2, *Metarhizium* at several concentrations and in combination with oil did not reduce attacks. In experiment 3, Acelepryn, BAS 320i, and Onyx did not reduce attacks in comparison to the Nontreated controls.

Table 97. Efficacy of Onyx with and without surfactants for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 1, Mizell, 2007.

Treatment	Rate (per 100 gal)	Mean Attacks per Bolt (14 DAT)
Onyx (bifenthrin)	32 fl oz	3.3 ± 1.7 a
Onyx + PentaBark	32 fl oz +1%	3.0 ± 1.1 a
Onyx + Previa	32 fl oz + 1%	2.0 ± 0.3 a
Nontreated		13.8 ± 1.5 b

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Table 98. Efficacy of *Metarhizium* for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 2, Mizell, 2007.

Treatment	Rate (per 100 gal)	Mean Attacks per Bolt (14 DAT)
<i>Metarhizium</i>	1.3 x 10 ⁹	19.8 ± 4.4 a
<i>Metarhizium</i>	3.9 x 10 ⁸	18.0 ± 3.6 a
<i>Metarhizium</i>	3.9 x 10 ⁹	14.5 ± 3.9 a
<i>Metarhizium</i> (oil)	3.9 x 10 ⁹	16.9 ± 4.4 a
Nontreated		18.3 ± 2.4 a

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Table 99. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Mimosa (*Albizia julibrissin*) bolts – Experiment 3, Mizell, 2007.

Treatment	Rate (per 100 gal)	Mean Attacks per Bolt (14 DAT)
Acelepryn (chlorantraniliprole) + PentaBark	32 fl oz +1%	11.0 ± 3.3 a
Acelepryn/DPX-E2Y45(chlorantraniliprole)	32 fl oz/100 gal	18.2 ± 1.5 a
BAS320i (metaflumizone)	16 fl oz/100 gal	11.3 ± 1.4 a
Onyx (bifenthrin)	32 fl oz/100 gal	11.8 ± 4.0 a
Nontreated		16.5 ± 3.6 a

^z Means followed by the same letter are significantly different from the control as determined by a least squares mean test at P<0.05.

Schultz 2009

In 2009, Schultz tested 6 products as either trunk sprays or soil drenches to determine whether any could prevent AAB infestation on magnolia. Applications were made on Apr 7 for Flagship 25WP and Safari 20SG as drenches. The remaining products were applied as trunk sprays on Apr 13: Each magnolia with the exception of the planned Nontreated without ethanol was injected with an ethanol solution on Apr 14 to more uniformly attract adult beetle attacks. Every 2 days over a 4 day period in late April, the number of holes created by adult beetles were counted.

On the first day of assessment, there were very few attacks and no significant differences between treatments with ethanol were observed; on the second assessment two and 4 days later Acelepryn showed higher number of attack holes than other treatments (Table 78).

Table 100. Efficacy Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Magnolia, Schultz, 2009.

Treatment	Rate	Average Number of Attack Holes (cumulative with previous date)		
		4/24	4/26	4/28
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	32 fl oz trunk spray	1.0 a	4.5 a	5.6 a
DPX-HGW86 / Mainspring (cyantraniliprole)	32 fl oz trunk spray	0.0 a	1.4 bc	2.8 abc
Flagship 25WG (thiamethoxam)	8 oz drench	1.75 a	3.4 ab	1.5 bc
Onyx (bifenthrin)	32 fl oz trunk spray	0.0 a	1.0 bc	1.5 bc
Safari 20SG (dinotefuran)	24 oz drench	0.75 a	2.0 abc	2.1 bc
Tolfenpyrad	21 fl oz trunk spray	0.75 a	2.6 abc	4.3 ab
Water		0.0 a	1.1 bc	3.5 ab
Control (no ETOH)		0.0 a	0.1 c	0.0c

Chong 2010

In 2010, Chong compared efficacy of Kontos and OnyxPro applied applied as trunk sprays to prevent AAB infestation on Eastern redbud. Both treatments were applied on Mar 26 (immediately after the first detection of adult flight with ethanol traps) and a second application of OnyxPro was applied on Apr 9 (during the peak flight of adult AAB). Frass tubes or entry holes on each tree were counted before the treatment and weekly for 5 weeks after the treatment. On May 21, about 50 days after the peak flight, glass vials were glued over the entry holes on the trunk with silicon sealant to capture emerging AAB's. The glass vials were collected on Jun 4. The percentage of holes that yielded AAB's or the percentage of emergence was used as an indication of potential systemic effects of Kontos against the AAB.

Kontos was not effective in preventing attacks by AAB during the adult flight period; on the other hand, the standard OnyxPro was effective (Table 101). Treated trees were attacked at the same intensity as the Nontreated trees. The systemic activity of Kontos did not reduce the number of entry holes that yielded offspring of the original attackers. Overall, offspring emerged from 91.6, 96.2 and 98.2% of the entry holes from trees treated with OnyxPro, Kontos and Nontreated, respectively.

Table 101. Efficacy Efficacy of insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Eastern Redbud (*Cercis canadensis*) 'Forest Pansy', Chong, 2010.

Percentage of Trees Attacked ^x							
Treatment	Rate per 100 gal	Pre- Trt	1 WAT	2 WAT	3 WAT	4 WAT	5 WAT
Kontos (spirotetramat)	3.4 fl oz	0 a	3.3 ± 2.1 ab	10.0 ± 2.6 ab	10.0 ± 2.6 ab	11.7 ± 3.1 a	11.7 ± 3.1 a
OnyxPro (bifenthrin)	6.4 fl oz	0 a	0 b	3.5 ± 2.2 b	3.5 ± 2.2 b	3.5 ± 2.2 b	3.5 ± 2.2 b
Nontreated		0 a	11.7 ± 4.0 a	13.3 ± 4.0 a	13.3 ± 4.0 a	13.3 ± 4.0 a	13.3 ± 4.0 a
Average Number of Holes Per Attacked Tree ^x							
Treatment	Rate per 100 gal	Pre- Trt	1 WAT	2 WAT	3 WAT	4 WAT	5 WAT
Kontos (spirotetramat)	3.4 fl oz	-	3.5 ± 0.5 a	5.8 ± 1.6 a	5.8 ± 1.6 a	5.8 ± 1.6 a	5.8 ± 1.6 a
OnyxPro (bifenthrin)	6.4 fl oz	-	-	6.0 a	6.0 a	6.0 a	6.0 a
Nontreated		-	5.3 ± 0.8 a	4.1 ± 0.4 a	4.1 ± 0.4 a	4.1 ± 0.4 a	4.1 ± 0.4 a

^x Means followed by the same letter do not significantly differ (Tukey's LSD test, P = 0.10).

Ludwig 2010

In 2010, Ludwig compared efficacy of several insecticides applied as drenches or sprays to prevent AAB infestation on Eastern redbud (Table 102). All treatments were applied once on Apr 22. To induce beetle attack, trees were injected with 75 ml of 25% ethanol 24 hours after the insecticides were applied. The total number of entrance holes in each tree was recorded 4 weeks after treatment. Unfortunately, majority of trees (including Nontreated) were not attacked so no conclusions can be made.

Table 102. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Redbud (*Cercis canadensis*), Ludwig 2010.

Treatment	Rate Per 100 Gal	Application Method	Mean No. of Holes Per Tree
DPX-E2Y45 / Acelepryn (chlorantraniliprole)	32 fl oz	Drench	0.7
DPX-HGW86 / Mainspring (cyantraniliprole)	32 fl oz	Drench	0.1
Flagship (thiamethoxam)	8 oz	Drench	0.1
OnyxPro (bifenthrin)	32 fl oz	Foliar	0.0
Safari 2G (dinotefuran)	2.2 grams per gal potting media	Drench	0.1
Safari 20SG (dinotefuran)	24 oz	Drench	0.3
Safari 20SG (dinotefuran) + Capsil	8 oz/gal + 2.5% v/v	Basal Spray	0.0
Tolfenpyrad	21 fl oz	Foliar	0.0
Tristar (acetamiprid) + Capsil	8 oz + 2.5% v/v	Spray	0.1
Uninjected Control			0.0
Injected Control			0.1

Chong 2018

In 2018, Chong compared efficacy of several insecticides applied as drenches or trunk sprays in two NC locations, a containerized nursery in Pikeville, and an in-ground nursery in Harmony, to prevent AAB infestation on Eastern redbud. Treatments were applied once or twice in each location on a schedule shown in Table 92 . Unfortunately, data collected at both locations were insufficient to generate useful result tables or make conclusion on the efficacy of the tested products in protecting redbud trees in containerized and in-ground nurseries. No phytotoxicity or residue was observed on the treated redbuds at both locations.

Table 103. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on Redbud (*Cercis canadensis*), ‘Forest Pansy’ Chong, 2018.

Treatment	Rate Per 100 Gal	Applic. Method	Applic. Dates	Phytotoxicity
Botaniguard MAXX (pyrethrins + <i>Beauveria bassiana</i> strain GHA	32 fl oz	Trunk spray	May 1 & 2; May 29 & 30	0.0
ESP 715 (ESP 715)	8.55 fl oz	Drench	April 17 & 18	0.0
Marathon II (imidacloprid)	0.2 fl oz	Drench	April 17 & 18	0.0
Pyridium (beta-cyfluthrin)	2.1 fl oz	Trunk spray	May 1 & 2	0.0
Talstar One (bifenthrin)	21.7 fl oz	Trunk spray	May 1 & 2; May 29 & 30	0.0
V-10433 (V-10433)	5.5 fl oz	Trunk spray	May 1 & 2	0.0
V-10433 (V-10433)	11 fl oz	Trunk spray	May 1 & 2	0.0
Nontreated	-	-	-	0.0

Chong 2019

In 2019, Chong compared efficacy of several insecticides applied as drenches or trunk sprays in two nurseries, to prevent Asian ambrosia beetle infestation on flowering dogwood. Treatments were applied once or twice in each location on a schedule shown in Table 104. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on flooded Flowering Dogwood (*Cornus florida*), 'Cherokee Princess', Chong, 2019.. Trees in the in-ground nursery that were not flooded were virtually not attacked, while trees that were flooded in the containerized nursery were heavily attacked; the data are presented in Table . Pyrethroids (Scimitar and Perm-UP) were the only products that had provided any significant reduction in ambrosia beetle attacks. When the ambrosia beetles became active in mid-April, trees treated with Mainspring + Scimitar, Mainspring-Scimitar and Perm-UP suffered significantly lower numbers of attacks when compared to water and other treatments (at least 69% reduction). Tank mix of Mainspring and Scimitar, as well as Perm-UP, continued to provide good reduction at 4 WAT onward. The good efficacy of pyrethroid products was likely the result of excellent performance at 2 WAT when the beetles were in their peak flight, of which the efficacy carried over to subsequent weeks. The experimental design did not allow for the assessment of whether the drench or trunk spray of Mainspring contributed to the observed efficacy of the tank mix (Mainspring + Scimitar) or rotation treatments (Mainspring – Scimitar). Trees treated with Botanigard MAXX had lower (but not statistically significant) number of attacks at 2 WAT, perhaps because of pyrethrins in the product. However, this treatment did not have any residual efficacy. No phytotoxicity or residue was observed on the treated flowering dogwood at both locations.

Table 104. Efficacy of several insecticides for Asian Ambrosia Beetle (*Xylosandrus crassiusculus*) on flooded Flowering Dogwood (*Cornus florida*), ‘Cherokee Princess’, Chong, 2019.

Treatment	Rate Per 100 Gal	Applic. Method	No. Attacks Per Tree ^x							
			Pre-treat	2 WAT	4 WAT	6 WAT	8 WAT	10 WAT	12 WAT	24 WAT
BCS-507	16 oz	Trunk spray; once	0	7.0 ± 1.7 a	8.7 ± 1.1ab	9.0 ± 1.0 a	9.0 ± 1.0 a	9.0 ± 1.0 a	9.0 ± 1.0 a	9.0 ± 1.0 a
Botanigard MAXX	32 fl oz	Trunk spray; 2 times, monthly	0	4.2 ± 1.1 abc	8.2 ± 1.2 ab	8.5 ± 1.2 ab	8.5 ± 1.2 ab	8.5 ± 1.2 ab	8.5 ± 1.2 ab	8.5 ± 1.2 ab
Perm-UP 3.2 EC	8 fl oz	Trunk spray; 3 times, monthly	0	0.3 ± 0.2 c	3.3 ± 1.3 c	3.3 ± 1.3 c	3.3 ± 1.3 c	3.3 ± 1.3 c	3.3 ± 1.3 c	3.3 ± 1.3 c
Rotation: ESP-715 followed by BCS- 507	8.55 fl oz – 16 oz	Trunk spray; once, ESP-715 3 weeks before BCS-507	0	6.3 ± 2.1 a	9.3 ± 2.1 a	9.5 ± 2.0 a	9.5 ± 2.0 a	9.5 ± 2.0 a	9.5 ± 2.0 a	9.5 ± 2.0 a
Rotation: Mainspring - Scimitar	0.2 fl oz/ft height – 5 fl oz	Mainspring drench 3 wks before Scimitar trunk spray; once	0	1.3 ± 0.8 bc	4.0 ± 1.8 bc	4.3 ± 2.0 bc	4.3 ± 2.0 bc	4.3 ± 2.0 bc	4.3 ± 2.0 bc	4.3 ± 2.0 bc
Rotation: Marathon II followed by BCS-507	0.2 fl oz/inch diameter – 16 oz	Marathon II drench 3 weeks before BCS-507 trunk spray; once	0	5.7 ± 1.9 ab	9.2 ± 0.5 a	9.7 ± 0.8 a	9.7 ± 0.8 a	9.7 ± 0.8 a	9.7 ± 0.8 a	9.7 ± 0.8 a
Tank mix: Mainspring + Scimitar	4 fl oz + 5 fl oz	Trunk spray; 3 times, monthly	0	0.7 ± 0.3 c	2.2 ± 0.9 c	2.5 ± 1.0 c	2.5 ± 1.0 c	2.5 ± 1.0 c	2.5 ± 1.0 c	2.5 ± 1.0 c
V-10433	5.5 fl oz	Trunk spray; once	0	6.5 ± 2.0 a	9.0 ± 2.2 a	9.3 ± 2.0 a	9.3 ± 2.0 a	9.3 ± 2.0 a	9.3 ± 2.0 a	9.3 ± 2.0 a
V-10433	11 fl oz	Trunk spray; once	0	6.8 ± 2.8 a	8.5 ± 2.6ab	9.3 ± 2.3 a	9.3 ± 2.3 a	9.3 ± 2.3 a	9.3 ± 2.3 a	9.3 ± 2.3 a
Water	-	Trunk spray; 3 times, monthly	0	6.7 ± 2.4 a	10.7 ± 2.1 a	11.3 ± 1.7 a	11.3 ± 1.7 a	11.3 ± 1.7 a	11.3 ± 1.7 a	11.3 ± 1.7 a

^x Data were log₁₀(x+0.1) transformed before ANOVA under RCBD. Means followed by the same letter do not significantly differ.

Comparative Efficacy on Ambrosia Beetle (*Xylosandrus germanus*)

The ambrosia beetle or black stem borer, *Xylosandrus germanus*, originated from Asia and now has dispersed to many regions of the US including the Northeast, South, Southeast and the Pacific Northwest. Like the Asian ambrosia beetle, the ambrosia beetle is known to attack both weakened and healthy plants. It is reported to attack over 200 species of plants and is increasingly being recognized as a key pest of ornamental nursery stock. Beyond direct damage to trees and shrubs, there is concern that *X. germanus* is responsible for carrying *Fusarium* fungus into trees as they tunnel into the wood (<http://oregonstate.edu/dept/nurspest/Xylosandrusgermanus.htm>).

Reding 2009

In 2009, Reding evaluated five insecticides applied as trunk sprays or soil drenches for control of the ambrosia beetle (*Xylosandrus germanus*) on magnolia, a known favorite host of *X. germanus*. The plants were purchased in 3 gallon pots from a nursery in Lake County, Ohio on March 17. Soil drenches were applied Apr 28 one week before ethanol injection and trunk sprays were applied May 4 one day before injection. Two sets of control plants were used, one injected with ethanol but Nontreated with insecticide and the other un-injected and Nontreated. On May 5 all magnolia trees, except the un-injected control plants, were injected with ethanol to attract ambrosia beetles and then placed along the wooded border of a field. Trees were watered as needed until the study was terminated on May 27. Beetle attacks (entrance holes in stems) were counted 2, 7, 14, and 23 days after treatment and exposure to beetles. All stems with attack holes were cut from the trees, labeled and brought back to the laboratory to determine survivability and ambrosia beetle species causing damage.

Attacks occurred on all treatments except for the un-injected controls (Table 105). Efficacy results showed a difference in treatments for the first few days but no statistical difference between treatments 5 days after exposure to ambrosia beetles. Flagship applied as drench and Tolfeprad trunk spray provided control that was almost comparable to the standard Onyx. Acelepryn, Safari and Scimitar were ineffective.

Table 105. Efficacy of several insecticides for Ambrosia Beetle (*Xylosandrus germanus*) on Magnolia (*Magnolia virginiana*), Reding, 2009.

Treatment	Rate per 100 gal	Application Method	Mean Attacks ^z	
			May 6 – 11	May 12 - 27
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	32 fl oz	Trunk Spray	2.7 a	13.9
Flagship 25WG (thiamethoxam)	8 oz	Drench	0.9 bc	17.9
Onyx (bifenthrin)	32 fl oz	Trunk Spray	0.4 c	13.9
Safari 20SG (dinotefuran)	24 oz	Drench	1.7 abc	10.0
Safari 20SG (dinotefuran)	24 oz	Trunk Spray	3.2 a	12.5
Scimitar GC(lambda-cyhalothrin)	5 fl oz	Trunk Spray	2.4 ab	17.3
Tolfeprad 15EC	21 fl oz	Trunk Spray	0.8 bc	11.0
Nontreated Control		-	2.6 a	13.9
Un-injected Control		-	0.0 c	0.0

^z Means within columns followed by the same letter are not significantly different ANOVA ($P = 0.05$), means separated by LSD ($\alpha = 0.05$).

Comparative Efficacy on Bronze Birch Borer (*Agrilus anxius*)

The bronze birch borer is a native insect which occurs throughout the range of birch in North America. It is a serious pest of ornamental birch plantings. The bronze birch borer attacks paper (canoe), European white (especially cutleaf variety), gray, yellow, and other birches, as well as poplar, cottonwood and willow. The larva bores in the trunk and larger limbs, often girdling them. Trees weakened by drought or injured are most susceptible to attack. Tree injury is caused by larval tunneling in the inner bark or cambium (<http://www.uri.edu/ce/factsheets/sheets/brbiborer.html>).

In 2006, Nielsen located infested trees in early May followed by applying treatments on May 24 when red horse chestnuts were in bloom. Trees were cut May 15, 2007, and all old emergence holes circled with a wax pencil. Tree trunks were placed upright in the shade at Wooster. New emergence holes (EH) were counted and trunk surface area measured on Aug 29 to evaluate treatment effects.

In the first experiment, Arena, DPX-E2Y45, and Safari reduced the number of exit holes, but not statistically (Table 106). In the second experiment, both NEI 25925 and Flagship significantly reduced infestations (Table 107).

Table 106. Efficacy of several insecticides for Bronze Birch Borer (*Agrilus anxius*) on Weeping European white birch (*Betula pendula*) – Experiment 1, Nielsen, 2006.

Treatment ^z	Rate per inch DBH	Number of Exit Holes ^y
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	14.8 ml	1.3
Arena 50WGD (clothianidin)	1.9 g	2.5
Safari 20SG (dinotefuran)	12 g	0
Nontreated		7.3

^z All treatments applied in 1 gal water as drench to root flare and soil.

^y Means within columns followed by the same letter are not significantly different according to Fisher-Hayton ($P < 0.05$).

Table 107. Efficacy of several insecticides for Bronze Birch Borer (*Agrilus anxius*) on Weeping European white birch (*Betula pendula*) – Experiment 2, Nielsen, 2006.

Treatment ^z	Rate	Number of Exit Holes
NEI 25925 9.25% (acetamiprid) + Capsil	4 ml/ inch DBH + 1%	3.1 a
Flagship 25 WG (thiamethoxam)	16 oz./acre (1200 trees/acre - nursery)	2.1 a
Nontreated		28.9 b

^z All treatments applied in 1 liter of water/tree spraying 18" of lower tree trunk.

^y Means within columns followed by the same letter are not significantly different according to Fisher-Hayton ($P < 0.05$).

Comparative Efficacy on Flatheaded Apple Tree Borer (*Chrysobothris femorata*)

Flatheaded apple tree borer (FHATB) is a native pest of apple, beech, cherry, chestnut, cotoneaster, dogwood, elm, flowering crabapple, hawthorn, hickory, linden, maple, mountain ash, oak, peach, pear, pecan, plum, poplar, quince, shadbush, sycamore, willow, and other hardwood species. This borer preferentially attacks diseased or dying trees, inhabiting all parts of the tree from the base of the trunk to the limbs. Its hosts are primarily newly transplanted nursery stock and to trees that have been weakened from causes such as inappropriate pruning, drought, or inadequate soil or nutrient conditions. Trees suffering from sun scald are regarded as being particularly subject to attack. Like many borers, the FHATB will often girdle a small tree, with a single larva sometimes capable of killing the tree (<http://www.nysipm.cornell.edu/factsheets/treefruit/pests/ab/ab.asp>).

In 2005, Potter tested Discus and Flagship as treatments for two cultivars of red maple. The infestation rate in control trees was > 60% in 2005 (Table 108). Trees that are stressed are more attractive to FHATB so the hot dry 2005 summer likely contributed to borer attacks in this trial. Although Discus significantly reduced the overall

proportion of infested trees, the level of control obtained would not be acceptable in nursery production, nor was it comparable to the > 95% control previously obtained with lindane or chlorpyrifos bark sprays.

Insecticide Trial 2. (Data not shown). In 2006, only 3 of the 300 total trees at the two study sites showed symptoms of FHATB attack, so no conclusions can be drawn regarding efficacy of the treatments. Two of those borers were in Nontreated trees; the other was in a tree that received DPX-E2Y45. The 2006 growing season was relatively cool and had record rainfall so absence of tree stress, as occurs in more typical Kentucky summers, may have contributed to the low incidence of borers.

Table 108. Efficacy of Discus and Flagship Drenches for Flatheaded Apple Tree Borer (*Chrysobothris femorata*) on Red Maples (*Acer rubrum*), Potter, 2005*.

Treatment	Number Infested Trees			Percent with No Attacks		
	October Glory	Red Sunset	Total	October Glory	Red Sunset	Total
Discus (imidacloprid + cyfluthrin)	9	2	11**	44%	88%	66%
Flagship (thiamethoxam)	10	6	16	38%	63%	50%
Nontreated	14	6	20	13%	63%	38%

* Not an IR-4 experiment.

Treatments were applied at high label rate with 1 L per tree. Assumed 0.18 g/1000 sq ft for flagship and 1.5 fl oz per inch DBH for Discus

** Proportion of trees infested is significantly lower for Discus than Nontreated ($\chi^2 = 5.07$, 1 df, $P = 0.02$) but does not differ between Flagship and Nontreated ($\chi^2 = 1.02$, 1 df, $P = 0.31$).

Comparative Efficacy on Banded Ash Clearwing Borer (*Podosesia aureocincta*)

The banded ash clearwing borer is a serious pest of ornamental ash plantings. Day flying females lay eggs in wounds and bark crevices. Hatching larvae chew into the bark and feed both laterally and vertically in the phloem tissue. Later most larvae excavate upward in the sapwood where they concentrate most of their feeding. Completed galleries are about 7 - 32 cm long and 5 - 7 mm in diameter. The sapwood galleries physically weaken the structure of the tree.

In 2008, Nielsen evaluated 5 products for controlling banded ash clearwing borer (Table 109). The green ash trees chosen for this project were heavily infested when treatments were applied on Aug 8, 2008. Presence or absence of new frass was recorded on Jul 13, 2009 as evidence of borer infestation; by this time the borer population had crashed. Frass indexing showed little frass production, even from Nontreated Check trees. No frass production was noted from any of the trees treated with Acelepryn, Aloft, or Tolfeprynad. Onyx reduced frass production. Tristar + Capsil, applied to bark from the soil to a height of 8', was ineffective.

No phytotoxicity was observed.

Table 109. Efficacy of several insecticides for Banded Ash Clearwing Borer (*Podosesia aureocincta*) on Green Ash (*Fraxinus pennsylvanica*); Nielsen, 2008.

Treatment ^z	Rate	Application Method	Frass Index ^y
Acelepryn / DPX-E2Y45 (chlorantraniliprole)	10 fl oz/100 gal	Trunk spray to just above first scaffold limbs	0
Aloft (clothianidin+bifenthrin)	32 fl oz/100 gal	Sprayed trunk and soil at base of tree	0
NEI-25925 (acetamiprid)	6 ml/inch DBH + Capsil	Applied 1 quart of mix/tree up to height of 8'	0.6
Onyx 2EC (bifenthrin)	12.8 fl oz/100 gal	Trunk spray to just above first scaffold limbs	0.2
Tolfenpyrad	24 fl oz/100 gal	Trunk spray to just above first scaffold limbs	0
Nontreated			0.6

^z Treatments were applied Aug 8, 2008 and evaluated Jul 13, 2009. Five plants per treatment were used.

^y Frass index 0-3 where 0 = no frass, 3 = heavy frass

Comparative Efficacy on Peachtree Borer (*Synanthedon exitiosa*)

The peachtree borer is a native insect that is a serious pest of all stone fruits in the genus *Prunus* (peach, cherry, plum, prune, nectarine, apricot) and ornamental shrubs. Larvae feed on the cambium of trunks and large roots forming galleries that are found from about the soil surface to a depth of nearly 30 cm. Extensive larval feeding can girdle and kill trees. Young trees are highly susceptible to severe damage by even a single larva.

In 2009, Nielsen tested six products for their residual efficacy in controlling peachtree borer. An experimental block of purple-leaf sand cherry used in this trial supported a moderately high level infestation of peachtree borer larvae. Treatments were applied on Jun 25 when Catalpa was in full bloom. Presence or absence of new orange frass and gummosis near base of plants were recorded on Sep 13 as evidence of borer infestation. Drench application of Discus, DPX-E2Y45, DPX-HGW86 and Safari, and spray application of Onyx, Scimitar and Tristar provided excellent control (Table 110). The top-dress treatment with Flagship 25WG was somewhat effective, but inadequate for nursery production. Tolfenpyrad spray was ineffective.

No phytotoxicity was observed.

Table 110. Efficacy of several insecticides for Peachtree Borer (*Synanthedon exitiosa*) on Sand Cherry (*Prunus cistina*), Nielsen, 2009.

Treatment ^z	Rate	Application Method	Percent Plants Infested 11 WAT
Acelepryn (chlorantraniliprole)	100 fl oz per 100 gal	Drench	0
Discus (imidacloprid+cyfluthrin)	44 ml/inch DBH	Drench	0
DPX-HGW86 (cyantraniliprole)	0.25 fl oz/quart	Drench	0
Flagship 25WG (thiamethoxam)	0.09 g/linear foot	Top-dress	15
Onyx 2EC (bifenthrin)	102 fl oz/100 gal.	Trunk spray to run-off	0
Safari 20SG (dinotefuran)	6 g/ft shrub height	Drench	0
Scimitar (lambda-cyhalothrin)	5 fl oz/100 gal + Capsil	Trunk spray to run-off	0
Tolfenpyrad	21 fl oz/100 gal	Trunk spray to run-off	25
Tristar 30SG (acetamiprid)	4 oz/inch DBH + Capsil	Trunk spray to run-off	0
Nontreated		-	65

^z Treatments were applied Jun 25 and evaluated Sep 13. Twenty plants per treatment were used.

Results: Efficacy Summary by Product

Acelepryn (chlorantraniliprole)

This product did not impact black vine weevil adults or Asian ambrosia beetle, but it did control adult Japanese beetles in most experiments. Acelepryn also provided excellent control of viburnum leaf beetle. Promising efficacy was observed among researchers for oriental beetle grubs and black vine weevil larvae. Also this product showed good efficacy on May/June beetle grubs, banded ash clearwing borer and peachtree borer. It provided good efficacy on Sri Lankan weevil, but poor efficacy on redheaded flea beetle in single experiments.

Aloft SC (clothianidin+bifenthrin)

In single trials, Aloft provided excellent control of Japanese beetle adults and banded ash clearwing borer. It provided good control of redheaded flea beetles in one hydrangea and two Itea trials, and commercially acceptable control of European elm leaf weevils in a single trial.

Ammo EC (cypermethrin)

This cypermethrin-based product provided reasonable reduction in Asian ambrosia beetle infestations of mimosa bolts.

Ancora/Preferal (*Isaria fumosorosea* Apopka Strain 97 (ATCC 20874))

This biological organism exhibited poor efficacy on Sri Lankan weevil in a single experiment. Poor efficacy on redheaded flea beetles was obtained from 6 of 8 trials.

Arena 50WDG/Celero 16WSG (clothianidin)

No significant reduction in black vine weevil adults or Asian ambrosia beetle infestation was observed with these two products. For Japanese beetle adults, efficacy levels were variable possibly due to differences in techniques and assessments among researchers. Excellent control of viburnum leaf beetle was achieved. Celero also provided good to excellent control of oriental beetle grubs. There were mixed levels of efficacy on black vine weevil larvae.

Asana XL (esfenvalerate)

In two out of four trials, Asana reduced Asian ambrosia beetle infestations. This warrants further exploration.

AzaGuard (azadirachtin)

This product provided poor efficacy on Sri Lankan weevil in a trial and Japanese beetle in two trials.

BAS 320i (metaflumizone)

Metaflumizone exhibited excellent efficacy for black vine weevil adults, Japanese beetle adults, red-headed flea beetle and viburnum leaf beetle. There was virtually no impact on Asian ambrosia beetle infesting mimosa bolts. Excellent efficacy was achieved with black vine weevil larvae. Promising efficacy on oriental beetle larvae was observed in one test.

BCS 507 (BCS 507) good to excellent efficacy for Japanese beetle adults in a single experiment, excellent efficacy on redheaded flea beetle, but poor control of Asian ambrosia beetle.

Bifenthrin 8%ME/Onyx/Talstar (bifenthrin)

These bifenthrin products provided excellent management of Asian ambrosia beetle. Onyx also gave excellent efficacy for Japanese beetle adults. Talstar NL exhibited excellent control of black vine weevil adults. In single trials, Talstar exhibited good control of oriental beetle larvae, and Onyx 2EC provided good to excellent control of ambrosia beetle, banded ash clearwing borer and peachtree borer. Bifenthrin provided good control of redheaded flea beetles in a single trial but Onyx gave poor control in two other trials.

BeetleGONE (*Bacillus thuringiensis galleriae* strain SDS-502)

This product provided good efficacy on Sri Lankan weevil in a single trial, and poor and good efficacy on redheaded flea beetles in 2 experiments.

BotaniGard ES (*Beauveria bassiana*)

This compound exhibited poor efficacy on Sri Lankan weevil in a single experiment. Three trials on Japanese beetle adults provided poor to mediocre efficacy. Generally poor efficacy on redheaded flea beetles was obtained from 3 trials.

BotaniGard MAXX/Xpectro (Pyrethrins + *Beauveria bassiana*)

This product exhibited poor efficacy on Sri Lankan weevil in one experiment. Five trials on Japanese beetle adults provided poor to excellent efficacy. Data collected were insufficient to make conclusions on efficacy of BotaniGard MAXX and other treatments on Ambrosia beetles in 2 experiments; in another trial, poor control was obtained.

CoreTec/Marathon/Merit/Xytect (imidacloprid)

Imidacloprid showed good efficacy for larvae of oriental beetle and May/June beetles, but no efficacy for larvae of black vine weevil and strawberry rootworm in single trials. This active ingredient provided excellent control of Japanese beetles and redheaded flea beetles, and commercially acceptable control of European elm leaf weevils. Data collected were insufficient to make conclusions on efficacy of Marathon and other treatments on Ambrosia beetles in 2 experiments; in another trial, Marathon in rotation with BCS 507 provided poor control.

DEET

This repellent exhibited inconsistent control of Asian ambrosia beetle infestations. There, however, was some indication of improved efficacy for conventional materials when tank mixed with DEET.

Discus (imidacloprid + cyfluthrin)

In two out of four trials, Discus provided a good reduction in Asian ambrosia beetle infestations. It also reduced oriental beetle grubs and peachtree borer, and provided good to excellent control of redheaded flea beetles.

DuraGard/Dursban/Lorsban 4E (chlorpyrifos)

Very little impact was observed for Asian ambrosia beetle with the Dursban formulation, but the Lorsban 4E formulation provided good efficacy.

Dylox 80S (trichlorfon)

Dylox was the only product that provided good control of 3rd instar oriental beetle larvae in one trial.

Flagship (thiamethoxam)

Flagship provided some control of Japanese beetle adults, variable control of black vine weevil, but good control of oriental beetle larvae and redheaded flea beetles. In single trials, it showed poor efficacy on ambrosia beetle and peachtree borer.

Grandevo (*Chromobacterium subtsugae* strain PRAA4-1T)

This biological organism provided variable control of redheaded flea beetles, and commercially acceptable control of European elm leaf weevils. It provided good efficacy on Sri Lankan weevil in a single trial.

Hachi-Hachi/Tolfenpyrad EC (tolfenpyrad)

In single experiments, this compound provided no control of black vine weevil adults, poor control of ambrosia beetle and peachtree borer, commercially acceptable control of European elm leaf weevils, and excellent control of banded ash clearwing borer. It exhibited poor to excellent control of redheaded flea beetles, variable control of Japanese beetle adults, and excellent control of viburnum leaf beetle.

KOC22018-8 (botanical oil blend)

This new product exhibited poor and good control of Japanese beetle adults in one , good efficacy on redheaded flea beetle in two experiments.

Lynx (pyrethrins)

This compound exhibited poor efficacy on Sri Lankan weevil in a single experiment.

Mainspring (cyantraniliprole)

This new product showed good efficacy on peachtree borer in one test. It exhibited variable efficacy on redheaded flea beetles in four trials. In an Asian Ambrosia beetle experiment, tank-mixing or rotation with Scimitar provided good control.

Metarhizium anisopliae

This biological organism was tested as two formulations (Met 52, Tick Ex EC). No efficacy was observed for viburnum leaf beetles, Asian ambrosia beetle, Japanese beetle adults or oriental beetle larvae. While good control of black vine weevil larvae in the greenhouse, little was seen with trials in field containers.

Perm-UP (permethrin) This compound exhibited excellent efficacy on redheaded flea beetle in a single experiment.

Pyridium (Beta-cyfluthrin)

This compound exhibited great efficacy on Japanese beetle adults, and excellent control of redheaded flea beetles in single experiments. Data collected were insufficient to make conclusions on efficacy of Pyridium and other treatments on Ambrosia beetles in 2 experiments.

Safari 2G/20SG (dinotefuran)

Variable levels of efficacy were observed among researchers working with Japanese beetle and redheaded flea beetle adults. Safari did provide very effective control of viburnum leaf beetle. It gave excellent control of oriental beetle, but had variable impact on black vine weevil larvae. In single trials, Safari showed no efficacy on ambrosia beetle, but commercially acceptable control of European elm leaf weevils, and good control of peachtree borer.

Sarissa (cyclaniliprole)

This compound exhibited good to excellent efficacy on Japanese beetle adults in 6 trials, but poor efficacy on Sri Lankan weevil in a single experiment. Poor to excellent efficacy on redheaded flea beetles was obtained from 6 trials.

Sevin (carbaryl)

This compound exhibited excellent efficacy on Japanese beetles in a two experiments.

TetraCURB Conc./Org. (rosemary oil) Both formulations of this product provided poor control of Japanese beetle adults in 2 trials. In one redheaded flea beetle trial, poor efficacy was obtained with the Concentrate formulation.

TriStar 8.5SL/30SG/NEI 25925 (acetamiprid)

Overall efficacy of this compound for Japanese beetle adults was good to excellent. In single trials, Tristar exhibited no efficacy for banded ash clearwing borer, good control of Sri Lankan weevil, and excellent control of peachtree borer. Generally good to excellent efficacy was obtained on redheaded flea beetles in 5 experiments.

UP-Star Gold (bifenthrin)

This compound exhibited excellent efficacy on redheaded flea beetles in a single experiment.

V-10433 (V-10433)

V-10433 exhibited poor to mediocre efficacy on Japanese beetle adults in 6 experiments, and variable control of redheaded flea beetles in 4 experiments. Poor control was obtained.

Venerate (Burkholderia rinojensis A396)

This compound exhibited poor efficacy on Sri Lankan weevil in a single experiment.

VST-06330

This compound exhibited poor efficacy on Japanese beetles in one experiment, and on redheaded flea beetles in 2 trials.

VST-06350

This compound exhibited poor efficacy on Japanese beetles in 2 trials and redheaded flea beetles in a single experiment.

XXpire (spinoteram+sulfoxaflor)

This new product exhibited variable control of redheaded flea beetles.

Phytotoxicity

Injury was only observed with RTSA-721 drench treatments on Crapemyrtle.

Table 111. Summary of Efficacy By Product – Borers and Foliar Feeding Beetles

Note: Table entries are sorted by crop Latin name. Only those trials received by 6/20/2022 are included in the table below.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
25516	Acelepryn (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata) 'Bailey'	Field Container	Alm	RI	2006	Foliar	Excellent control up to 31 DAT using 10 fl oz per 100 gal.
25516	Acelepryn (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Excellent control with 8 fl oz per 100 gal applied twice biweekly.
34637	Acelepryn (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 2: Some reduction in adult viability with 6 fl oz per 100 gal applied biweekly, but feeding damage was greatly reduced.
33791	Acelepryn (Chlorantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	No statistically relevant reduction in a low population with 16 fl oz per 100 gal, and no difference in feeding damage from nontreated controls; no injury or impact on growth observed.
33791	Acelepryn (Chlorantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2019	Foliar	Poor control with 16 fl oz per 100 gal applied twice biweekly.
33791	Acelepryn (Chlorantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Poor efficacy with 6 and 8 fl oz per 100 gal applied 4 times at 10-17 days interval.
26100	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Bronze Birch Borer (Agrilus anxius)	European White Birch (Betula pendula) 'Youngi'	Field Container	Nielsen	OH	2006	Drench	Very low infestation.
25909	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Flatheaded Apple Tree Borer (Chrysobothris femorata)	Maple, Red (Acer rubrum) 'October Glory'	Field In-Ground	Potter	KY	2006	Trunk spray	Very little insect pressure so no conclusions can be drawn.
25512	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Flea Beetle, Garden, Epitrix sp. (Epitrix sp.)	Evening Primrose; Sundrops (Oenothera sp.) O. speciosa var. berlandieri	Field Container	Schultz	VA	2006	Drench	Inconclusive; population too low.
27619	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Black Vine Weevil - adults (Otiorhynchus sulcatus - adults)	Hybrid Yew (Taxus X media)	Field Container	Nielsen	OH	2007	Foliar	No efficacy at 10 oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
27838	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Banded Ash Clearwing Borer (Podosesia aureocincta)	Ash (Fraxinus sp.) F. pennsylvanica	Field In-Ground	Nielsen	OH	2008	Trunk spray	Low infestation. Excellent efficacy at 1qt per 100 gal.
26757	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Drench	No significant reduction of feeding damage; no phytotoxicity and growth effect.
26757	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Foliar	No difference among treatment or untreated plants in this study; no phytotoxicity and no impacts on growth.
26757	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus)	Field In-Ground	Reding	OH	2007	Foliar	No significant reduction of feeding damage at 10 oz per 100 gal; no phytotoxicity and growth effect.
25521	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Cherry, Sargent (Prunus sargentii)	Field Container	Alm	RI	2006	Foliar	Excellent control up to 31 DAT.
25521	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Cherry, Sargent (Prunus sargentii)	Field Container	Alm	RI	2007	Foliar	Excellent control of Japanese beetle adults at 0.026, 0.052 and 0.104 lb ai per 100 gal; equal to bifenthrin.
26948	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	Excellent control at 10 fl oz per 100 gal.
26948	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) R. x odorata 'Tiffany'	Field Container	Davis	MI	2009	Foliar	Significantly reduced number of adults and feeding damage at 10 fl oz per 100 gal; inferior to Flagship.
26948	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Julia Child Butter Gold'	Field Container	Schultz	VA	2007	Foliar	Excellent efficacy.
27788	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Blackberry (Rubus sp.) R. idaeus	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
26488	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. gracilistyla 'Melanostachys'	Field Container	Braman	GA	2006	Foliar	Good efficacy.
26488	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. hakuro nishiki	Field Container	Braman	GA	2007	Foliar	Excellent efficacy at 10 oz per 100 gal.
27783	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Grape (Non-Bearing) (Vitis sp.) 'Seyval Blanc'	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
25734	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.) V. dentatum	Field Container	Costa	VT	2006	Foliar	Effective control; equal to standard permethrin.
25734	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field Container	Costa	VT	2007	Foliar	Some reduction in defoliation severity and extent using 10 fl oz per 100 gal, equivalent level to permethrin.
29862	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field In-Ground	Weston	NY	2007	Foliar	Effective control at 10 oz per 100 gal; equal to imidacloprid.
29292	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Drench	Excellent control at 0.25 fl. Oz./quart water; 1-qt per inch caliper.
28851	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Joe Pye Weed (Eupatorium maculatum)	Field Container	Kunkel	DE	2008	Foliar	No significant control at 10 oz per 100 gal; data not reliable due to high Control mortality and unfavorable environment.
29134	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Drench	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
29872	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Maple, Red (Acer rubrum)	Field Container	Schultz	VA	2008	Trunk spray	Trial 1: Too few attacks for biological significance. Several additional experiments were conducted to refine methodologies for future research.
29872	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Maple, Red (Acer rubrum)	Field Container	Schultz	VA	2008	Trunk spray	Trial 2: Too few attacks for biological significance. Several additional experiments were conducted to refine methodologies for future research.
25481	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Monticello - Results inconclusive due to low numbers of attacks per bolt.
25481	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Quincy - Results inconclusive due to low numbers of attacks per bolt.
25481	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2007	Tree bolt immersion	No control at 32 oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
30497	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010		Untreated trees not attacked; no conclusions can be made.
26826	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Kousa (Cornus kousa)	Field Container	Reding	OH	2007	Trunk spray	No infestation; no phytotoxicity.
29493	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Sweet Bay (Magnolia virginiana)	Field Container	Schultz	VA	2009	Trunk spray	No control at 32 fl oz per 100 gal.
31423	Acelepryn (Dupont) (Chlorantraniliprole)	IRAC 28	Ambrosia Beetle (Xylosandrus germanus)	Sweet Bay (Magnolia virginiana)	Field Container	Reding	OH	2009	Trunk spray	Did not reduce X. germanus beetle attacks at 32 fl oz per 100 gal.
34340	Acephate 97UP (Acephate)	IRAC 1B	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata)	Field Container	Sadof	IN	2018	Drench	No significant control from Acephate at 12 oz per 100 gal and other treatments. Heavy rain during the 9 weeks of this study (~12 inches) may have contributed to product failures.
33980	Acephate Pro 75 WSP (Acephate)	IRAC 1B	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Poor control.
27925	Allectus SC (Bifenthrin + Imidacloprid)	IRAC 3 + IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Maple, Red (Acer rubrum)	Field Container	Schultz	VA	2008	Trunk spray	Trial 1: Too few attacks for biological significance. Several additional experiments were conducted to refine methodologies for future research.
27925	Allectus SC (Bifenthrin + Imidacloprid)	IRAC 3 + IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Maple, Red (Acer rubrum)	Field Container	Schultz	VA	2008	Trunk spray	Trial 2: Too few attacks for biological significance. Several additional experiments were conducted to refine methodologies for future research.
31711	Aloft SC (Clothianadin + bifenthrin)	IRAC 4A + IRAC 3	European Elm Flea Weevil (Orchestes alni)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Foliar	Mediocre but commercially acceptable feeding damage reduction with 15 fl oz per 100 gal applied once.
27841	Aloft SC (Clothianadin + bifenthrin)	IRAC 4A + IRAC 3	Banded Ash Clearwing Borer (Podosesia aureocincta)	Ash (Fraxinus sp.) F. pennsylvanica	Field In-Ground	Nielsen	OH	2008	Sprench	Low infestation. Excellent efficacy at 10 fl oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
28156	Aloft SC (Clothianadin + bifenthrin)	IRAC 4A + IRAC 3	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	Excellent control at 8 fl oz oz per 100 gal.
31716	Aloft SC (Clothianadin + bifenthrin)	IRAC 4A + IRAC 3	Japanese Beetle - adults (Popillia japonica - adults)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Foliar	Did not reduce feeding damage with 15 fl oz per 100 gal applied once.
31588	Aloft SC (Clothianadin + bifenthrin)	IRAC 4A + IRAC 3	Red Headed Flea Beetle (Systema frontalis)	Hydrangea (Hydrangea sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a high leaf damage with 14.9 fl oz per 100 gal applied once; better than Tristar applied once.
31580	Aloft SC (Clothianadin + bifenthrin)	IRAC 4A + IRAC 3	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a low leaf damage with 14.9 fl oz per 100 gal applied once; better than Tristar applied once.
31580	Aloft SC (Clothianadin + bifenthrin)	IRAC 4A + IRAC 3	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Foliar	Excellent reduction of low to moderate leaf damage with 15 fl oz per 100 gal; better than Onyx.
27963	Ammo EC (Cypermethrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 1: About 50% reduction in attacks per bolt treated with 66.3 g ai per 100 gal.
27963	Ammo EC (Cypermethrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 4: Virtually no impact on attacks per bolt treated with 144.9 g ai per 100 gal.
33881	Ancora (Isaria fumosorosea Apopka Strain 97 (ATCC 20874))	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	Statistically significant reduction in a low population 3 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
33881	Ancora (Isaria fumosorosea Apopka Strain 97 (ATCC 20874))	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Bokrathirteen' ('Sweet Summer')	Field Container	Gilrein	NY	2018	Foliar	No significant control with 1 lb per 100 gal applied once.
26098	Arena 50WDG (Clothianadin)	IRAC 4A	Bronze Birch Borer (Agrilus anxius)	European White Birch (Betula pendula) 'Youngi'	Field Container	Nielsen	OH	2006	Drench	Very low infestation.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
27965	Asana XL (Esfenvalerate)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 2: Approximately 50% reduction in attacks per bolt treated with 8 g ai per 100 gal.
27965	Asana XL (Esfenvalerate)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 3: Approximately 60 to 80% reduction in attacks per bolt treated with 76.5 and 153.5 g ai per 100 gal.
27965	Asana XL (Esfenvalerate)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 4: Virtually no impact on attacks per bolt treated with 95.9 g ai per 100 gal.
29522	Astro (Permethrin)	IRAC 3A	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Foliar	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
32656	AzaGuard (Azadirachtin)	IRAC UN	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 16 fl oz per 100 gal applied 3 times weekly; comparable to the standard Tristar.
34152	AzaGuard (Azadirachtin)	IRAC UN	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Poor control with 16 fl oz per 100 gal applied 3 times weekly.
33982	Azatin O (Azadirachtin)	IRAC UN	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Poor control.
26181	Azatin XL (Azadirachtin)	IRAC UN	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt Immersion	Experiment 1: Poor efficacy with a 2% rate.
27617	BAS 320i (Metaflumizone)	IRAC 22B	Black Vine Weevil - adults (Otiorhynchus sulcatus - adults)	Rhododendron (Rhododendron sp.)	Field Container	Nielsen	OH	2007	Foliar	Excellent efficacy at 16 oz per 100 gal.
26825	BAS 320i (Metaflumizone)	IRAC 22B	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus)	Field In-Ground	Reding	OH	2007	Foliar	No significant reduction of feeding damage at 4.5 oz per 100 gal; no phytotoxicity and growth effect.
26945	BAS 320i (Metaflumizone)	IRAC 22B	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Sunsprite'	Field Container	Alm	RI	2008	Foliar	Poor control at 16 fl oz per 100 gal; inferior to standard.
26945	BAS 320i (Metaflumizone)	IRAC 22B	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	Excellent control at 16 oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
26945	BAS 320i (Metaflumizone)	IRAC 22B	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) R. x odorata 'Tiffany	Field Container	Davis	MI	2009	Foliar	Significantly reduced number of adults and feeding damage at 16 fl oz per 100 gal; inferior to Flagship.
26945	BAS 320i (Metaflumizone)	IRAC 22B	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Julia Child Butter Gold'	Field Container	Schultz	VA	2007	Foliar	Excellent efficacy.
26402	BAS 320i (Metaflumizone)	IRAC 22B	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field Container	Costa	VT	2007	Foliar	Some reduction in defoliation severity and extent using 16 oz per 100 gal, equivalent level to permethrin.
29865	BAS 320i (Metaflumizone)	IRAC 22B	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field In-Ground	Weston	NY	2007	Foliar	Effective control at 16 oz per 100 gal; equal to imidacloprid.
28850	BAS 320i (Metaflumizone)	IRAC 22B	Red Headed Flea Beetle (Systema frontalis)	Joe Pye Weed (Eupatorium maculatum)	Field Container	Kunkel	DE	2008	Foliar	No significant control at 16 oz per 100 gal; data not reliable due to high mortality in the untreated and unfavorable environment.
26849	BAS 320i (Metaflumizone)	IRAC 22B	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2007	Tree bolt immersion	No significant control at 16 oz per 100 gal.
34145	BCS-507 (BCS-507)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Excellent control with 16 oz per 100 gal applied twice biweekly.
33786	BCS-507 (BCS-507)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	Statistically relevant reduction in a low population with 16 oz per 100 gal 3 and 7 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
33786	BCS-507 (BCS-507)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2019	Foliar	Excellent control with 16 fl oz per 100 gal applied twice biweekly; comparable to the standard Tristar.
33875	BCS-507 (BCS-507)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Villanassery Joseph	GA	2019	Trunk Spray	Efficacy unknown; No statistical difference among treatments
34235	BCS-507 (BCS-507)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Trunk Spray	Georgetown, SC: No control with 16 oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34243	BCS-507 (BCS-507)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive
32657	beetleGONE! tlc (Bacillus thuringiensis subsp. galleriae Strain SDS-502)	IRAC 11A	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 16 lb per 100 gal + NuFilm applied 3 times weekly; comparable to the standard Tristar.
33976	beetleGONE! tlc (Bacillus thuringiensis subsp. galleriae Strain SDS-502)	IRAC 11A	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Poor control.
33842	beetleGONE! tlc (Bacillus thuringiensis subsp. galleriae Strain SDS-502)	IRAC 11A	Red Headed Flea Beetle (Systema frontalis)	Forsythia (Forsythia sp.)	Field Container	Kunkel	DE	2017	Foliar	Significantly reduced feeding damage with 16 oz applied 3 times weekly; inferior to UpStar Gold.
32638	beetleGONE! tlc (Bacillus thuringiensis subsp. galleriae Strain SDS-502)	IRAC 11A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Baby Lace'	Field Container	Chong	SC	2017	Foliar	Did not significantly reduce adults nor consistently reduce defoliation with 16 lb per 100 gal + LI 700 applied 3 times weekly.
32638	beetleGONE! tlc (Bacillus thuringiensis subsp. galleriae Strain SDS-502)	IRAC 11A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) Vanilla StrawberryTM	Field Container	Frank	NC	2017	Foliar	Poor beetle toxicity and feeding damage reduction with 16 lb per 100 gal + Capsil in lab bioassays; inferior to the standard Marathon.
32638	beetleGONE! tlc (Bacillus thuringiensis subsp. galleriae Strain SDS-502)	IRAC 11A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Pee Gee'	Field Container	Gilrein	NY	2016	Foliar	Poor control with 16 lb per 100 gal + NuFilm P.
33168	beetleGONE! tlc (Bacillus thuringiensis subsp. galleriae Strain SDS-502)	IRAC 11A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Cloyd	KS	2016	Foliar	Mediocre efficacy with 2.2 lb per 100 gal applied Jul 1, 8, 15, 22, Aug 12 and Sep 2.
34339	Benefit 60 WP (imidacloprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata)	Field Container	Sadof	IN	2018	Drench	No significant control from Benefit 60WP at 0.7 oz per 100 gal and other treatments. Heavy rain during the 9 weeks of this study (~12 inches) may have contributed to product failures.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
27966	Bifenthrin 8% ME (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 2: Approximately 75% reduction in attacks per bolt treated with 42.5 and 85 g ai per 100 gal.
27966	Bifenthrin 8% ME (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 3: Almost complete reduction in attacks per bolt treated with 170 g ai per 100 gal.
32658	BotaniGard ES (BioWorks) (Beauveria bassiana Strain GH)	IRAC UNF	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 32 fl oz per 100 gal applied once; inferior to the standard Tristar.
32887	BotaniGard ES (BioWorks) (Beauveria bassiana Strain GH)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Pink Velour'	Field Container	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; no significant differences between treatments.
32641	BotaniGard ES (BioWorks) (Beauveria bassiana Strain GH)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Louis Philippe'	Field Container	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; significantly reduced % leaf damage with 32 fl oz per 100 gal applied once; inferior to Tristar.
32641	BotaniGard ES (BioWorks) (Beauveria bassiana Strain GH)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Louis Phillippe'	Field Container	Addesso	TN	2017	Foliar	Did not significantly reduce feeding damage with 32 fl oz per 100 gal applied twice.
33597	BotaniGard ES (BioWorks) (Beauveria bassiana Strain GH)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.)	Greenhouse	Braman	GA	2017	Foliar	Poor control with 32 fl oz per 100 gal; inferior to the standard Marathon II.
32641	BotaniGard ES (BioWorks) (Beauveria bassiana Strain GH)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Radrazz'	Field Container	Persad	OH	2017	Foliar	74% mortality of adults and significant reduction of leaf damage with 32 fl oz per 100 gal applied 3 times weekly; comparable to Talstar.
33608	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GH)	IRAC 3A + IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi' and 'Pink Velour'	Field Container	Addesso	TN	2018	Foliar	Poor efficacy with 11 and 22 fl oz per 100 gal applied twice; much inferior to the standard Tristar.
34146	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GH)	IRAC 3A + IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Poor to mediocre control with 32 fl oz per 100 gal applied twice biweekly.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34338	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata)	Field Container	Sadof	IN	2018	Drench	No significant control from Botanigard Maxx at 32 fl oz per 100 gal and other treatments. Heavy rain during the 9 weeks of this study (~12 inches) may have contributed to product failures.
34338	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata)	Field Container	Sadof	IN	2018	Foliar	No significant control from Botanigard Maxx at 32 fl oz per 100 gal and other treatments. Heavy rain during the 9 weeks of this study (~12 inches) may have contributed to product failures.
33640	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Harmony: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33640	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Pikeville: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33876	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Villanassery Joseph	GA	2019	Trunk spray	Efficacy unknown; No statistical difference among treatments
34234	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Trunk spray	Georgetown, SC: No significant control with 32 fl oz per 100 gal.
34242	BotaniGard MAXX (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive
34318	BW133 (BW133)	FRAC NC	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Poor control with 5 lb per 100 gal applied twice.
34472	BW133 (BW133)	FRAC NC	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Poor control with 5 lb per 100 gal applied 4 times every 3-7 days.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34374	BW133 (BW133)	FRAC NC	Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Moonrock'	Field Container	Gilrein	NY	2020	Foliar	Good efficacy with 5 lb per 100 gal applied 3 times weekly; inferior to the standard Tristar. Moderate phytotoxicity.
34319	BW238 ES (BW238 ES)		Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Crape Myrtle (<i>Lagerstroemia indica</i>)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Poor control with 2 qt per 100 gal applied twice.
34473	BW238 ES (BW238 ES)		Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Linden, Shamrock (<i>Tilia cordata</i>)	Field In-Ground	Potter	KY	2020	Foliar	Poor control with 2 qt per 100 gal applied twice weekly.
34375	BW238 ES (BW238 ES)		Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Moonrock'	Field Container	Gilrein	NY	2020	Foliar	Good efficacy with 2 qt per 100 gal applied 3 times weekly; inferior to the standard Tristar. Minor phytotoxicity.
34320	BW238 WP (BW238 WP)		Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Crape Myrtle (<i>Lagerstroemia indica</i>)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Poor control with 2 lb per 100 gal applied twice.
34474	BW238 WP (BW238 WP)		Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Linden, Shamrock (<i>Tilia cordata</i>)	Field In-Ground	Potter	KY	2020	Foliar	Poor control with 2 lb per 100 gal applied twice weekly.
25908	Celero 16WSG (Clothianadin)	IRAC 4A	Flatheaded Apple Tree Borer (<i>Chrysobothris femorata</i>)	Maple, Red (<i>Acer rubrum</i>) 'October Glory'	Field In-Ground	Potter	KY	2006	Drench	Very little insect pressure so no conclusions can be drawn.
25511	Celero 16WSG (Clothianadin)	IRAC 4A	Flea Beetle, Garden, <i>Epitrix</i> sp. (<i>Epitrix</i> sp.)	Evening Primrose; Sundrops (<i>Oenothera</i> sp.) <i>O. speciosa</i> var. <i>berlandiere</i>	Field Container	Schultz	VA	2006	Drench	Population declined but numbers too low for statistical analysis
27618	Celero 16WSG (Clothianadin)	IRAC 4A	Black Vine Weevil - adults (<i>Otiorhynchus sulcatus</i> - adults)	Hybrid Yew (<i>Taxus X media</i>)	Field Container	Nielsen	OH	2007	Foliar	Poor efficacy at 4 oz per 100 gal.
26756	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose-Of-Sharon (<i>Hibiscus syriacus</i>) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Drench	No significant reduction of feeding damage; no phytotoxicity and growth effect.
26756	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose-Of-Sharon (<i>Hibiscus syriacus</i>) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Foliar	No difference among treatment or untreated plants in this study; no phytotoxicity and no impacts on growth.
26756	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose-Of-Sharon (<i>Hibiscus syriacus</i>)	Field In-Ground	Reding	OH	2007	Foliar	No significant reduction of feeding damage at 4 oz per 100 gal; no phytotoxicity and growth effect.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
25520	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Cherry, Sargent (Prunus sargentii)	Field Container	Alm	RI	2006	Foliar	Excellent control up to 10 DAT.
26946	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Julia Child Butter Gold'	Field Container	Schultz	VA	2007	Foliar	Excellent efficacy.
27787	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Blackberry (Rubus sp.) R. idaeus	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
26487	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. gracilistyla 'Melanostachys'	Field Container	Braman	GA	2006	Foliar	Poor efficacy.
26487	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. hakuro nishiki	Field Container	Braman	GA	2007	Foliar	Fair efficacy at 4 oz per 100 gal.
25515	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata) 'Bailyei'	Field Container	Alm	RI	2006	Foliar	Good control up to 19 DAT using 6 oz per 100 gal.
27782	Celero 16WSG (Clothianadin)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Grape (Non-Bearing) (Vitis sp.) 'Seyval Blanc'	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
25733	Celero 16WSG (Clothianadin)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.) V. dentatum	Field Container	Costa	VT	2006	Foliar	Effective control; equal to standard permethrin.
25733	Celero 16WSG (Clothianadin)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field Container	Costa	VT	2007	Foliar	Some reduction in defoliation severity and extent using 4 oz per 100 gal, equivalent level to permethrin.
29866	Celero 16WSG (Clothianadin)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field In-Ground	Weston	NY	2007	Foliar	Effective control at 4 oz per 100 gal; better than imidacloprid.
25479	Celero 16WSG (Clothianadin)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 4: Some impact on attacks per bolt treated with 144.9 g ai per 100 gal.
25479	Celero 16WSG (Clothianadin)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Monticello - Results inconclusive due to low numbers of attacks per bolt.
25479	Celero 16WSG (Clothianadin)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Quincy - Results inconclusive due to low numbers of attacks per bolt.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
25479	Celero 16WSG (Clothianadin)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Mediocre efficacy (about 50% reduction in attacks per bolt) with 8 oz per 100 gal.
26828	Celero 16WSG (Clothianadin)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Kousa (Cornus kousa)	Field Container	Reding	OH	2007	Trunk spray	No infestation; no phytotoxicity.
26180	DEET (DEET)	?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree Bolt Immersion	Experiment 1: Excellent control at 40% and 90%.
26180	DEET (DEET)	?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - some reduction in attacks on bolts with 40%, but not statistically significant.
26180	DEET (DEET)	?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no statistically significant differences among treatments.
25477	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Flea Beetle, Garden, Epitrix sp. (Epitrix sp.)	Evening Primrose; Sundrops (Oenothera sp.) O. speciosa var. berlandieri	Field Container	Schultz	VA	2006	Drench	Inconclusive; population too low.
29298	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Drench	Excellent control at 44 ml per inch dbh.
31893	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Foliar	Excellent reduction of low to moderate leaf damage with 50 oz per 100 gal; better than Onyx.
31893	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Frank	NC	2013	Foliar	Results inconclusive due to low infestation; some plant damage reduction with 50 fl oz per 100 gal.
29523	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Drench	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
24764	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 4: Approximately 50% reduction in attacks per bolt treated with 144.9 g ai per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
24764	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Monticello - Results inconclusive due to low numbers of attacks per bolt.
24764	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Quincy - Results inconclusive due to low numbers of attacks per bolt.
24764	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Approximate 70% reduction in attacks per bolt treated with 100 oz per 100 gal.
24764	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - no statistical differences among treatments.
24764	Discus (Imidacloprid + cyfluthrin)	IRAC 4A + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no statistical differences among treatments.
29293	DPX-HGW86 (Cyantraniliprole)	IRAC 28	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Drench	Excellent control at 0.25 fl. oz./quart water; 1-qt per inch caliper.
29135	DPX-HGW86 (Cyantraniliprole)	IRAC 28	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Drench	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
30498	DPX-HGW86 (Cyantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010		Untreated trees not attacked; no conclusions can be made.
29861	DPX-HGW86 (Cyantraniliprole)	IRAC 28	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Sweet Bay (Magnolia virginiana)	Field Container	Schultz	VA	2009	Trunk spray	No significant control at 32 fl oz per 100 gal.
25480	Dursban (Chlorpyrifos)	IRAC 1B	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 1: Some reduction in attacks per bolt treated with 92 g ai per 100 gal.
25480	Dursban (Chlorpyrifos)	IRAC 1B	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Poor efficacy.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31589	Flagship 0.22G (Thiamethoxam)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Hydrangea (Hydrangea sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Foliar	Flagship 25WG used. Significantly reduced a high leaf damage with 8 oz per 100 gal applied once; comparable to Tristar applied once.
26099	Flagship 25WG (Thiamethoxam)	IRAC 4A	Bronze Birch Borer (Agrilus anxius)	European White Birch (Betula pendula) 'Crimson Frost'	Field Container	Nielsen	OH	2006	Trunk/ground spray	Data inconclusive.
25791	Flagship 25WG (Thiamethoxam)	IRAC 4A	Flea Beetle, Garden, Epitrix sp. (Epitrix sp.)	Evening Primrose; Sundrops (Oenothera sp.) O. speciosa var. berlandieri	Field Container	Schultz	VA	2006	Drench	Inconclusive; population too low.
31672	Flagship 25WG (Thiamethoxam)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) R. x odorata 'Tiffany'	Field Container	Davis	MI	2009	Surface of potting media	Significantly reduced number of adults and feeding damage at 2.2 g per gal media.
26491	Flagship 25WG (Thiamethoxam)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. gracilistyla 'Melanostachys'	Field Container	Braman	GA	2006	Foliar	Fair efficacy.
29294	Flagship 25WG (Thiamethoxam)	IRAC 4A	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Top Dress	Inadequate control at 0.09 g per linear ft.
28852	Flagship 25WG (Thiamethoxam)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Joe Pye Weed (Eupatorium maculatum)	Field Container	Kunkel	DE	2008	Foliar	No significant control at 8 oz per 100 gal; data not reliable due to high Control mortality and unfavorable environment.
31581	Flagship 25WG (Thiamethoxam)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012		Good reduction of beetles and reduced damage to leaves but not significantly with 1 and 2 lb per 100 gal applied twice.
31581	Flagship 25WG (Thiamethoxam)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a low leaf damage with 8 oz per 100 gal applied once; better than Tristar applied once.
29136	Flagship 25WG (Thiamethoxam)	IRAC 4A	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Directed	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
29873	Flagship 25WG (Thiamethoxam)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Maple, Red (Acer rubrum)	Field Container	Schultz	VA	2008	Trunk spray	Trial 2: Too few attacks for biological significance. Several additional experiments were conducted to refine methodologies for future research.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
30499	Flagship 25WG (Thiamethoxam)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010	Drench	Untreated trees not attacked; no conclusions can be made.
29495	Flagship 25WG (Thiamethoxam)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Sweet Bay (Magnolia virginiana)	Field Container	Schultz	VA	2009	Drench	No significant control at 8 oz per 100 gal.
31424	Flagship 25WG (Thiamethoxam)	IRAC 4A	Ambrosia Beetle (Xylosandrus germanus)	Sweet Bay (Magnolia virginiana)	Field Container	Reding	OH	2009	Drench	Significantly reduced X. germanus beetle attacks for less than 5 days at 8 oz per 100 gal; comparable to Onyx. Beyond 5 days, all treatments were attacked, possibly due to high rate of ethanol injection.
31708	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	European Elm Flea Weevil (Orchestes alni)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Foliar	Poor but commercially acceptable feeding damage reduction with 1 and 2 lb per 100 gal applied once.
31713	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Japanese Beetle - adults (Popillia japonica - adults)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Foliar	Did not reduce feeding damage with 1 and 2 lb per 100 gal applied 4 times.
31585	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Red Headed Flea Beetle (Systema frontalis)	Hydrangea (Hydrangea sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a high leaf damage with 1 and 2 lb per 100 gal applied twice; comparable to Tristar applied once.
32558	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) 'Autumn Joy'	Field Container	Kunkel	DE	2013	Foliar	Significantly reduced feeding damage with 1 and 2 lb per 100 gal applied twice; inferior to the standard Scimitar.
31577	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a low leaf damage with 1 and 2 lb per 100 gal applied twice; comparable to Tristar applied once.
31577	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Foliar	Poor reduction of low to moderate leaf damage with 2 lb per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31577	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Frank	NC	2013	Foliar	Results inconclusive due to low infestation.
31577	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Kunkel	DE	2013	Foliar	Reduced feeding damage, though not significantly, with 1 and 2 lb per 100 gal applied twice; inferior to the standard Scimitar.
32552	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	FRAC NC	Red Headed Flea Beetle (Systema frontalis)	Woodland Sage (Salvia nemorosa) 'Blue Hill'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 1 and 2 lb per 100 gal applied twice; inferior to the standard Scimitar.
31707	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	European Elm Flea Weevil (Orchestes alni)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Foliar	Poor but commercially acceptable feeding damage reduction with 21 and 32 fl oz per 100 gal applied once.
27621	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Black Vine Weevil - adults (Otiorhynchus sulcatus - adults)	Hybrid Yew (Taxus X media)	Field Container	Nielsen	OH	2007	Foliar	No efficacy at 21 fl oz per 100 gal.
27840	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Banded Ash Clearwing Borer (Podosesia aureocincta)	Ash (Fraxinus sp.) F. pennsylvanica	Field In-Ground	Nielsen	OH	2008	Trunk spray	Low infestation. Excellent efficacy at 24 fl oz per 100 gal.
26951	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	Virtually no control at 21 fl oz per 100 gal.
26951	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) R. x odorata 'Tiffany'	Field Container	Davis	MI	2009	Foliar	Poor efficacy at 21 fl oz per 100 gal + NIS.
26951	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Julia Child Butter Gold'	Field Container	Schultz	VA	2007	Foliar	Good efficacy; some foliar feeding damage.
31712	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Japanese Beetle - adults (Popillia japonica - adults)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Foliar	Did not reduce feeding damage with 21 and 32 fl oz per 100 gal applied once.
26404	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field Container	Costa	VT	2007	Foliar	Some reduction in defoliation severity and extent using 21 fl (?) oz per 100 gal, equivalent level to permethrin.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
29869	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field In-Ground	Weston	NY	2007	Foliar	Effective control at 21 oz per 100 gal; equal to imidacloprid.
29296	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Foliar	Poor control at 21 fl oz per 100 gal.
28853	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Joe Pye Weed (Eupatorium maculatum)	Field Container	Kunkel	DE	2008	Foliar	No significant control at 21 oz per 100 gal; data not reliable due to high Control mortality and unfavorable environment.
31584	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Hydrangea (Hydrangea sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a high leaf damage with 21 and 32 fl oz per 100 gal applied once; better than Tristar applied once.
31576	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a low leaf damage with 21 and 32 fl oz per 100 gal applied once; inferior to Tristar applied once.
31576	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Foliar	Poor reduction of low to moderate leaf damage with 32 fl oz per 100 gal.
31576	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Frank	NC	2013	Foliar	Results inconclusive due to low infestation.
29139	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Foliar	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
30500	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010	Foliar	Untreated trees not attacked; no conclusions can be made.
29496	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Sweet Bay (Magnolia virginiana)	Field Container	Schultz	VA	2009	Trunk spray	No control at 21 fl oz per 100 gal.
31425	Hachi-Hachi EC (Tolfenpyrad)	IRAC 21A	Ambrosia Beetle (Xylosandrus germanus)	Sweet Bay (Magnolia virginiana)	Field Container	Reding	OH	2009	Trunk spray	Significantly reduced X. germanus beetle attacks for < 5 days at 21 fl oz per 100 gal; comparable to Onyx.
32654	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 21 and 27 fl oz per 100 gal applied once; comparable to the standard Tristar.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
33977	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Good control with 21 and 27 fl oz er 100 gal.
33838	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Forsythia (Forsythia sp.)	Field Container	Kunkel	DE	2017	Foliar	Did not reduce feeding damage with 27 fl oz per 100 gal applied once.
32636	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Baby Lace'	Field Container	Chong	SC	2017	Foliar	Did not significantly reduce adults nor consistently reduce defoliation with 27 fl oz per 100 gal applied once.
32636	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) Vanilla StrawberryTM	Field Container	Frank	NC	2017	Foliar	Poor beetle toxicity and feeding damage reduction with 27 fl oz per 100 gal in lab bioassays; inferior to the standard Marathon.
32636	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry	Field Container	Frank	NC	2019	Foliar	Statistically relevant reduction in a low population with 27 fl oz per 100 gal 3 and 7 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
32636	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Pee Gee'	Field Container	Gilrein	NY	2016	Foliar	Excellent control with 21 and 27 fl oz per 100 gal; comparable to the standard Scimitar.
32636	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Bokrathirteen' ('Sweet Summer')	Field Container	Gilrein	NY	2018	Foliar	Great control with 27 fl oz per 100 gal applied once; almost comparable to the standard Tristar.
32636	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata)	Field Container	Sadof	IN	2018	Foliar	No significant control from Hachi Hachi SC at 27 fl oz per 100 gal and other treatments. Heavy rain during the 9 weeks of this study (~12 inches) may have contributed to product failures.
32560	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) 'Autumn Joy'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 21 and 32 fl oz per 100 gal + Capsil applied once; inferior to the standard Scimitar applied twice.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
32548	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Cloyd	KS	2016	Foliar	Mediocre efficacy with 21 fl oz per 100 gal applied Jul 1, 8, 15, 22, Aug 12 and Sep 2.
32548	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 21 and 32 fl oz per 100 gal + Capsil applied once; inferior to the standard Scimitar applied twice.
32554	Hachi-Hachi SC (Tolfenpyrad)	IRAC 21A	Red Headed Flea Beetle (Systema frontalis)	Woodland Sage (Salvia nemorosa) 'Blue Hill'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 21 and 32 fl oz per 100 gal + Capsil applied once; inferior to the standard Scimitar applied twice.
34321	ISM-555 (ISM-555, A21377X)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Good to excellent control with 5.76 fl oz per 100 gal applied once or twice.
34321	ISM-555 (ISM-555, A21377X)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Foliar	Great efficacy with 5.76 fl oz per 100 gal applied 2 or 3 times weekly; almost comparable to the standard Onyx.
34475	ISM-555 (ISM-555, A21377X)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Good to excellent control with 5.76 fl oz per 100 gal + Capsil applied twice weekly or biweekly.
34475	ISM-555 (ISM-555, A21377X)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 2: No reduction in adult viability with 5.76 fl oz per 100 gal applied weekly or biweekly, but feeding damage was reduced.
34377	ISM-555 (ISM-555, A21377X)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2020	Foliar	Great efficacy with 5.76 fl oz per 100 gal + Capsil applied 3 times weekly; comparable to standard Tristar. No phytotoxicity.
34377	ISM-555 (ISM-555, A21377X)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Fire and Ice'	Field Container	Gilrein	NY	2021	Foliar	Excellent efficacy with 3.84 and 5.76 fl oz per 100 gal. No crop injury observed.
34377	ISM-555 (ISM-555, A21377X)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Great efficacy with 3.84 and 5.76 fl oz per 100 gal + Capsil applied 4 times at 10-17 days interval.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34148	KOC22018-8 (Botanical Oil Blend)		Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Linden, Shamrock (<i>Tilia cordata</i>)	Field Container	Potter	KY	2019	Foliar	Poor to mediocre control with 128 fl oz per 100 gal applied 3 times weekly.
33789	KOC22018-8 (Botanical Oil Blend)		Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	No statistically relevant reduction in a low population with 128 fl oz per 100 gal, and no difference in feeding damage from nontreated controls; no injury or impact on growth observed.
33789	KOC22018-8 (Botanical Oil Blend)		Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Moonrock'	Field Container	Gilrein	NY	2019	Foliar	Great control with 128 fl oz per 100 gal applied 3 times weekly; slightly inferior to the standard Tristar.
29904	Kontos (BYI 8330 240SC) (Spirotetramat)	IRAC 23	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	American Redbud, Eastern Redbud (<i>Cercis canadensis</i>) 'Forest Pansy'	Field Container	Chong	SC	2010	Trunk spray	No significant control at 3.4 fl oz per 100 gal.
27967	Lorsban 4E (Chlorpyrifos)	IRAC 1B	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 3: Approximately 80% reduction in attacks per bolt treated with 767.5 g ai per 100 gal.
27967	Lorsban 4E (Chlorpyrifos)	IRAC 1B	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 4: Approximately 50% reduction in attacks per bolt treated with 144.9 g ai per 100 gal.
32659	Lynx EC 5.0 (Pyrethrins)	IRAC 3A	Sri Lankan Weevil (<i>Myllocerus undatus</i>)	Shoeblackplant; Chinese hibiscus (<i>Hibiscus rosa-sinensis</i>) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 16 oz per 100 gal applied twice weekly; comparable to the standard Tristar.
33790	Mainspring GNL 200SC (Cyantraniliprole)	IRAC 28	Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	Statistically relevant reduction in a low population with 27 fl oz per 100 gal 3 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
33790	Mainspring GNL 200SC (Cyantraniliprole)	IRAC 28	Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Moonrock'	Field Container	Gilrein	NY	2019	Foliar	Excellent control with 16 fl oz per 100 gal applied once; comparable to the standard Tristar.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
32555	Mainspring GNL 200SC (Cyantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) 'Autumn Joy'	Field Container	Kunkel	DE	2013	Foliar	Significantly reduced feeding damage with 8 and 16 fl oz per 100 gal applied twice; inferior to the standard Scimitar.
31892	Mainspring GNL 200SC (Cyantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Foliar	Poor reduction of low to moderate leaf damage with 16 oz per 100 gal.
31892	Mainspring GNL 200SC (Cyantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Frank	NC	2013	Foliar	Results inconclusive due to low infestation; some plant damage reduction with 8 and 16 fl oz per 100 gal.
31892	Mainspring GNL 200SC (Cyantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Kunkel	DE	2013	Foliar	Reduced feeding damage, though not significantly, with 16 fl oz per 100 gal applied twice; inferior to the standard Scimitar.
32549	Mainspring GNL 200SC (Cyantraniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Woodland Sage (Salvia nemorosa) 'Blue Hill'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 8 and 16 fl oz per 100 gal applied twice; inferior to the standard Scimitar.
31591	Marathon 1% granular (Imidacloprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Hydrangea (Hydrangea sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Broadcast	Significantly reduced a high leaf damage with 7 g per pot; better than Tristar applied once.
31583	Marathon 1% granular (Imidacloprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012	Broadcast	Significantly reduced a low leaf damage with 7 g per pot; better than Tristar applied once.
33596	Marathon II (Imidacloprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.)	Greenhouse	Braman	GA	2017	Foliar	Excellent control with 50 ml per 100 gal.
33581	Marathon II (Imidacloprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) Vanilla Strawberry™	Field Container	Frank	NC	2017	Foliar	Good beetle toxicity and feeding damage reduction with 1.7 fl oz per 100 gal in lab bioassays; best treatment.
34322	MBI 203 SC2 (MBI 203)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Poor control with 128 fl oz per 100 gal applied twice.
34322	MBI 203 SC2 (MBI 203)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Foliar	Good efficacy with 128 fl oz per 100 gal applied twice weekly; almost comparable to the standard Onyx.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34476	MBI 203 SC2 (MBI 203)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Poor control with 128 fl oz per 100 gal applied twice weekly.
34476	MBI 203 SC2 (MBI 203)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 2: No reduction in adult viability or feeding damage with 128 fl oz per 100 gal applied weekly.
34378	MBI 203 SC2 (MBI 203)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2020	Foliar	Poor efficacy with 128 fl oz per 100 gal applied twice weekly; inferior to the standard Tristar. No phytotoxicity.
34378	MBI 203 SC2 (MBI 203)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Fire and Ice'	Field Container	Gilrein	NY	2021	Foliar	No effective population reduction with 128 fl oz per 100 gal. No injury observed.
34378	MBI 203 SC2 (MBI 203)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Poor efficacy with 5 fl oz per 100 gal + Capsil applied 4 times at 10-17 days interval.
34323	MBI 306 (MBI 306)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Poor control with 5 fl oz per 100 gal applied twice.
34323	MBI 306 (MBI 306)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Foliar	Poor efficacy with 2.5 AND 5 fl oz per 100 gal applied twice weekly.
34477	MBI 306 (MBI 306)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Poor control with 5 fl oz per 100 gal applied twice weekly.
34477	MBI 306 (MBI 306)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 2: Some reduction in adult viability with 5 fl oz applied weekly but no impact on feeding damage; no reduction in adult viability or feeding damage with 2.5 fl oz applied weekly.
34379	MBI 306 (MBI 306)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2020	Foliar	Poor efficacy with 5 fl oz per 100 gal applied twice weekly; inferior to the standard Tristar. No phytotoxicity.
34379	MBI 306 (MBI 306)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Fire and Ice'	Field Container	Gilrein	NY	2021	Foliar	No effective population reduction with 5 fl oz per 100 gal. No injury observed.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34379	MBI 306 (MBI 306)		Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Poor efficacy with 2.5 and 5 fl oz per 100 gal + Capsil applied 4 times at 10-17 days interval.
32660	MBI-203 WDG (<i>Chromobacterium subtsugae</i>)		Sri Lankan Weevil (<i>Myllocerus undatus</i>)	Shoeblackplant; Chinese hibiscus (<i>Hibiscus rosa-sinensis</i>) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 3 lb per 100 gal + NuFilm applied 3 times weekly; better than the standard Tristar.
27620	Met52 (<i>Metarhizium anisopliae</i> strain F52)	FRAC NC & IRAC UNF	Black Vine Weevil - adults (<i>Otiorhynchus sulcatus</i> - adults)	Hybrid Yew (<i>Taxus X media</i>)	Field Container	Nielsen	OH	2007	Foliar	No efficacy at 29 fl oz per 100 gal.
26403	Met52 (<i>Metarhizium anisopliae</i> strain F52)	FRAC NC & IRAC UNF	Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>)	Arrowwood (<i>Viburnum</i> sp.)	Field Container	Costa	VT	2007	Foliar	No efficacy at 29 oz per 100 gal.
29867	Met52 (<i>Metarhizium anisopliae</i> strain F52)	FRAC NC & IRAC UNF	Viburnum Leaf Beetle (<i>Pyrrhalta viburni</i>)	Arrowwood (<i>Viburnum</i> sp.)	Field In-Ground	Weston	NY	2007	Foliar	No efficacy at 29 oz per 100 gal
26850	Met52 (<i>Metarhizium anisopliae</i> strain F52)	FRAC NC & IRAC UNF	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2007	Tree bolt immersion	No significant control at rates up to 3.9 X 10,000,000,000.
26101	NEI 25925 (Acetamiprid)	IRAC 4A	Bronze Birch Borer (<i>Agrilus anxius</i>)	European White Birch (<i>Betula pendula</i>) 'Crimson Frost'	Field Container	Nielsen	OH	2006	Trunk spray	Data inconclusive.
25912	NEI 25925 (Acetamiprid)	IRAC 4A	Flatheaded Apple Tree Borer (<i>Chrysobothris femorata</i>)	Maple, Red (<i>Acer rubrum</i>) 'October Glory'	Field In-Ground	Potter	KY	2006	Trunk spray	Very little insect pressure so no conclusions can be drawn.
25910	Onyx (Bifenthrin)	IRAC 3A	Flatheaded Apple Tree Borer (<i>Chrysobothris femorata</i>)	Maple, Red (<i>Acer rubrum</i>) 'October Glory'	Field In-Ground	Potter	KY	2006	Trunk spray	Very little insect pressure so no conclusions can be drawn.
25831	Onyx (Bifenthrin)	IRAC 3A	Flea Beetle, Garden, <i>Epitrix</i> sp. (<i>Epitrix</i> sp.)	Evening Primrose; Sundrops (<i>Oenothera</i> sp.) <i>O. speciosa</i> var. <i>berlandieri</i>	Field Container	Schultz	VA	2006	Foliar	Inconclusive; population too low.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
29303	Onyx (Bifenthrin)	IRAC 3A	Banded Ash Clearwing Borer (<i>Podosesia aureocincta</i>)	Ash (<i>Fraxinus</i> sp.) <i>F. pennsylvanica</i>	Field In-Ground	Nielsen	OH	2008	Trunk spray	Low infestation. Fair efficacy at 12.8 fl oz per 100 gal.
34509	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Crape Myrtle (<i>Lagerstroemia indica</i>)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Excellent control with 26 fl oz per 10-gal
25524	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Cherry, Sargent (<i>Prunus sargentii</i>)	Field Container	Alm	RI	2006	Foliar	Excellent control up to 31 DAT.
25524	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Cherry, Sargent (<i>Prunus sargentii</i>)	Field Container	Alm	RI	2007	Foliar	Excellent control of Japanese beetle adults at 0.2 lb ai per 100 gal.
26949	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose (<i>Rosa</i> sp.) 'Sunsprite'	Field Container	Alm	RI	2008	Foliar	Standard treatment; effective control at 6.4 fl oz per 100 gal.
26949	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose (<i>Rosa</i> sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	Excellent control at 12.8 fl oz per 100 gal.
26949	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose (<i>Rosa</i> sp.) 'Julia Child Butter Gold'	Field Container	Schultz	VA	2007	Foliar	Excellent efficacy.
27791	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Blackberry (<i>Rubus</i> sp.) <i>R. idaeus</i>	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
25519	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Linden, Shamrock (<i>Tilia cordata</i>) 'Bailyei'	Field Container	Alm	RI	2006	Foliar	Good control up to 31 DAT using 12.8 fl oz per 100 gal.
27786	Onyx (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Grape (Non-Bearing) (<i>Vitis</i> sp.) 'Seyval Blanc'	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
29300	Onyx (Bifenthrin)	IRAC 3A	Peachtree Borer (<i>Synanthedon exitiosa</i>)	Plum (<i>Prunus</i> sp.) <i>P. cistina</i>	Field In-Ground	Nielsen	OH	2009	Foliar	Excellent control at 3.2 pt per 100 gal
31587	Onyx (Bifenthrin)	IRAC 3A	Red Headed Flea Beetle (<i>Systema frontalis</i>)	Hydrangea (<i>Hydrangea</i> sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a high leaf damage with 12.8 fl oz per 100 gal applied once; comparable to Tristar applied once.
31579	Onyx (Bifenthrin)	IRAC 3A	Red Headed Flea Beetle (<i>Systema frontalis</i>)	Virginia Sweetspire (<i>Itea virginica</i>)	Field Container	Braman	GA	2013	Foliar	Poor reduction of low to moderate leaf damage with 12.8 fl oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31579	Onyx (Bifenthrin)	IRAC 3A	Red Headed Flea Beetle (<i>Systema frontalis</i>)	Virginia Sweetspire (<i>Itea virginica</i>)	Field Container	Frank	NC	2013	Foliar	Results inconclusive due to low infestation; some plant damage reduction with 40 fl oz per 100 gal.
29524	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Redbay (<i>Xyleborus glabratus</i>)	Redbay (<i>Persea borbonia</i>)	Field In-Ground	Pena	FL	2009	Foliar	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 1: About 50% reduction in attacks per bolt treated with 212.3 g ai per 100 gal.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 2: Approximately 99% reduction in attacks per bolt treated with 212.3 g ai per 100 gal.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 3: 100% reduction in attacks per bolt treated with 212.3 g ai per 100 gal.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 4: Approximately 70% reduction in attacks per bolt treated with 144.9 g ai per 100 gal.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Monticello - Results inconclusive due to low numbers of attacks per bolt.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 5: Quincy - Results inconclusive due to low numbers of attacks per bolt.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Effective control using 32 fl oz per 100 gal.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - no significant differences among treatments.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	Mimosa Silk Tree (<i>Albizia julibrissin</i>)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no significant differences among treatments.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2007	Tree bolt immersion	Test 1: significant control at 16 oz per 100 gal.
25220	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2007	Tree bolt immersion	Test 2: no significant control at 32 oz per 100 gal.
26137	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	SC	2010	Trunk spray	Effective control at 6.4 fl oz per 100 gal.
26137	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field In-Ground	Ludwig	TX	2004	Trunk spray	Inconclusive results due to application timing.
30504	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010		Untreated trees not attacked; no conclusions can be made.
27721	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Kousa (Cornus kousa)	Field Container	Reding	OH	2007	Trunk spray	No infestation; no phytotoxicity.
29497	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Sweet Bay (Magnolia virginiana)	Field Container	Schultz	VA	2009	Trunk spray	No significant control at 32 fl oz per 100 gal.
31426	Onyx (Bifenthrin)	IRAC 3A	Ambrosia Beetle (Xylosandrus germanus)	Sweet Bay (Magnolia virginiana)	Field Container	Reding	OH	2009	Trunk spray	Significantly reduced X. germanus beetle attacks for < 5 days at 32 fl oz per 100 gal.
26429	Perm-Up 3.2EC (Permethrin)	IRAC 3A	Flea Beetle, Garden, Epitrix sp. (Epitrix sp.)	Evening Primrose; Sundrops (Oenothera sp.) O. speciosa var. berlandieri	Field Container	Schultz	VA	2006	Foliar	Inconclusive; population too low.
34987	Perm-Up 3.2EC (Permethrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Villanassery Joseph	GA	2019	Trunk spray	Efficacy unknown; No statistical difference among treatments
34241	Perm-Up 3.2EC (Permethrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Trunk spray	Georgetown, SC: Excellent control with 8 fl oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34249	Perm-Up 3.2EC (Permethrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive
34620	Pradia (Cyclaniliprole + Flonicamid)	IRAC 28 + IRAC 29	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Mediocre efficacy with 17 fl oz per 100 gal applied 4 times at 10-17 days interval.
26492	Precise Acephate (Acephate)	IRAC 1B	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. gracilistyla 'Melanostachys'	Field Container	Braman	GA	2006	Broadcast	Fair efficacy.
33978	Pre-fe-ral (Biobest) (Paecilomyces fumosoroseus)	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Poor control.
32661	Preferal (SePro) (Isaria fumosoroseus)	FRAC NC & IRAC UNF	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 1 lb per 100 gal applied once; inferior to the standard Tristar.
33843	Preferal (SePro) (Isaria fumosoroseus)	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Forsythia (Forsythia sp.)	Field Container	Kunkel	DE	2017	Foliar	Did not significantly reduce feeding damage with 8 oz per 100 gal applied twice biweekly.
32639	Preferal (SePro) (Isaria fumosoroseus)	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Baby Lace'	Field Container	Chong	SC	2017	Foliar	Did not significantly reduce adults nor consistently reduce defoliation with 1 lb per 100 gal applied once.
32639	Preferal (SePro) (Isaria fumosoroseus)	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) Vanilla StrawberryTM	Field Container	Frank	NC	2017	Foliar	Poor beetle toxicity and feeding damage reduction with 1 lb per 100 gal in lab bioassays; inferior to the standard Marathon.
32639	Preferal (SePro) (Isaria fumosoroseus)	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Pee Gee'	Field Container	Gilrein	NY	2016	Foliar	Poor control with 1 lb per 100 gal.
33169	Preferal (SePro) (Isaria fumosoroseus)	FRAC NC & IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Cloyd	KS	2016	Foliar	Good efficacy with 1 lb per 100 gal applied Jul 1, 8, 15, 22, Aug 12 and Sep 2.
28002	Proclaim (Enamectin benzoate)	IRAC 6	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - no statistical difference among treatments.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
28002	Proclaim (Emamectin benzoate)	IRAC 6	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no statistical difference among treatments.
33610	Pyridium (Beta-cyfluthrin)	IRAC 3A	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi' and 'Pink Velour'	Field Container	Addesso	TN	2018	Foliar	Great efficacy with 2 fl oz per 100 gal applied once; slightly inferior to the standard Tristar.
33880	Pyridium (Beta-cyfluthrin)	IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Bokrathirteen' ('Sweet Summer')	Field Container	Gilrein	NY	2018	Foliar	Excellent control with 2.1 fl oz per 100 gal applied once; comparable to the standard Tristar.
33641	Pyridium (Beta-cyfluthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Harmony: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33641	Pyridium (Beta-cyfluthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Pikeville: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33877	Rotation: ESP 715 / BCS-507 (Fluopyram / BCS-507)	FRAC 7 / ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Villanassery Joseph	GA	2019	Trunk spray	Efficacy unknown; No statistical difference among treatments
34236	Rotation: ESP 715 / BCS-507 (Fluopyram / BCS-507)	FRAC 7 / ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Trunk spray	Georgetown, SC: No significant control with 8.55 fl oz/16 oz per 100 gal.
34244	Rotation: ESP 715 / BCS-507 (Fluopyram / BCS-507)	FRAC 7 / ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive
33643	Rotation: ESP 715 / Pyridium (Fluopyram / beta-cyfluthrin)	FRAC 7 / IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Harmony: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33643	Rotation: ESP 715 / Pyridium (Fluopyram / beta-cyfluthrin)	FRAC 7 / IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Pikeville: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34240	Rotation: Mainspring / Scimitar (cyantraniliprole / lambda cyhalothrin)	IRAC 28 / IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Drench / Trunk spray	Georgetown, SC: Mediocre control with 0.2 fl oz per ft ht/5 fl oz per 100 gal; inferior to the standard Perm-Up.
34248	Rotation: Mainspring / Scimitar (cyantraniliprole / lambda cyhalothrin)	IRAC 28 / IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Drench / Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive
33878	Rotation: Marathon II / BCS-507 (Imidacloprid / BCS-507)	IRAC 4A / ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Villanassery Joseph	GA	2019	Drench / Trunk spray	Efficacy unknown; No statistical difference among treatments
34237	Rotation: Marathon II / BCS-507 (Imidacloprid / BCS-507)	IRAC 4A / ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Drench / Trunk Spray	Georgetown, SC: No significant control with 0.2 fl oz per in diam/16 oz per 100 gal.
34245	Rotation: Marathon II / BCS-507 (Imidacloprid / BCS-507)	IRAC 4A / ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Drench / Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive
33644	Rotation: Marathon II / Pyridium (Imidacloprid / beta-cyfluthrin)	IRAC 4A / IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Drench / Trunk spray	Harmony: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33644	Rotation: Marathon II / Pyridium (Imidacloprid / beta-cyfluthrin)	IRAC 4A / IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Drench / Trunk spray	Pikeville: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
34324	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Basal spray	Good control with 100 ml per gal applied once.
34324	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Drench	Poor control with 10 ml + 120 ml water per inch caliper applied once.
34324	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Good control with 650 ml per 100 gal applied once.
34324	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Basal spray	Poor efficacy with 100 ml per 1 gal applied once.
34324	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Drench	Poor efficacy with 10 ml + 120 ml water / inch caliper @ 1 inch height applied once.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34324	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Foliar	Good efficacy with 650 ml per 100 gal applied once; almost comparable to the standard Onyx.
34478	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Basal spray	Poor control with 100 ml per gal applied once.
34478	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Drench	Poor control with 10 ml + 115 ml water per inch DBH applied once.
34478	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Excellent control with 650 ml per 100 gal applied once.
34478	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 1: Excellent mortality and reduction in feeding damage with 650 ml per 100 gal.
34478	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Soil Injection	Experiment 1: No mortality or reduction in feeding damage with adult confinement on leaves with 10ml per inch caliper soil injection approximately 2 months after application.
34478	RTSA 721 (RTSA 721)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Tree Injection	Experiment 1: No mortality or reduction in feeding damage with adult confinement on leaves with 100ml/gal trunk injection approximately 2 months after application.
26102	Safari 20SG (Dinotefuran)	IRAC 4A	Bronze Birch Borer (Agrilus anxius)	European White Birch (Betula pendula) 'Youngi'	Field Container	Nielsen	OH	2006	Drench	Very low infestation.
25911	Safari 20SG (Dinotefuran)	IRAC 4A	Flatheaded Apple Tree Borer (Chrysobothris femorata)	Maple, Red (Acer rubrum) 'October Glory'	Field In-Ground	Potter	KY	2006	Drench	Very little insect pressure so no conclusions can be drawn.
25513	Safari 20SG (Dinotefuran)	IRAC 4A	Flea Beetle, Garden, Epitrix sp. (Epitrix sp.)	Evening Primrose; Sundrops (Oenothera sp.) O. speciosa var. berlandieri	Field Container	Schultz	VA	2006	Drench	Population declined but numbers too low for statistical analysis.
31709	Safari 20SG (Dinotefuran)	IRAC 4A	European Elm Flea Weevil (Orchestes alni)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012		Mediocre but commercially acceptable feeding damage reduction with 12 g per DBH inch applied once at dormancy.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
27623	Safari 20SG (Dinotefuran)	IRAC 4A	Black Vine Weevil - adults (Otiorhynchus sulcatus - adults)	Hybrid Yew (Taxus X media)	Field Container	Nielsen	OH	2007	Foliar	No efficacy at 8 oz per 100 gal.
26758	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Drench	No significant reduction of feeding damage; no phytotoxicity and growth effect.
26758	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Foliar	No difference among treatment or untreated plants in this study; no phytotoxicity and no impacts on growth.
25522	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Cherry, Sargent (Prunus sargentii)	Field Container	Alm	RI	2006	Foliar	Virtually no control.
26947	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	Excellent control at 8 oz per 100 gal.
26947	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) R. x odorata 'Tiffany'	Field Container	Davis	MI	2009	Drench	Significantly reduced number of adults and feeding damage at 24 oz per 100 gal; inferior to Flagship.
26947	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Julia Child Butter Gold'	Field Container	Schultz	VA	2007	Drench	Excellent efficacy.
27789	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Blackberry (Rubus sp.) R. idaeus	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
26489	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. gracilistyla 'Melanostachys'	Field Container	Braman	GA	2006	Foliar	Poor efficacy.
26489	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. hakuro nishiki	Field Container	Braman	GA	2007	Foliar	Fair efficacy at 8 oz per 100 gal.
25517	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata) 'Bailyei'	Field Container	Alm	RI	2006	Foliar	Good control up to 19 DAT using 6 fl oz per 100 gal.
31714	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Drench	Did not reduce feeding damage with 12 g per DBH inch applied once at dormancy.
27784	Safari 20SG (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Grape (Non-Bearing) (Vitis sp.) 'Seyval Blanc'	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
25735	Safari 20SG (Dinotefuran)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.) V. dentatum	Field Container	Costa	VT	2006	Foliar	Effective control; equal to standard permethrin.
25735	Safari 20SG (Dinotefuran)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field Container	Costa	VT	2007	Foliar	Some reduction in defoliation severity and extent using 4 oz per 100 gal, equivalent level to permethrin.
29868	Safari 20SG (Dinotefuran)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field In-Ground	Weston	NY	2007	Foliar	Effective control at 8 oz per 100 gal; better than imidacloprid.
29295	Safari 20SG (Dinotefuran)	IRAC 4A	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Drench	Excellent control at 60 g per ft height.
33981	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Effective control.
28855	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Joe Pye Weed (Eupatorium maculatum)	Field Container	Kunkel	DE	2008	Foliar	No significant control at 8 oz per 100 gal; data not reliable due to high Control mortality and unfavorable environment.
31586	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Hydrangea (Hydrangea sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Drench	Significantly reduced a high leaf damage with 8 oz per 100 gal applied once; better than Tristar applied once.
32559	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) 'Autumn Joy'	Field Container	Kunkel	DE	2013	Drench	Significantly reduced feeding damage with 24 oz per 100 gal applied once; comparable to the standard Scimitar applied twice.
32559	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) 'Autumn Joy'	Field Container	Kunkel	DE	2013	Foliar	Significantly reduced feeding damage with 8 oz per 100 gal applied once; inferior to the standard Scimitar applied twice.
31578	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012	Drench	Significantly reduced a low leaf damage with 8 oz per 100 gal applied once; comparable to Tristar applied once.
31578	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Drench	Excellent reduction of low to moderate leaf damage with 24 oz per 100 gal; better than Onyx.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31578	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Foliar	Excellent reduction of low to moderate leaf damage with 8 oz per 100 gal; better than Onyx.
31578	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Cloyd	KS	2016	Foliar	Good efficacy with 8 oz per 100 gal applied Jul 1, 8, 15, 22, Aug 12 and Sep 2.
31578	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Frank	NC	2013	Foliar	Results inconclusive due to low infestation; some plant damage reduction with 8 and 24 oz per 100 gal.
31578	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Kunkel	DE	2013	Drench	Did not significantly reduce feeding damage with 24 oz per 100 gal applied once; inferior to the standard Scimitar applied twice.
31578	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 8 oz per 100 gal applied once; inferior to the standard Scimitar applied twice.
32553	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Woodland Sage (Salvia nemorosa) 'Blue Hill'	Field Container	Kunkel	DE	2013	Drench	Reduced feeding damage, though not significantly, with 24 oz per 100 gal applied once; inferior to the standard Scimitar applied twice.
32553	Safari 20SG (Dinotefuran)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Woodland Sage (Salvia nemorosa) 'Blue Hill'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 8 oz per 100 gal applied once; inferior to the standard Scimitar applied twice.
29137	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Drench	Trial not successful - no damage developed after infesting trees with beetles 3 different times.
29874	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Maple, Red (Acer rubrum)	Field Container	Schultz	VA	2008	Trunk spray	Trial 2: Too few attacks for biological significance. Several additional experiments were conducted to refine methodologies for future research.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
30501	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010	Basal spray	Untreated trees not attacked; no conclusions can be made.
30501	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010	Drench	Untreated trees not attacked; no conclusions can be made.
26827	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Kousa (Cornus kousa)	Field Container	Reding	OH	2007	Drench	No infestation; no phytotoxicity.
29492	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Sweet Bay (Magnolia virginiana)	Field Container	Schultz	VA	2009	Drench	No significant control at 24 oz per 100 gal.
31427	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle (Xylosandrus germanus)	Sweet Bay (Magnolia virginiana)	Field Container	Reding	OH	2009	Drench	Reduced X. germanus beetle attacks for < 5 days, though not significantly different from untreated, at 24 oz per 100 gal.
31427	Safari 20SG (Dinotefuran)	IRAC 4A	Ambrosia Beetle (Xylosandrus germanus)	Sweet Bay (Magnolia virginiana)	Field Container	Reding	OH	2009	Trunk spray	Did not reduce X. germanus beetle attacks at 24 oz per 100 gal.
27923	Safari 2G (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Sunsprite'	Field Container	Alm	RI	2008	Broadcast	Effective control at 60 g per plant; equal to standard.
27923	Safari 2G (Dinotefuran)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) R. x odorata 'Tiffany'	Field Container	Davis	MI	2009	Surface of potting media	Significantly reduced number of adults and feeding damage at 2.2 g per gal media; inferior to Flagship.
26613	Safari 2G (Dinotefuran)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.)	Field Container	Costa	VT	2007	Top Dress	No efficacy at 2.2g per gal potting media.
30502	Safari 2G (Dinotefuran)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010	Soil Incorporation	Untreated trees not attacked; no conclusions can be made.
32655	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 22 and 27 fl oz per 100 gal applied once; comparable to the standard Tristar.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
32888	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Pink Velour'	Field Container	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; no significant differences between treatments.
32888	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi' and 'Pink Velour'	Field Container	Addesso	TN	2018	Foliar	Great efficacy with 22 and 27 fl oz per 100 gal applied once; slightly inferior to the standard Tristar.
32642	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Louis Philippe'	Field Container	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; significantly reduced % leaf damage with 22 and 27 fl oz per 100 gal applied twice biweekly; comparable to Tristar.
32642	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Louis Phillippe'	Field Container	Addesso	TN	2017	Foliar	Excellent reduction of feeding damage with 22 and 27 fl oz per 100 gal applied once; comparable to the standard Tristar.
33598	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.)	Greenhouse	Braman	GA	2017	Foliar	Excellent control with 22 and 27 fl oz per 100 gal; comparable to the standard Marathon II.
32642	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Radrazz'	Field Container	Persad	OH	2017	Foliar	100% mortality of adults and significant reduction of leaf damage with 22 and 27 fl oz per 100 gal applied twice biweekly; comparable to Talstar.
34147	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Excellent control with 27 fl oz per 100 gal applied twice biweekly.
32637	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Good control with 22 and 27 fl oz per 100 gal.
33839	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Forsythia (Forsythia sp.)	Field Container	Kunkel	DE	2017	Foliar	Did not significantly reduce feeding damage with 22 fl oz per 100 gal applied twice biweekly.
33160	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Baby Lace'	Field Container	Chong	SC	2017	Foliar	Did not significantly reduce adults nor consistently reduce defoliation with 22 fl oz per 100 gal applied twice biweekly.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
33160	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) Vanilla StrawberryTM	Field Container	Frank	NC	2017	Foliar	Poor beetle toxicity and mediocre feeding damage reduction with 22 fl oz per 100 gal in lab bioassays; inferior to the standard Marathon.
33160	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	Statistically relevant reduction in a low population with 27 fl oz per 100 gal 3 and 7 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
33160	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Pee Gee'	Field Container	Gilrein	NY	2016	Foliar	Excellent control with 22 and 27 fl oz per 100 gal; comparable to the standard Scimitar.
33160	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Bokrathirteen' ('Sweet Summer')	Field Container	Gilrein	NY	2018	Foliar	Excellent control with 22 fl oz per 100 gal applied twice biweekly; comparable to the standard Tristar.
33160	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2019	Foliar	Excellent control with 22 fl oz per 100 gal applied once; comparable to the standard Tristar.
33160	Sarisa 50SL (Cyclaniliprole)	IRAC 28	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Mediocre efficacy with 11 fl oz per 100 gal applied 4 times at 10-17 days interval.
31673	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) R. x odorata 'Tiffany'	Field Container	Davis	MI	2009	Surface of potting media	Significantly reduced number of adults and feeding damage at 2.2 g per gal media; inferior to Flagship.
26493	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. gracilistyla 'Melanostachys'	Field Container	Braman	GA	2006	Foliar	Excellent efficacy.
29299	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Foliar	Excellent control at 5 fl oz per 100 gal + Capsil.
33161	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Baby Lace'	Field Container	Chong	SC	2017	Foliar	Did not significantly reduce adults nor consistently reduce defoliation with 5 fl oz per 100 gal + Capsil applied 3 times weekly.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
33161	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Pee Gee'	Field Container	Gilrein	NY	2016	Foliar	Excellent control with 3.2 fl oz per 100 gal.
32557	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) 'Autumn Joy'	Field Container	Kunkel	DE	2013	Foliar	Significantly reduced feeding damage with 5 fl oz per 100 gal applied twice; best treatment.
32547	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Kunkel	DE	2013	Foliar	Significantly reduced feeding damage with 5 fl oz per 100 gal applied twice; best treatment.
32551	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Woodland Sage (Salvia nemorosa) 'Blue Hill'	Field Container	Kunkel	DE	2013	Foliar	Significantly reduced feeding damage with 5 fl oz per 100 gal applied twice; best treatment.
29494	Scimitar CS (Lambda-cyhalothrin)	IRAC 3A	Ambrosia Beetle (Xylosandrus germanus)	Sweet Bay (Magnolia virginiana)	Field Container	Reding	OH	2009	Trunk spray	Did not reduce X. germanus beetle attacks at 5 fl oz per 100 gal.
35017	Sevin (zeta-cypermethrin - consumer label) (zeta-cypermethrin)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 1: Excellent reduction in adult populations and feeding damage with 4 fl oz per gallon.
34153	Sevin SL (Carbaryl)	IRAC 1A	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Excellent control with 1.5 fl oz per gal applied twice biweekly.
34481	Sevin SL (Carbaryl)	IRAC 1A	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Excellent control with 1.5 fl oz per gal applied twice biweekly.
34325	SP3014 (SP3014)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Poor control with 26 fl oz per 100 gal applied twice.
34325	SP3014 (SP3014)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Foliar	Poor efficacy with 13 and 26 fl oz per 100 gal applied 3 times weekly.
34479	SP3014 (SP3014)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Poor control with 26 fl oz per 100 gal + Capsil applied 3 times weekly.
34479	SP3014 (SP3014)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 2: No reduction in adult viability or feeding damage with 26 fl oz per 100 gal applied weekly.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34380	SP3014 (SP3014)		Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Moonrock'	Field Container	Gilrein	NY	2020	Foliar	Poor efficacy with 26 fl oz per 100 gal applied 3 times weekly; inferior to the standard Tristar. No phytotoxicity.
34380	SP3014 (SP3014)		Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Fire and Ice'	Field Container	Gilrein	NY	2021	Foliar	Good efficacy with 26 fl oz per 100 gal but no effective population reduction with 13 fl oz per 100 gal. No crop injury observed.
34380	SP3014 (SP3014)		Red Headed Flea Beetle (<i>Systema frontalis</i>)	Panicle Hydrangea (<i>Hydrangea paniculata</i>) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Poor efficacy with 13 and 26 fl oz per 100 gal + Capsil applied 4 times at 10-17 days interval.
28157	Talstar Flowable Insecticide/Miticide (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose (<i>Rosa</i> sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	Excellent control at 21.7 fl oz per 100 gal.
28157	Talstar Flowable Insecticide/Miticide (Bifenthrin)	IRAC 3A	Japanese Beetle - adults (<i>Popillia japonica</i> - adults)	Rose (<i>Rosa</i> sp.) 'Radrazz'	Field Container	Persad	OH	2017	Foliar	100% mortality of adults and significant reduction of leaf damage with 10 fl oz per 100 gal applied twice biweekly.
28854	Talstar Flowable Insecticide/Miticide (Bifenthrin)	IRAC 3A	Red Headed Flea Beetle (<i>Systema frontalis</i>)	Joe Pye Weed (<i>Eupatorium maculatum</i>)	Field Container	Kunkel	DE	2008	Foliar	No significant control at 20 oz per 100 gal; data not reliable due to high Control mortality and unfavorable environment.
26138	Talstar Flowable Insecticide/Miticide (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	American Redbud, Eastern Redbud (<i>Cercis canadensis</i>) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Harmony: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
26138	Talstar Flowable Insecticide/Miticide (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	American Redbud, Eastern Redbud (<i>Cercis canadensis</i>) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Pikeville: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
26138	Talstar Flowable Insecticide/Miticide (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (<i>Xylosandrus crassiusculus</i>)	American Redbud, Eastern Redbud (<i>Cercis canadensis</i>)	Field In-Ground	Ludwig	TX	2004	Trunk spray	Results inconclusive due to application timing.
27622	Talstar NF (Bifenthrin)	IRAC 3A	Black Vine Weevil - adults (<i>Otiorhynchus sulcatus</i> - adults)	Hybrid Yew (<i>Taxus X media</i>) 'Densiformis'	Field Container	Nielsen	OH	2007	Foliar	Excellent efficacy at 0.2 lb ai per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
25482	Talstar NF (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 1: Some reduction in attacks per bolt treated with 41 g ai per 100 gal.
25482	Talstar NF (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 3: Almost complete reduction in attacks per bolt treated with 135.3 g ai per 100 gal.
25482	Talstar NF (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 4: Approximately 66% reduction in attacks per bolt treated with 144.9 g ai per 100 gal.
25482	Talstar NF (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - no statistical difference among treatments.
25482	Talstar NF (Bifenthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no statistical difference among treatments.
28000	Tank Mix: Discus + DEET (Imidacloprid + cyfluthrin + DEET)	IRAC 4A + IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Excellent control using 100 oz Discus per 100 gal and 40% DEET.
28000	Tank Mix: Discus + DEET (Imidacloprid + cyfluthrin + DEET)	IRAC 4A + IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - Approximately 80% reduction in bolt attacks but not statistically significant.
28000	Tank Mix: Discus + DEET (Imidacloprid + cyfluthrin + DEET)	IRAC 4A + IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - No statistical difference among treatments.
34239	Tank Mix: Mainspring GNL + Scimitar GC (cyantraniliprole + lambda cyhalothrin)	IRAC 28 + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Drench / Trunk spray	Georgetown, SC: Good control with 4 fl oz/5 fl oz per 100 gal; inferior to the standard Perm-Up.
34247	Tank Mix: Mainspring GNL + Scimitar GC (cyantraniliprole + lambda cyhalothrin)	IRAC 28 + IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Drench / Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
27997	Tank Mix: Onyx + Azatin XL (Bifenthrin + Neem Oil Extract)	IRAC 3A + IRAC 18B	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Approximately 66% reduction in bolt attacks, but not statistical significant.
27998	Tank Mix: Onyx + Celero 16WSG (Bifenthrin + clothianadin)	IRAC 3A + IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Excellent control at 32 oz + 8 oz per 100 gal.
27996	Tank Mix: Onyx + Deet (Bifenthrin + DEET)	IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Great control with 32 oz Onyx + 40% DEET.
27996	Tank Mix: Onyx + Deet (Bifenthrin + DEET)	IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - some reduction in bolt attacks but not statistically significant.
27996	Tank Mix: Onyx + Deet (Bifenthrin + DEET)	IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no statistical difference among treatments.
27999	Tank Mix: Onyx + Dursban (Bifenthrin + chlorpyrifos)	IRAC 3A + IRAC 1B	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 1: Excellent control using 32 oz + 32 oz per 100 gal.
33983	Tank Mix: Sarissa + Azatin O (Cyclaniliprole + azadirachtin)	IRAC 28 +	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Good control.
28003	Tank Mix: Talstar L&T + DEET (Bifenthrin + DEET)	IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - approximately 80% reduction in attacks per bolt, but no statistical significance.
28003	Tank Mix: Talstar L&T + DEET (Bifenthrin + DEET)	IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no statistical difference among treatments.
28001	Tank Mix: Tempo 2 + DEET (Cyfluthrin + DEET)	IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - approximate 90% reduction in attacks per bolt, but not statistically different.
28001	Tank Mix: Tempo 2 + DEET (Cyfluthrin + DEET)	IRAC 3A + ?	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 3: Quincy - no statistical differences among treatments.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
26182	Tempo 20WP (Cyfluthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Monticello - no statistical difference among treatments.
26182	Tempo 20WP (Cyfluthrin)	IRAC 3A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2006	Tree bolt immersion	Experiment 2: Quincy - no statistical difference among treatments.
34149	TetraCURB Concentrate (Rosemary Oil)	IRAC UNE	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Poor control with 128 fl oz per 100 gal applied 3 times weekly.
33787	TetraCURB Concentrate (Rosemary Oil)	IRAC UNE	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	Statistically relevant reduction in a low population with 128 fl oz per 100 gal 3 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
33787	TetraCURB Concentrate (Rosemary Oil)	IRAC UNE	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2019	Foliar	Mediocre control with 128 fl oz per 100 gal applied 3 times weekly; inferior to the standard Tristar.
34150	TetraCURB Organic (Rosemary Oil)	IRAC UNE	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Poor control with 128 fl oz per 100 gal applied 3 times weekly.
33788	TetraCURB Organic (Rosemary Oil)	IRAC UNE	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	No statistically relevant reduction in a low population with 128 fl oz per 100 gal, and no difference in feeding damage from nontreated controls; no injury or impact on growth observed.
27964	Thiodan EC (Endosulfan)	IRAC 2A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Mimosa Silk Tree (Albizia julibrissin)	Field Container	Mizell	FL	2005	Tree bolt immersion	Experiment 1: Very little impact on attacks per bolt treated with 83 g ai per 100 gal.
27720	TickEx EC (Metarhizium anisopliae)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus)	Field In-Ground	Reding	OH	2007	Foliar	No significant reduction of feeding damage at 29 oz per 100 gal; no phytotoxicity and growth effect.
26950	TickEx EC (Metarhizium anisopliae)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Sunsprite'	Field Container	Alm	RI	2008	Foliar	Poor control at 29 fl oz per 100 gal; inferior to standard.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
26950	TickEx EC (Metarhizium anisopliae)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Blushing'	Field Container	Braman	GA	2008	Foliar	No control at 29 fl oz per 100 gal.
26950	TickEx EC (Metarhizium anisopliae)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Julia Child Butter Gold'	Field Container	Schultz	VA	2007	Foliar	Poor efficacy - very slow acting and short residual.
26547	TickEx EC (Metarhizium anisopliae)	IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. hakuro nishiki	Field Container	Braman	GA	2007	Foliar	No efficacy at 29 oz per 100 gal.
25514	TriStar 30SG (Acetamiprid)	IRAC 4A	Flea Beetle, Garden, Epitrix sp. (Epitrix sp.)	Evening Primrose; Sundrops (Oenothera sp.) O. speciosa var. berlandieri	Field Container	Schultz	VA	2006	Foliar	Inconclusive; population too low.
27839	TriStar 30SG (Acetamiprid)	IRAC 4A	Banded Ash Clearwing Borer (Podosesia aureocincta)	Ash (Fraxinus sp.) F. pennsylvanica	Field In-Ground	Nielsen	OH	2008	Trunk spray	Low infestation. No efficacy at 6 ml per inch dbh.
25523	TriStar 30SG (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Cherry, Sargent (Prunus sargentii)	Field Container	Alm	RI	2006	Foliar	100 % control up to 31 DAT.
25736	TriStar 30SG (Acetamiprid)	IRAC 4A	Viburnum Leaf Beetle (Pyrrhalta viburni)	Arrowwood (Viburnum sp.) V. dentatum	Field Container	Costa	VT	2006	Foliar	Effective control; equal to standard permethrin.
29297	TriStar 30SG (Acetamiprid)	IRAC 4A	Peachtree Borer (Synanthedon exitiosa)	Plum (Prunus sp.) P. cistina	Field In-Ground	Nielsen	OH	2009	Foliar	Excellent control at 4 fl oz per inch dbh + Capsil.
31590	TriStar 30SG (Acetamiprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Hydrangea (Hydrangea sp.) 'White Diamonds'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a high leaf damage with 8 g per 100 gal applied once.
31582	TriStar 30SG (Acetamiprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Henry's Garnet'	Field Container	Braman	GA	2012	Foliar	Significantly reduced a low leaf damage with 8 g per 100 gal applied once.
31582	TriStar 30SG (Acetamiprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Cloyd	KS	2016	Foliar	Good efficacy with 12 oz per 100 gal applied Jul 1, 8, 15, 22, Aug 12 and Sep 2.
29140	TriStar 30SG (Acetamiprid)	IRAC 4A	Ambrosia Beetle, Redbay (Xyleborus glabratus)	Redbay (Persea borbonia)	Field In-Ground	Pena	FL	2009	Foliar	Trial not successful - no damage developed after infesting trees with beetles 3 different times.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
29875	TriStar 30SG (Acetamiprid)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Maple, Red (Acer rubrum)	Field Container	Schultz	VA	2008	Trunk spray	Trial 2: Too few attacks for biological significance. Several additional experiments were conducted to refine methodologies for future research.
30503	TriStar 30SG (Acetamiprid)	IRAC 4A	Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Ludwig	TX	2010	Foliar	Untreated trees not attacked; no conclusions can be made.
26759	TriStar 70WSP (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Drench	No significant reduction of feeding damage; no phytotoxicity and growth effect.
26759	TriStar 70WSP (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose-Of-Sharon (Hibiscus syriacus) 'Notwoodone'	Field In-Ground	Reding	OH	2006	Foliar	No difference among treatment or untreated plants in this study; no phytotoxicity and no impacts on growth.
27790	TriStar 70WSP (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Blackberry (Rubus sp.) R. idaeus	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
26490	TriStar 70WSP (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. gracilistyla 'Melanostachys'	Field Container	Braman	GA	2006	Foliar	Excellent efficacy.
26490	TriStar 70WSP (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Willow (Salix sp.) S. hakuro nishiki	Field Container	Braman	GA	2007	Foliar	Provided about 50% efficacy 4 DAT however, there was a significant reduction in feeding although adult japanese beetle were still alive using 96 g per 100 gal.
25518	TriStar 70WSP (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata) 'Bailyei'	Field Container	Alm	RI	2006	Foliar	100 % control up to 31 DAT using 3.38 oz per 100 gal.
27785	TriStar 70WSP (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Grape (Non-Bearing) (Vitis sp.) 'Seyval Blanc'	Field Container	Alm	RI	2007	Foliar	Infestations too low to determine efficacy.
32890	TriStar 8.5SL (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Pink Velour'	Field Container	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; no significant differences between treatments.
32890	TriStar 8.5SL (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi' and 'Pink Velour'	Field Container	Addesso	TN	2018	Foliar	Excellent efficacy with 25.3 fl oz per 100 gal applied once.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
32886	TriStar 8.5SL (Acetamiprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Louis Philippe'	TBD	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; significantly reduced % leaf damage with 25.3 fl oz per 100 gal applied twice biweekly.
33882	TriStar 8.5SL (Acetamiprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	Statistically relevant reduction in a low population with 25 fl oz per 100 gal 3 and 7 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
33882	TriStar 8.5SL (Acetamiprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Bokrathirteen' ('Sweet Summer')	Field Container	Gilrein	NY	2018	Foliar	Excellent control with 25.3 fl oz per 100 gal applied twice biweekly.
33882	TriStar 8.5SL (Acetamiprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2020	Foliar	Great efficacy with 25.3 fl oz per 100 gal applied once. No phytotoxicity.
33882	TriStar 8.5SL (Acetamiprid)	IRAC 4A	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Fire and Ice'	Field Container	Gilrein	NY	2021	Foliar	Excellent efficacy with 25.3 fl oz per 100 gal. No crop injury observed.
33845	UP-Star Gold (Bifenthrin)	IRAC 3A	Red Headed Flea Beetle (Systema frontalis)	Forsythia (Forsythia sp.)	Field Container	Kunkel	DE	2017	Foliar	Significantly reduced feeding damage with 8 fl oz per 100 gal applied twice biweekly; best product.
33609	V-10433 (V-10433)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi' and 'Pink Velour'	Field Container	Addesso	TN	2018	Foliar	Mediocre efficacy with 11 and 22 fl oz per 100 gal applied twice; much inferior to the standard Tristar.
34326	V-10433 (V-10433)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica)	Shadehouse/ Lath House	Addesso	TN	2020	Foliar	Poor control with 13 fl oz per 100 gal applied twice.
34326	V-10433 (V-10433)		Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Hopi'	Shadehouse/ Lath House	Addesso	TN	2021	Foliar	Good efficacy with 13 fl oz per 100 gal applied 5 times at 3-4 days interval; almost comparable to the standard Onyx.
34151	V-10433 (V-10433)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field Container	Potter	KY	2019	Foliar	Poor control with 22 fl oz per 100 gal applied 3 times weekly.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34480	V-10433 (V-10433)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2020	Foliar	Poor control with 11 fl oz per 100 gal applied 4 times every 3-4 days.
34480	V-10433 (V-10433)		Japanese Beetle - adults (Popillia japonica - adults)	Linden, Shamrock (Tilia cordata)	Field In-Ground	Potter	KY	2021	Foliar	Experiment 2: No reduction in adult viability or feeding damage with 11 fl oz per 100 gal applied every 3 - 4 days.
33792	V-10433 (V-10433)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Vanilla Strawberry'	Field Container	Frank	NC	2019	Foliar	Statistically relevant reduction in a low population with 22 fl oz per 100 gal 3 d after treatment, but no differences in feeding damage from nontreated controls; no injury or impact on growth observed.
33792	V-10433 (V-10433)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Bokrathirteen' ('Sweet Summer')	Field Container	Gilrein	NY	2018	Foliar	Poor control with 11 and 22 fl oz per 100 gal applied 3 times weekly.
33792	V-10433 (V-10433)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Moonrock'	Field Container	Gilrein	NY	2019	Foliar	Mediocre control with 22 fl oz per 100 gal applied 3 times weekly; inferior to the standard Tristar.
33792	V-10433 (V-10433)		Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Fire and Ice'	Field Container	Gilrein	NY	2021	Foliar	No effective population reduction with 11 fl oz per 100 gal. No injury observed.
33642	V-10433 (V-10433)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Harmony: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33642	V-10433 (V-10433)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis) 'Forest Pansy'	Field In-Ground	Chong	NC	2018	Trunk	Pikeville: Data collected were insufficient to make conclusions on efficacy of treatments; no phytotoxicity.
33879	V-10433 (V-10433)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	American Redbud, Eastern Redbud (Cercis canadensis)	Field Container	Villanassery Joseph	GA	2019	Trunk spray	Efficacy unknown; No statistical difference among treatments
34238	V-10433 (V-10433)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida) 'Cherokee Princess'	Field Container	Chong	SC	2019	Trunk spray	Georgetown, SC: No significant control with 5.5 and 11 fl oz per 100 gal.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
34246	V-10433 (V-10433)		Ambrosia Beetle, Granulate/Asian (Xylosandrus crassiusculus)	Dogwood, Flowering (Cornus florida)	Field In-Ground	Chong	SC	2019	Trunk spray	Low infestation using trees without flooding or ethanol stress; results inconclusive
34966	Velifer (Beauveria bassiana strain PPRI 5339)	IRAC UNF	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Fire and Ice'	Field Container	Gilrein	NY	2021	Foliar	Some mortality observed with 13 fl oz per 100 gal. No injury observed.
32662	Venerate CG (Burkholderia rinojensis strain A396)	FRAC NC & IRAC UNB	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 1 gal per 100 gal + NuFilm applied 3 times weekly; comparable to the standard Tristar.
33145	VST-006350 (Liquid) (GS-omega/kappa-Hxtx-Hv1a)	IRAC 32	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Louis Phillippe'	Field Container	Addesso	TN	2017	Foliar	Did not reduce feeding damage with 1 liter per 30 gal applied twice.
33840	VST-006350 (Liquid) (GS-omega/kappa-Hxtx-Hv1a)	IRAC 32	Red Headed Flea Beetle (Systema frontalis)	Forsythia (Forsythia sp.)	Field Container	Kunkel	DE	2017	Foliar	Did not reduce feeding damage with 3.33 liters per 100 gal + LI-700 applied 3 times weekly.
33146	VST-006350 (Liquid) (GS-omega/kappa-Hxtx-Hv1a)	IRAC 32	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Baby Lace'	Field Container	Chong	SC	2017	Foliar	Did not significantly reduce adults nor consistently reduce defoliation with 3.33 L per 100 gal + LI 700 applied 3 times weekly.
33146	VST-006350 (Liquid) (GS-omega/kappa-Hxtx-Hv1a)	IRAC 32	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) Vanilla StrawberryTM	Field Container	Frank	NC	2017	Foliar	Poor beetle toxicity and feeding damage reduction with 2 lb + Capsil per 100 gal in lab bioassays; inferior to the standard Marathon.
33146	VST-006350 (Liquid) (GS-omega/kappa-Hxtx-Hv1a)	IRAC 32	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata)	Field Container	Gilrein	NY	2016	Foliar	Poor control with 94.6 L per 100 gal.
32663	Xpectro OD (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Sri Lankan Weevil (Myllocerus undatus)	Shoeblackplant; Chinese hibiscus (Hibiscus rosa-sinensis) 'Double Peach'	Field Container	Dale	FL	2016	Foliar	Significantly reduced leaf feeding damage with 32 oz per 100 gal applied twice weekly; inferior to the standard Tristar.
32889	Xpectro OD (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Crape Myrtle (Lagerstroemia indica) 'Pink Velour'	Field Container	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; no significant differences between treatments.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
32643	Xpectro OD (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Louis Philippe'	Field Container	Addesso	TN	2016	Foliar	Number of beetles and % leaf damage too low for reliable evaluations; significantly reduced % leaf damage with 32 fl oz per 100 gal applied twice weekly; inferior to Tristar.
33599	Xpectro OD (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.)	Greenhouse	Braman	GA	2017	Foliar	Poor control with 32 fl oz per 100 gal; inferior to the standard Marathon II.
32643	Xpectro OD (Pyrethrins + Beauveria bassiana Strain GHA)	IRAC 3A + IRAC UNF	Japanese Beetle - adults (Popillia japonica - adults)	Rose (Rosa sp.) 'Radrazz'	Field Container	Persad	OH	2017	Foliar	83% mortality of adults and significant reduction of leaf damage with 32 fl oz per 100 gal applied twice biweekly; inferior to Talstar.
33979	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Maple, Red (Acer rubrum)	Field Container	Frank	NC	2016	Foliar	Effective control.
32640	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Panicle Hydrangea (Hydrangea paniculata) 'Limelight'	Field Container	Sadof	IN	2017	Foliar	Mediocre efficacy with 2.5 oz per 100 gal applied 4 times at 10-17 days interval.
32556	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) Hydrangea paniculata 'Pee Gee'	Field Container	Gilrein	NY	2016	Foliar	Excellent control with 3.5 oz per 100 gal + Capsil; comparable to the standard Scimitar.
32556	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Witch's Moneybags (Hylotelephium telephium) 'Autumn Joy'	Field Container	Kunkel	DE	2013	Foliar	Significantly reduced feeding damage with 2.75 and 3.5 oz per 100 gal + Capsil applied twice; slightly inferior to the standard Scimitar.
31891	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Braman	GA	2013	Foliar	Mediocre reduction of low to moderate leaf damage with 3.5 oz per 100 gal.
31891	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica)	Field Container	Frank	NC	2013	Foliar	Results inconclusive due to low infestation.

PR#	Product (Active Ingredients)	MOA Class	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31891	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Virginia Sweetspire (Itea virginica) 'Little Henry'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 2.75 and 3.5 oz per 100 gal + Capsil applied twice; inferior to the standard Scimitar.
32550	Xxpire 40WG (Spinetoram + sulfoxaflor)	IRAC 5 + IRAC 4C	Red Headed Flea Beetle (Systema frontalis)	Woodland Sage (Salvia nemorosa) 'Blue Hill'	Field Container	Kunkel	DE	2013	Foliar	Did not significantly reduce feeding damage with 2.75 and 3.5 oz per 100 gal + Capsil applied twice; inferior to the standard Scimitar.
31710	Xytect 2F (Imidacloprid)	IRAC 4A	European Elm Flea Weevil (Orchestes alni)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Drench	Good control with 0.2 fl oz per DBH inch applied once at dormancy. Best treatment.
31715	Xytect 2F (Imidacloprid)	IRAC 4A	Japanese Beetle - adults (Popillia japonica - adults)	Elm (Ulmus sp.) 'Patriot'	Field In-Ground	Jones	WI	2012	Drench	Significantly reduced feeding damage with 0.2 fl oz per DBH inch applied once at dormancy.

Table 112. Summary of Efficacy By Product – White Grubs and Weevils

Note: Table entries are sorted by crop Latin name. Only those trials received by 5/12/2010 are included in the table below.

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
31409	A16901B 45WG (Cyantraniliprole + thiamethoxam)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field In-Ground	Freiberger	NJ	2012	Broadcast & water-in	Excellent efficacy with 0.095 g product per 9 sq ft using 0.5 gal water.	N	20130412a.pdf
30397	A16901B 45WG (Cyantraniliprole + thiamethoxam)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2010	Drench	Significant reduction in grub population at 0.095g per 9 sq ft drench; no impact on plant height or width.	N	20110331h.pdf
25123	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Holly (Ilex sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Excellent reduction of grubs with Aug drench of 0.8 oz per 100 gal; great control with Nov application.	Y	20080522a.pdf
29498	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Oak, White (Quercus alba)	Field Container	Reding	OH	2009	Drench	100 % control at 32 fl oz per 100 gal.	N	20091130p.pdf
26936	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Rhododendron (Rhododendron sp.) 'Scintillation'	Field Container	Alm	RI	2008	Drench	Excellent control at 0.08 fl oz per 100 gal; equal to standard.	Y	20081113b.pdf
26761	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field Container	Reding	OH	2006	Drench	No infestation developed; no phytotoxicity or impact on growth.	Y	20070410b.pdf
26761	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris) S. chinensis	Field Container	Reding	OH	2009	Drench	100 % control at 32 fl oz per 100 gal.	Y	20091130p.pdf
31669	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Yew (Taxus media) 'Nigra'	Field Container	Alm	RI	2006	Drench	Excellent control of oriental beetle larvae.	N	20061212w.pdf
25124	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Good reduction of grubs with Aug drench of 0.8 oz per 100 gal; good control with Nov application.	Y	20080522a.pdf
25124	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2009	Drench	An 87% reduction in grub counts using 0.8 fl oz per 100 gal drench.	Y	20110324a.pdf
25124	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2010	Drench	Significant reduction in grub population at 0.8 fl oz per 9 sq ft drench; no impact on plant height or width.	Y	20110331h.pdf
29117	Acelepryn 1.67 (Chlorantraniliprole)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Drench	37 % to not significant control at 0.8 and 3.2 fl oz per 100 gal.	N	20090429a.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
26469	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Drench	Great control with 6.5 fl oz per 100 gal, but poor control with 0.8 fl oz per 100 gal as curative drenches.	N	20070418g.pdf
26469	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Soil Incorporation	Excellent efficacy with 5, 10, and 20 ppm preplant soil incorporation.	N	20070418g.pdf
26469	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Drench	Poor control at 0.8 fl oz per 100 gal.	N	20080910c.pdf
26469	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Poor control at 1, 2 and 4 ppm.	N	20080910c.pdf
26754	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Stonecrop (Sedum spurium) 'Vera Jameson'	Field Container	Reding	OH	2006	Drench	0.8 oz ai per 100 gal; excellent efficacy on larvae; no significant reduction of adult feeding damage; no phytotoxicity; plant not significantly taller than Nontreated.	N	20070412m.pdf
26091	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Drench	Excellent efficacy with 16 oz per 100 gal drenched pre-infestation.	N	20070228a.pdf
26091	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Yew (Taxus media) 'Densifomis'	Field Container	Nielsen	OH	2007	Drench	Excellent efficacy at 0.8 fl oz per 100 gal.	N	20071220b.pdf
26091	Acelepryn 1.67 (Chlorantraniliprole)	Black Vine Weevil - grubs (Otorhynchus sulcatus - grubs)	Yew (Taxus media) 'Nigra'	Field Container	Alm	RI	2006	Drench	No significant control of black vine weevil.	N	20061212w.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
27611	Acelepryn 1.67 (Chlorantraniliprole)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2007	Drench	Low infestation levels and no statistical significance among treatments.	N	20071219p.pdf
26495	Acelepryn 1.67 (Chlorantraniliprole)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2006	Drench	No statistical difference among treatments.	Y	20070411a.pdf
26495	Acelepryn 1.67 (Chlorantraniliprole)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2007	Drench	Only ~10 % grub survival; no statistical difference among treatments.	Y	20080116f.pdf
27916	Arena 50WDG (Clothianadin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Drench	No efficacy using 1.28 oz per 100 gal curative drench.	N	20070418g.pdf
27916	Arena 50WDG (Clothianadin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Excellent control at 49 mg per pot.	N	20080910c.pdf
31410	AzaGuard (Azadirachtin)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field In-Ground	Freiberger	NJ	2012	Drench	Poor efficacy with 16 fl oz per 100 gal using 0.5 gal diluent per 9 sq ft.	N	20130412a.pdf
26938	BAS 320i (Metaflumizone)	Oriental Beetle (Anomala orientalis)	Rhododendron (Rhododendron sp.) 'Scintillation'	Field Container	Alm	RI	2008	Drench	100 % control at 16 fl oz per 100 gal; equal to standard.	N	20081113b.pdf
30336	BAS 320i (Metaflumizone)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2009	Drench	A 24% reduction in grub counts using 16 oz per 100 gal.	N	20110324a.pdf
29113	BAS 320i (Metaflumizone)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Drench	No efficacy at 16 fl oz per 100 gal.	N	20090429a.pdf
26471	BAS 320i (Metaflumizone)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Drench	Excellent efficacy using 50 and 100 ppm as curative drenches.	N	20070418g.pdf
26471	BAS 320i (Metaflumizone)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Soil Incorporation	Excellent efficacy with 25, 50, 100, and 200 ppm preplant soil incorporation.	N	20070418g.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
26471	BAS 320i (Metaflumizone)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Drench	Excellent control at 16 fl oz per 100 gal.	N	20080910c.pdf
26471	BAS 320i (Metaflumizone)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Excellent control at 4 ppm, less effective at 1 and 2 ppm.	N	20080910c.pdf
27614	BAS 320i (Metaflumizone)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Rhododendron (Rhododendron sp.) R. roseum elegans	Field Container	Nielsen	OH	2007	Drench	Excellent efficacy at 16 oz per 100 gal.	N	20071220b.pdf
27610	BAS 320i (Metaflumizone)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2007	Drench	Low infestation levels and no statistical significance among treatments.	N	20071219p.pdf
27675	BAS 320i (Metaflumizone)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2007	Drench	Only ~10 % grub survival; no statistical difference among treatments.	N	20080116f.pdf
27917	BotaniGard ES (BioWorks) (Beauveria bassiana)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Drench	Poor control at 18.7 mg per pot WP or 39 uL per pot ES formulations.	N	20080910c.pdf
27917	BotaniGard ES (BioWorks) (Beauveria bassiana)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Poor control at 18.7 mg per pot WP or 39 uL per pot ES formulations.	N	20080910c.pdf
25119	Celero 16WSG (Clothianidin)	Oriental Beetle (Anomala orientalis)	Holly (Ilex sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Good reduction of grubs with Aug drench of 0.5 oz per 100 gal; good control with Nov application.	Y	20080522a.pdf
26760	Celero 16WSG (Clothianidin)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field Container	Reding	OH	2006	Drench	No infestation developed; no phytotoxicity or impact on growth.	Y	20070410b.pdf
31670	Celero 16WSG (Clothianidin)	Oriental Beetle (Anomala orientalis)	Yew (Taxus media) 'Nigra'	Field Container	Alm	RI	2006	Drench	Good control of oriental beetle larvae.	N	20061212w.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
25120	Celero 16WSG (Clothianidin)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Good reduction of grubs with Aug drench of 0.5 oz per 100 gal; good control with Nov application.	Y	20080522a.pdf
26753	Celero 16WSG (Clothianidin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Stonecrop (Sedum spurium) 'Vera Jameson'	Field Container	Reding	OH	2006	Drench	0.64 oz ai per 100 gal; excellent efficacy on larvae; significantly reduced adult feeding damage; no phytotoxicity; plant significantly taller than Nontreated.	Y	20070412m.pdf
26090	Celero 16WSG (Clothianidin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Drench	Excellent efficacy with 20 oz per 100 gal drenched pre-infestation.	Y	20070228a.pdf
26090	Celero 16WSG (Clothianidin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media) 'Densiformis'	Field Container	Nielsen	OH	2007	Drench	Excellent efficacy at 16 oz per 100 gal.	Y	20071220b.pdf
26090	Celero 16WSG (Clothianidin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media) 'Nigra'	Field Container	Alm	RI	2006	Drench	No control of black vine weevil.	Y	20061212w.pdf
27609	Celero 16WSG (Clothianidin)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2007	Drench	Low infestation levels and no statistical significance among treatments.	N	20071219p.pdf
26494	Celero 16WSG (Clothianidin)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2006	Drench	No statistical difference among treatments.	Y	20070411a.pdf
26494	Celero 16WSG (Clothianidin)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2007	Drench	Only ~10 % grub survival; no statistical difference among treatments.	Y	20080116f.pdf
25125	Discus (Imidacloprid + cyfluthrin)	Oriental Beetle (Anomala orientalis)	Holly (Ilex sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Excellent reduction of grubs with Aug drench of 0.8 oz per 100 gal; good control with Nov application.	Y	20080522a.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
25126	Discus (Imidacloprid + cyfluthrin)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Some reduction of grubs with Aug drench of 0.8 oz per 100 gal; no statistical difference from Nontreated with Nov application.	Y	20080522a.pdf
25126	Discus (Imidacloprid + cyfluthrin)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2010	Drench	Some reduction in grub population at 17 fl oz per 3000 sq ft drench; no impact on plant height or width.	Y	20110331h.pdf
26857	Discus (Imidacloprid + cyfluthrin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Drench	No control with 13 fl oz per 100 gal as curative drench.	Y	20070418g.pdf
27613	DuraGuard (Chlorpyrifos)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2007	Drench	Low infestation levels and no statistical significance among treatments.	N	20071219p.pdf
29110	Dylox 80 SP (Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Drench	93 % control at 3.75 oz per 100 gal.	Y	20090429a.pdf
31412	Flagship 0.22G (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field In-Ground	Freiberger	NJ	2012	Broadcast & water-in	Poor efficacy with 11.25 g product per 9 sq ft.	Y	20130412a.pdf
30337	Flagship 0.22G (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2009	Broadcast	A 75% reduction in grub counts using 11.25 g product per 9 sq ft.	Y	20110324a.pdf
30337	Flagship 0.22G (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2010	Broadcast	Good reduction in grub population at 11.25 g per 9 sq ft drench; no impact on plant height or width.	Y	20110331h.pdf
29115	Flagship 0.22G (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Broadcast	47 % control at 6 g per pot.	Y	20090429a.pdf
25121	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Holly (Ilex sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Excellent reduction of grubs with Aug drench of 10 oz per 100 gal; excellent control with Nov application.	Y	20080522a.pdf
29500	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Oak, White (Quercus alba)	Field Container	Reding	OH	2009	Drench	100 % control at 8 oz per 100 gal.	Y	20091130p.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
29499	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field Container	Freiberger	NJ	2012	Drench	Mediocre efficacy with 0.1 g product per 9 sq ft in 0.5 gal water.	Y	20130412a.pdf
29499	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris) S. chinensis	Field Container	Reding	OH	2009	Drench	100 % control at 8 oz per 100 gal.	Y	20091130p.pdf
25122	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2005	Drench	Some reduction of grubs with Aug drench of 10 oz per 100 gal; no statistical difference from Nontreated with Nov application.	Y	20080522a.pdf
25122	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2009	Drench	An 87% reduction in grub counts using 0.1 g product per 9 sq ft in ½ gal water.	Y	20110324a.pdf
25122	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2010	Drench	Significant reduction in grub population at 0.1 g per 9 sq ft drench; no impact on plant height or width.	Y	20110331h.pdf
29116	Flagship 25WG (Thiamethoxam)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Drench	46 and 40 % control at 8 and 17 oz per 100 gal.	Y	20090429a.pdf
26858	Flagship 25WG (Thiamethoxam)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Drench	No control with 8 oz per 100 gal as curative drench.	Y	20070418g.pdf
26096	Flagship 25WG (Thiamethoxam)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Drench	Great efficacy with 8 oz per 100 gal drench pre-infestation.	Y	20070228a.pdf
25032	Flagship 25WG (Thiamethoxam)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2005	Drench	Little efficacy using 0.18 oz per 1000 sq ft drenched post-infestation.	N	20060626a.pdf
26497	Flagship 25WG (Thiamethoxam)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2006	Drench	No statistical difference among treatments.	Y	20070411a.pdf
30399	Grandevo (MBI 203 DF) (Chromobacterium subtsugae NRRL B-30655)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2010	Drench	No impact on grub population at 4 fl oz per tree; no impact on plant height or width.	N	20110331h.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
27920	Hachi-Hachi EC (Tolfenpyrad)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Inconclusive data due to low Nontreated population; excellent root rating at 10 ppm.	N	20080910c.pdf
29114	Hexacide (Rosemary Oil)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Drench	31 % control at 1.5 qt per 100 gal.	N	20090429a.pdf
26498	Mach 2 Granular (Halofenazide)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2006	Broadcast to media surface	No statistical difference among treatments.	N	20070411a.pdf
26097	Mach 2 Liquid (Halofenazide)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Drench	Little efficacy with 2 lb ai per acre drenched pre-infestation.	N	20070228a.pdf
25034	Mach 2 Liquid (Halofenazide)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2005	Drench	Little efficacy using 2.9 fl oz per 1000 sq ft drenched post-infestation.	N	20060626a.pdf
26093	Marathon 1% granular (Imidacloprid)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Soil Incorporation	Good efficacy with 0.1 g ai per gal soil as soil incorporation.	Y	20070228a.pdf
25035	Marathon 60WP (Imidacloprid)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2005	Drench	Little efficacy using 20 g per 3000 sq ft drenched post-infestation.	N	20060626a.pdf
30338	Marathon II (Imidacloprid)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field Container	Freiberger	NJ	2009	Drench	An 86% reduction in grub counts using 0.6 fl oz per 1,000 sq ft in 10 gal water.	Y	20110324a.pdf
26824	Met52 (Metarhizium anisopliae strain F52)	Oriental Beetle (Anomala orientalis)	Dogwood, Kousa (Cornus kousa)	Field Container	Reding	OH	2007	Drench	No infestation; no phytotoxicity.	N	20080128h.pdf
26939	Met52 (Metarhizium anisopliae strain F52)	Oriental Beetle (Anomala orientalis)	Rhododendron (Rhododendron sp.) 'Scintillation'	Field Container	Alm	RI	2008	Soil Incorporation	No significant control at 20 g per 4 gal media.	N	20081113b.pdf
27722	Met52 (Metarhizium anisopliae strain F52)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field Container	Reding	OH	2007	Drench	No infestation; no phytotoxicity.	N	20080128h.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
26467	Met52 (Metarhizium anisopliae strain F52)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Soil Incorporation	No efficacy using 450,000,000 spores per L.	Y	20070418g.pdf
26467	Met52 (Metarhizium anisopliae strain F52)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Drench	Inconclusive data due to low Nontreated population; excellent root rating at 2.9 g per pot.	Y	20080910c.pdf
26467	Met52 (Metarhizium anisopliae strain F52)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Inconclusive data due to low Nontreated population; excellent root rating at 2.9 g per pot.	Y	20080910c.pdf
26092	Met52 (Metarhizium anisopliae strain F52)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Soil Incorporation	Little efficacy with 6.25 g per pot incorporated into soil pre-infestation.	Y	20070228a.pdf
26092	Met52 (Metarhizium anisopliae strain F52)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media) 'Densifloris'	Field Container	Nielsen	OH	2007	Soil Incorporation	Poor efficacy at 6.25 g per pot.	Y	20071220b.pdf
27612	Met52 (Metarhizium anisopliae strain F52)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2007	Drench	Low infestation levels and no statistical significance among treatments.	N	20071219p.pdf
27674	Met52 (Metarhizium anisopliae strain F52)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2007	Drench	Only ~10 % grub survival; no statistical difference among treatments	N	20080116f.pdf
30398	NI-CH001 (NI-CH001)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field In-Ground	Freiberger	NJ	2010	Drench	No impact on grub population at 1 oz per gal; no impact on plant height or width.	N	20110331h.pdf
26790	Ornazin 3%EC (Azadirachtin)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2005	Drench	Little efficacy using 10 oz per 100 gal drenched post-infestation.	N	20060626a.pdf
25279	Orthene TTO 97 (Valent) (Acephate)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2005	Drench	Excellent efficacy using 12 oz per 100 gal drenched post-infestation.	Y	20060626a.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
26094	Precise Acephate (Acephate)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Soil Incorporation	Little efficacy wit 6 g ai per pot incorporated into soil pre-infestation.	Y	20070228a.pdf
26499	Precise Acephate (Acephate)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2006	Broadcast to media surface	No statistical difference among treatments.	Y	20070411a.pdf
26823	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Dogwood, Kousa (Cornus kousa)	Field Container	Reding	OH	2007	Drench	No infestation; no phytotoxicity.	Y	20080128h.pdf
29501	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Oak, White (Quercus alba)	Field Container	Reding	OH	2009	Drench	100 % control at 24 oz per 100 gal.	N	20091130p.pdf
26937	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Rhododendron (Rhododendron sp.) 'Scintillation'	Field Container	Alm	RI	2008	Drench	100 % control at 24 oz per 100 gal; equal to standard.	Y	20081113b.pdf
25796	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field Container	Reding	OH	2005	Drench	Excellent control at 6, 12 and 24 oz per 100 gal.	Y	20060214a.pdf
25796	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field Container	Reding	OH	2006	Drench	No infestation developed; no phytotoxicity or impact on growth.	Y	20070410b.pdf
25796	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field Container	Reding	OH	2007	Drench	No infestation; no phytotoxicity.	Y	20080128h.pdf
25796	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris) S. chinensis	Field Container	Reding	OH	2009	Drench	100 % control at 24 oz per 100 gal.	Y	20091130p.pdf
31671	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Yew (Taxus sp.) 'Nigra'	Field Container	Alm	RI	2006	Drench	Excellent control of oriental beetle larvae.	N	20061212w.pdf
30339	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field Container	Freiberger	NJ	2009	Drench	A 21% reduction in grub counts using 12 grams per inch dbh for trees.	N	20110324a.pdf
29112	Safari 20SG (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Drench	No significant control at 24 oz per 100 gal.	N	20090429a.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
26470	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Drench	Good efficacy using 24 oz per 100 gal.	Y	20070418g.pdf
26470	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Excellent control at 24 oz per 100 gal.	Y	20080910c.pdf
25795	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Spruce, White; Cat (Picea glauca)	Field Container	Reding	OH	2005	Drench	No infestation developed.	Y	20060214a.pdf
25200	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Azalea, & Rhododendron (Rhododendron sp.) 'Nova Zembla'	Field Container	Reding	OH	2004	Drench	Very little efficacy at 0.75, 1.5 and 3.0 lb product per 100 gal when drenched onto mature and pupating larvae; no injury at any rate.	Y	20080522b.pdf
25201	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Stonecrop (Sedum sp.) 'Vera Jameson'	Field Container	Reding	OH	2004	Drench	Excellent efficacy at 0.75, 1.5, and 3.0 lb product per 100 gal when drenched shortly before adults laid eggs; no injury at any rate.	Y	20080522c.pdf
25201	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Stonecrop (Sedum sp.) 'Vera Jameson'	Field Container	Reding	OH	2005	Drench	No efficacy at 6, 12, and 24 oz product per 100 gal with drench application prior to adults laying eggs; no injury.	Y	20060214a.pdf
26755	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Stonecrop (Sedum spurium) 'Vera Jameson'	Field Container	Reding	OH	2006	Drench	4.8 oz ai per 100 gal; excellent efficacy on larvae; significantly reduced adult feeding damage; no phytotoxicity; plant significantly taller than Nontreated	Y	20070412m.pdf
25432	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus sp.)	Field Container	Nielsen	OH	2006	Drench	Excellent efficacy with 6.8 g ai per pot drenched pre-infestation.	Y	20070228a.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
25432	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus sp.) 'Brownii'	Field Container	Reding	OH	2004	Foliar	Good to excellent efficacy at 0.75, 1.5, and 3.0 lb product per 100 gal with foliar application shortly before adults laid eggs; no injury at any rate.	Y	20080522d.pdf
25432	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus sp.) 'Brownii'	Field Container	Reding	OH	2005	Drench	No larvae in any treatment at final rating; no injury.	Y	20060214a.pdf
25432	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus sp.) 'Densiformis'	Field Container	Nielsen	OH	2007	Drench	Excellent efficacy at 12 oz per 100 gal.	Y	20071220b.pdf
25432	Safari 20SG (Dinotefuran)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus sp.) 'Nigra'	Field Container	Alm	RI	2006	Drench	No significant control of black vine weevil.	Y	20061212w.pdf
27608	Safari 20SG (Dinotefuran)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2007	Drench	Low infestation levels and no statistical significance among treatments.	N	20071219p.pdf
26496	Safari 20SG (Dinotefuran)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2006	Drench	No statistical difference among treatments.	Y	20070411a.pdf
30340	Safari 2G (V-10112 2G) (Dinotefuran)	Oriental Beetle (Anomala orientalis)	Arborvitae (Thuja sp.)	Field Container	Freiberger	NJ	2009	Broadcast	A 29% reduction in grub counts using 120 grams per inch dbh for tree.	Y	20110324a.pdf
26609	Safari 2G (V-10112 2G) (Dinotefuran)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.) 'Idea'	Field Container	Cowles	CT	2008	Soil Incorporation	Poor control at 1.23 g per pot.	Y	20080910c.pdf
27616	Safari 2G (V-10112 2G) (Dinotefuran)	Black Vine Weevil - grubs (Otiorynchus sulcatus - grubs)	Yew (Taxus media) 'Densiformis'	Field Container	Nielsen	OH	2007	Soil Incorporation	Excellent efficacy at 2.2 g per pot.	Y	20071220b.pdf

PR#	Product (Active Ingredients)	Target	Crop	Production Site	Researcher	Trial State	Trial Year	Application Type	Results	EPA Reg	File Name
25033	Talstar Flowable Insecticide/Miticide (Bifenthrin)	Strawberry Rootworm (Paria fragariae ssp. Fragariae)	Azalea (Rhododendron sp.)	Field Container	Hesselein	AL	2005	Drench	Highly efficacious using 25 fl oz per 100 gal drenched post-infestation.	N	20060626a.pdf
26500	Talstar Flowable Insecticide/Miticide (Bifenthrin)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2006	Drench	No statistical difference among treatments.	N	20070411a.pdf
26500	Talstar Flowable Insecticide/Miticide (Bifenthrin)	Japanese Beetle - grubs (Popillia japonica - grubs)	Rose (Rosa sp.) 'Caldwell Pink'	Field Container	Braman	GA	2007	Drench	Only ~10 % grub survival; no statistical difference among treatments.	N	20080116f.pdf
27969	Talstar NG (Bifenthrin)	Oriental Beetle (Anomala orientalis)	Rhododendron (Rhododendron sp.) 'Scintillation'	Field Container	Alm	RI	2008	Soil Incorporation	Standard treatment; 100 % control at 33.7 g per gal media.	Y	20081113b.pdf
26856	Talstar NG (Bifenthrin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Strawberry (Non-Bearing) (Fragaria sp.)	Field Container	Cowles	CT	2007	Soil Incorporation	Excellent control with 10 ppm preplant soil incorporation.	Y	20070418g.pdf
26095	Talstar NG (Bifenthrin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media)	Field Container	Nielsen	OH	2006	Soil Incorporation	Poot efficacy with 25 ppm incorporated into soil pre-infestation.	Y	20070228a.pdf
26095	Talstar NG (Bifenthrin)	Black Vine Weevil - grubs (Otiorhynchus sulcatus - grubs)	Yew (Taxus media) 'Densiformis'	Field Container	Nielsen	OH	2007	Soil Incorporation	Poor efficacy at 25 ppm ai.	Y	20071220b.pdf
31411	Tank Mix: AzaGuard + TerraClean 5.0 (azadirachtin +)	Oriental Beetle (Anomala orientalis)	Lilac, Common (Syringa vulgaris)	Field In-Ground	Freiberger	NJ	2012	Drench	Poor efficacy with 16 fl oz + 1 gal per 100 gal using 0.5 gal diluent per 9 sq ft.	N	20130412a.pdf
29111	TickEx EC (Metarhizium anisopliae)	Oriental Beetle (Anomala orientalis)	Turf (Turf)	Greenhouse	Gilrein	NY	2007	Drench	No efficacy at 21 and 29 fl oz per 100 gal.	N	20090429a.pdf

Label Suggestions

Based upon data accumulated through the IR-4 research program in 2005-2021, we suggest that registrants consider the following updates to their current product labels:

- Arena/Celero
 - add drench applications at time of male oriental beetle flight to these labels
 - add viburnum leaf beetle to these labels
 - if additional favorable data are available, add soil incorporation, pre-infestation drench and post-infestation drench to the Celero 20SG label to control black vine weevil larvae
- BAS 320i
 - if additional favorable data are available, add Japanese beetle adults to the initial label
 - if additional favorable data are available, add viburnum leaf beetle to the initial label
 - if additional favorable data are available, add black vine weevil to the initial label
- DPX-E2Y45/Acelepryn
 - add drench applications at time of male oriental beetle flight to the initial label
 - include both Japanese beetle adults and larvae at appropriate rates in the initial label for ornamental horticulture crops
 - add viburnum leaf beetle on the label at 16 oz per 100 gal
 - if additional favorable data are available, add soil incorporation and pre-infestation drench to the label to control black vine weevil larvae
- Hachi-Hachi/Tolfenpyrad
 - if additional favorable data are available, add Japanese beetle adults to the initial label
 - if additional favorable data are available, add viburnum leaf beetle to the initial label
- Safari 20SG
 - add pre-infestation drench to the label to control black vine weevil larvae
 - add viburnum leaf beetle to the existing label at 8 oz per 100 gal
 - if additional favorable data are available, add drench applications at time of male oriental beetle flight to this label
- TriStar
 - if additional favorable data are available, add viburnum leaf beetle to the label

Appendix 1: Contributing Researchers

Dr. Karla M Addesso	Tennessee State University Otis L. Floyd Nursery Research Center 472 Cadillac Lane McMinnville TN 37110
Dr. Scott Alm	University of Rhode Island Dept. of Plant Science 9 E. Alumni Ave., Suite 7 Kingston, RI02881
Dr. Eileen Buss	University of Florida Entomology & Nematology Department P. O. Box 110620 Gainesville, FL 32611-0620
Dr. Kris Braman	University of Georgia Department of Entomology Griffin Campus Griffin, GA30223-1797
Dr. Juang-Horng Chong	Clemson University Department of Entomology, Soils & Plant Sciences Clemson, SC 29634
Dr. Raymond A Cloyd	Kansas State University Department of Entomology 239 W. Waters Hall Manhattan, KS 66506
Dr. Scott Costa	University of Vermont Dept. of Plant & Soil Sci. 206 Hills Bldg. Burlington, VT
Dr. Rich Cowles	Connecticut Agricultural Experiment Station 153 Cook Hill Road P.O. Box 248 Windsor, CT06095
Dr. Adam Dale	University of Florida Entomology & Nematology Department P. O. Box 110620 Gainesville, FL 32611-0620
Mr. Terry Davis (retired)	Michigan State University Department of Entomology East Lansing, MI 48824-1115

Dr. Steve Frank	North Carolina State University Department of Entomology Raleigh, NC 27695
Mr. Tom Freiburger (<i>past affiliate</i>)	Rutgers University Tree and Small Fruit Experiment Station 283 Rt. 539 Cream Ridge, NJ 08514
Mr. Dan Gilrein	Cornell Cooperative Extension Long Island Horticulture Research & Experiment Station 3059 Sound Avenue Riverhead, NY 11901
Dr. Charles Hesselein (<i>retired</i>)	Alabama Cooperative Extension Mobile Ornamental Horticulture Research Center P.O. Box 8276 Mobile, AL 36689
Mr. Grant Jones (<i>past affiliate</i>)	Davey Tree 1500 N. Mantua St. Kent, OH 44240
Dr. Brian Kunkel	University of Delaware 248A Townsend Hall Newark, DE 19716
Dr. Scott Ludwig (<i>past affiliate</i>)	Texas AgriLife Cooperative Extension P.O. Box 38 Overton, TX 75684
Dr. Russ Mizell	University of Florida AREC, Route 4, Box 4092 Monticello, FL 32344-9304
Dr. Dave Nielsen (<i>retired</i>)	Ohio State University, OARDC Department of Entomology Wooster, OH 44691
Dr. Jorge Pena	University of Florida Tropical Research and Education Center 18905 S. W. 280 St. Homestead, FL 33031
Dr. Anand Persad (<i>past affiliate</i>)	The Davey Institute 1500 N, Mantua St. Kent, OH 44240
Dr. Dan Potter	University of Kentucky Department of Entomology S-225 Agriculture Science Building North Lexington, KY 40546-0091

Dr. Michael Reding	USDA-ARS Hort Insects Lab 1680 Madison Ave. Wooster, OH, 44691
Dr. Cliff Sadof	Purdue University Department of Entomology 901 West State Street West Lafayette, IN 47907
Dr. Pete Schultz	Virginia Technical Institute Hampton Roads AREC 1444 Diamond Springs Road Virginia Beach, VA 23455
Dr. Shimat Villanassery Joseph	University of Georgia 285 Higgins Road Griffin, GA 30223
Dr. Paul Weston (<i>past affiliate</i>)	Cornell University Department of Entomology Ithaca, NY 14853