



[Environment Horticulture Program Research Summaries](#)

## **IR-4 Environmental Horticulture Program S-Metolachlor Crop Safety**

**Author: Cristi L. Palmer  
Date: September 13, 2022**

### **Acknowledgements**

**Ely Vea  
Susan Bierbrunner  
Nikoya Lightbourne**

This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award numbers 2015-34383-23710, 2017-34383-27100, 2019-34383-29973, 2020-34383-32455 and 2021-343830-34848 with substantial cooperation and support from the State Agricultural Experiment Stations and USDA-ARS.

## Table of Contents

Table of Contents .....	2
Table of Tables .....	3
Abstract .....	4
Introduction.....	5
Materials and Methods.....	5
Results and Summary .....	5
Phytotoxicity .....	5
Label Suggestions .....	19
Appendix 1: Contributing Researchers .....	20

## Table of Tables

Table 1.	List of Pennant Magnum treated crops with no or minimal transitory injury. ....	6
Table 3.	List of Pennant Magnum treated crops with no or minimal transitory injury seen at the 1X rate, but the 2X or 4X rate did cause significant phytotoxicity .....	6
Table 4.	List of Pennant Magnum treated crops exhibiting significant injury. ....	6
Table 2.	List of Pennant Magnum treated crops where more research is needed to clarify response.....	7
Table 2.	List of Pennant Magnum treated crops with insufficient trials to determine response	7
Table 5.	Detailed Summary of 2005 Crop Safety Testing with Pennant Magnum .....	8

## **Abstract**

From 2004 to 2021, IR-4 completed 196 trials on Pennant Magnum (s-metolachlor). The data contained in this report was generated to register uses of s-metolachlor on and around environmental horticulture plants with over-the-top applications. The s-metolachlor rates in the testing program were 2.5, 5.0 and 10.0 pounds active ingredient per acre (lb ai per A) as the 1X, 2X and 4X rates with 4, 6, or 8 week intervals between applications. Pennant Magnus has been applied to 80 plant genera or species. Of these, 11 plant species exhibited no or minimal transient injury after application at all three rates. Ten (10) crops exhibited no phytotoxicity at 2.5 lb ai per acre but did have some injury at 5.0 and/or 10.0 lb ai per acre. Twenty-two (22) crops exhibited significant phytotoxicity at even the lowest rate. For nine crops, the response among sites was variable, and 30 crops have less than three trials completed.

## Introduction

Control of broadleaved weeds and sedges in the production of woody and herbaceous perennials can be problematic because nurseries grow many different types of plants and not all genera or species are listed on labels. These weeds can also be difficult to control in landscape settings for the same reason. Since 2004, Pennant Magnum (s-metolachlor) has been screened for crop safety on 80 plant genera or species. The early research activities led to an expansion of the label in 2010.

## Materials and Methods

Prior to 2007, two applications were made approximately 30 days apart. In the 2007 protocol, two applications of Pennant Magnum were made approximately 8 weeks apart. Beginning in 2008, two applications of Pennant Magnum were made approximately 6 weeks apart. The application rates were 2.5, 5, and 10 lb ai per acre, plus a water treated control. A minimum of four plants (replicate treatments) were required with many researchers exceeding this minimum. Phytotoxicity was recorded on a scale of 0 to 10 (0 = no phytotoxicity; 10 = complete kill) at 1, 2, 4, 8, and 12 weeks after initial application. Some researchers also included readings 3 to 4 days after the initial and second applications. The following protocols were used: 04-001, 05-001, 06-009, 07-010, 18-020, 19-020, and 22-017. Please visit <https://www.ir4project.org/ehc/ehc-registration-support-research/env-hort-researcher-resources/#Protocols> to view and download these protocols.

Pennant Magnum was supplied to researchers (See list of researchers in Appendix 2) by Syngenta.

## Results and Summary

### *Phytotoxicity*

Based on the type and nature of injury seen with Pennant Magnum applications in the research conducted from 2004 to 2022, tested plant species were placed into four categories: 1) no significant Phytotoxicity or growth differences from the untreated check or any injury was transitory, 2) injury was seen but additional research is warranted to clarify response, 3) no or minimal transitory injury seen at the 1X rate, but the 2X and/or 4X rates did cause significant phytotoxicity, 4) Significant injury sufficient to recommend growers not utilize this product.

In general, Pennant Magnum exhibited no or minimal negative impact on a range of plant species (Table 1). Eleven plant genera or species fell into this category. Some minimal injury may be acceptable for growers if applications are made several weeks to months in advance of crop sale particularly for woody ornamental crops. Ten crop species, exhibited no or little injury at the 2 lb ai per acre rate, but significant phytotoxicity occurred at the 4 lb ai per acre rate (Table 2). It may be prudent to either conduct additional trials or place language on the label indicating applications of Pennant Magnum are considered safe at the 2 lb ai per acre rate but any higher rate may cause unacceptable injury.

There were twenty-two crops that exhibited damage sufficient to recommend growers not utilize Pennant Magnum as an over-the-top treatment for pre-emergent weed control (Table 3).

There were nine species of plants for which the results were unclear and are suggested as candidates for further research: *Asarum sp.*, *Aubrieta sp.*, *Bergenia sp.*, *Eupatorium maculatum*, *Eupatorium rugosum*, *Eupatorium purpureum*, *Panicum virgatum*, *Polenium sp.* and *Sempervivium sp.* (Table 4). There were 30 plant species or genera where IR-4 has conducted less than three trials.

Please see Table 6 for a list of individual trial summaries for Pennant Magnum.

**Table 1. List of Pennant Magnum treated crops with no or minimal transitory injury.**

<i>Iberis sempervirens</i> <sup>1</sup>	<i>Penstemon digitalis</i>
<i>Kniphofia uvaria</i> <sup>1</sup>	<i>Ruellia carolinensis</i> <sup>1</sup>
<i>Lavandula angustifolia</i> <sup>1</sup>	<i>Sempervivum tectorum</i> <sup>1</sup>
<i>Ligularia stenocephala</i>	<i>Solidago rugosa</i>
<i>Linum perenne</i>	<i>Solidago sempervirens</i> <sup>1</sup>
<i>Nepeta faassenii</i>	

**Table 2. List of Pennant Magnum treated crops with no or minimal transitory injury seen at the 1X rate, but the 2X or 4X rate did cause significant phytotoxicity**

<i>Baptisia australis</i>	<i>Helianthus salicifolius</i>
<i>Cimicifuga racemosa</i>	<i>Helleborus niger</i>
<i>Gazania sp.</i>	<i>Hibiscus syriacus</i> <sup>1</sup>
<i>Gomphrena sp.</i>	<i>Santolina chamaecyparissus</i>
<i>Helenium autumnale</i>	<i>Verbena canadensis</i>

**Table 3. List of Pennant Magnum treated crops exhibiting significant injury.**

<i>Agastache aurantiaca</i>	<i>Echinacea purpurea</i>
<i>Agastache rugosa x foeniculum</i>	<i>Epimedium x rubrum</i>
<i>Alchemilla mollis</i>	<i>Gerbera jamesonii</i>
<i>Amsonia hubrichtii</i>	<i>Heuchera sanguinea</i>
<i>Amsonia tabernaemontana</i>	<i>Nepeta nervosa</i>
<i>Armeria maritima</i>	<i>Origanum libanoticum</i>
<i>Bergenia cordifolia</i>	<i>Pentas sp.</i>
<i>Chelone lyonii</i>	<i>Primula malacoides</i>
<i>Chrysogonum virginianum</i>	<i>Tiarella cordifolia</i>
<i>Delosperma nubigenum</i>	<i>Tradescantia andersoniana</i>
<i>Digitalis thapsi</i>	<i>Tradescantia ohiensis</i>

**Table 4. List of Pennant Magnum treated crops where more research is needed to clarify response**

<i>Asarum candanense</i>	<i>Eupatorium purpureum</i>
<i>Aubrieta sp.</i>	<i>Panicum virgatum</i>
<i>Bergenia sp.</i>	<i>Polemonium sp.</i>
<i>Eupatorium maculatum</i>	<i>Sempervivum sp.</i>
<i>Eupatorium rugosum</i>	

**Table 5. List of Pennant Magnum treated crops with insufficient trials to determine response**

<i>Actaea racemosa var. racemosa</i>	<i>Liriope muscari</i> <sup>1</sup>
<i>Agastache rupestris</i>	<i>Opuntia humifusa</i> <sup>1</sup>
<i>Antennaria parvifolia</i>	<i>Penstemon x mexicali</i>
<i>Asarum chinensis</i>	<i>Phlox paniculata</i>
<i>Athyrium nipponicum</i>	<i>Phlox subulata</i>
<i>Clematis sp.</i>	<i>Phormium colinsoi</i> <sup>1</sup>
<i>Cosmos sp.</i>	<i>Pulmonaria sp.</i>
<i>Cuphea hyssopifolia</i>	<i>Rudbeckia hirta</i>
<i>Helianthemum nummularium</i>	<i>Ruellia brittoniana</i>
<i>Heliopsis helianthoides</i>	<i>Santolina sp.</i>
<i>Helleborus sp.</i>	<i>Scabiosa sp.</i>
<i>Hibiscus mosheutos</i> <sup>1</sup>	<i>Solidago speciosa</i>
<i>Lantana hybrida</i> <sup>1</sup>	<i>Symphotrichum ericoides</i>
<i>Lantana montevidensi</i> <sup>1</sup>	<i>Vernonia noveboracensis</i> <sup>1</sup>
<i>Leucanthemum x superbum</i>	<i>Zinnia sp.</i>

<sup>1</sup> Registered already on EPA label.

**Table 6. Detailed Summary of 2005 Crop Safety Testing with Pennant Magnum**

Note: Table entries are sorted by crop Latin name. Only those trials with research reports received by 9/9/2022 are listed below.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
23828	Black Bugbane ( <i>Actaea racemosa</i> var. <i>racemosa</i> )	Field Container	Neal	NC	2005	Over the top	Little to no injury at 2.5 lb ai per acre; moderate to severe at 5 lb ai per acre and 10 lb ai per acre, respectively
24724	Giant Hyssop, Sunset ( <i>Agastache rupestris</i> ) 'Mango Tango'	Field Container	Senesac	NY	2022	Over the top	Minor to severe injury increasing with rate (2.5, 5.0 10.0 lb ai per acre).
24723	Giant Hyssop ( <i>Agastache</i> sp.) <i>A. rugosa</i> X <i>foeniculum</i> 'Blue Fortune'	Field Container	Boydston	WA	2005	Over the top	Minor to severe injury increasing with rate (2.5, 5.0, 10 lb ai per acre).
24723	Giant Hyssop ( <i>Agastache</i> sp.) <i>A. aurantiaca</i> 'Coronado'	Field Container	Klett	CO	2005	Over the top	Fort Collins: Minor to severe injury increasing with rate (2.5, 5.0, 10 lb ai per acre).
24723	Giant Hyssop ( <i>Agastache</i> sp.) <i>A. aurantiaca</i> 'Coronado'	Field Container	Klett	CO	2005	Over the top	Grand Junction: Moderate injury at all rates (2.5, 5.0, 10 lb ai per acre).
24723	Giant Hyssop ( <i>Agastache</i> sp.) <i>A. rugosa</i> X <i>foeniculum</i> 'Blue Fortune'	Field Container	Mathers (OSU)	OH	2005	Over the top	All rates (2.5, 5.0. 10.0 lb ai per acre) caused slight to moderate injury (leaf distortion and burning)
23819	Lady's-Mantle ( <i>Alchemilla</i> sp.) <i>A. mollis</i> 'Thriller'	Field Container	Ahrens/Mervosh	CT	2005	Over the top	Slight injury at 2.5 lb ai per acre and 5 lb ai per acre rates, moderate injury at 10 lb ai per acre
23819	Lady's-Mantle ( <i>Alchemilla</i> sp.) <i>A. mollis</i> 'Selection 25'	Field Container	Boydston	WA	2005	Over the top	No injury with 2.5, 5.0 and 10 lb ai per acre.
23819	Lady's-Mantle ( <i>Alchemilla</i> sp.) <i>A. mollis</i>	Field Container	Derr	VA	2004	Over the top	Slight injury at 2.5 lb ai per acre rate, moderate injury at 5 lb ai per acre and 10 lb ai per acre
23819	Lady's-Mantle ( <i>Alchemilla</i> sp.) <i>A. mollis</i>	Field Container	Lieth	CA	2004	Over the top	All rates caused severe leaf necrosis
23819	Lady's-Mantle ( <i>Alchemilla</i> sp.)	Field Container	Reding	OH	2005	Over the top	Slight injury at 2.5 lb ai per acre rate; slight to moderate injury at 5 lb ai per acre and 10 lb ai per acre
23819	Lady's-Mantle ( <i>Alchemilla</i> sp.) <i>A. mollis</i>	Field Container	Senesac	NY	2004	Over the top	No injury at 2.5 lb ai per acre and 5 lb ai per acre rates, slight injury at 10 lb ai per acre
34865	Hubricht's Bluestar, Arkansas Amsonia ( <i>Amsonia hubrichtii</i> )	Field Container	Larson	WA	2022	Over the top	Moderate leaf burning and no flowers with 10 lb ai per acre.
34865	Hubricht's Bluestar, Arkansas Amsonia ( <i>Amsonia hubrichtii</i> ) 'Butterscotch'	Field Container	Senesac	NY	2022	Over the top	Moderate to severe injury increasing with rate (2.5, 5.0 10.0 lb ai per acre).
23820	Eastern Bluestar ( <i>Amsonia tabernaemontana</i> ) <i>A. tabernaemontana</i>	Field Container	Gilliam	AL	2004	Over the top	All rates caused serious leaf burn (2.5, 5.0, 10.0 lb ai per acre).
23820	Eastern Bluestar ( <i>Amsonia tabernaemontana</i> ) <i>A. hubrichtii</i>	Field Container	Neal	NC	2004	Over the top	All rates caused significant injury (2.5, 5.0, 10.0 lb ai per acre).
23820	Eastern Bluestar ( <i>Amsonia tabernaemontana</i> ) <i>A. hubrichtii</i>	Field Container	Senesac	NY	2004	Over the top	Slight injury at 2.5 lb ai per acre rate and moderate injury at 5 lb ai per acre and 10 lb ai per acre rates.



PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
23822	Pussy-Toes, Small-leaf ( <i>Antennaria parvifolia</i> )	Field Container	Neal	NC	2004	Over the top	All rates caused moderate to severe injury (2.5, 5.0, 10.0 lb ai per acre).
23822	Pussy-Toes, Small-leaf ( <i>Antennaria parvifolia</i> )	Field Container	Senesac	NY	2004	Over the top	No injury with 2.5, 5.0, 10.0 lb ai per acre.
24729	Thrift Sea Pink ( <i>Armeria maritima</i> ) 'Splendens'	Field Container	Boydston	WA	2005	Over the top	Moderate to severe injury increasing with rate (2.5, 5.0 and 10 lb ai per acre). Single application caused purpling of leaves; second application caused additional injury.
24729	Thrift Sea Pink ( <i>Armeria maritima</i> ) 'Splendens'	Field Container	Gilliam	AL	2005	Over the top	All rates caused moderate to severe injury (burn and discoloration) and stunting (2.5, 5, and 10 lb ai per acre).
24729	Thrift Sea Pink ( <i>Armeria maritima</i> ) 'Dusseldorf'	Field Container	Lieth	CA	2005	Over the top	All rates caused unacceptable growth suppression
23823	Canadian Wildginger ( <i>Asarum canadense</i> ) <i>A. chinensis</i>	Field Container	Neal	NC	2005	Over the top	No injury at all rates (2.5, 5.0, 10.0 lb ai per acre).
23823	Canadian Wildginger ( <i>Asarum canadense</i> )	Field Container	Senesac	NY	2005	Over the top	Moderate to severe injury increasing with rate (2.5, 5.0, 10.0 lb ai per acre).
31435	Canadian Wildginger ( <i>Asarum canadense</i> )	Field Container	Senesac	NY	2022	Over the top	Severe injury with 2.5, 5.0 and 10.0 lb ai per acre.
24738	Fern, Lady ( <i>Athyrium nipponicum</i> )	Field Container	Derr	VA	2005	Over the top	No injury at 2.5 lb ai per acre and 5 lb ai per acre rate, slight injury at 10 lb ai per acre
24738	Fern, Lady ( <i>Athyrium nipponicum</i> ) 'Pretum'	Field Container	Mathers (OSU)	OH	2005	Over the top	Slight injury (frond scorching) at 2.5 lb ai per acre and 5 lb ai per acre rates; moderate injury from 10 lb ai per acre
26018	Lilacbush ( <i>Aubrieta</i> sp.) 'Large Flo'	Greenhouse	Freiberger	NJ	2006	Over the top	Light to high injury increasing with rates (2.5, 5 and 10 lb ai per acre)
23824	Lilacbush ( <i>Aubrieta</i> sp.) 'Whitewell Gem'	Field Container	Lieth	CA	2004	Over the top	No injury but reduced plant width
26018	Lilacbush ( <i>Aubrieta</i> sp.) 'Whitewell Gem'	Greenhouse	Mickelbart	IN	2006	Over the top	No significant injury or growth reduction at 2.5 lb ai per acre; significant injury at 5 and 10 lb ai
23824	Lilacbush ( <i>Aubrieta</i> sp.) 'Whitewell Gem'	Field Container	Neal	NC	2004	Over the top	All rates caused significant injury (2.5, 5.0, 10.0 lb ai per acre).
23824	Lilacbush ( <i>Aubrieta</i> sp.)	Field Container	Reding	OH	2005	Over the top	No injury
23825	Blue Wild Indigo ( <i>Baptisia australis</i> )	Field Container	Derr	VA	2004	Over the top	All rates caused slight transient injury (2.5. 5.0. 10.0 lb ai per acre). Excellent efficacy for large crabgrass and spotted spurge
23825	Blue Wild Indigo ( <i>Baptisia australis</i> )	Field Container	Gilliam	AL	2004	Over the top	No injury at 2.5 lb ai per acre and 5 lb ai per acre rates, slight injury at 10 lb ai per acre
23825	Blue Wild Indigo ( <i>Baptisia australis</i> )	Field Container	Neal	NC	2004	Over the top	No injury at 2.5 lb ai per acre rate, acceptable at 5 lb ai per acre and unacceptable at 10 lb ai per acre

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
23825	Blue Wild Indigo ( <i>Baptisia australis</i> )	Field Container	Senesac	NY	2005	Over the top	No injury at all rates (2.5, 5.0, 10.0 lb ai per acre).
24739	Siberian Tea ( <i>Bergenia cordifolia</i> ) 'Rotblum'	Field Container	Lieth	CA	2005	Over the top	Significant injury and unacceptable growth suppression
26021	Siberian Tea ( <i>Bergenia cordifolia</i> ) 'Winter Glow'	Greenhouse	Mickelbart	IN	2006	Over the top	No significant injury or growth reduction (2.5, 5 and 10 lb ai per acre)
24739	Siberian Tea ( <i>Bergenia cordifolia</i> ) 'New Hybrid'	Field Container	Neal	NC	2005	Over the top	Slight to moderate injury at 2.5 lb ai per acre rate. Severe at 5 lb ai per acre and 10 lb ai per acre
23826	Turtlehead, Snakehead ( <i>Chelone</i> sp.) <i>C. lyonii</i> 'Hot Lips'	Field Container	Derr	VA	2004	Over the top	Slight injury at 2.5 lb ai per acre rate, moderate at 5 lb ai per acre severe at 10 lb ai per acre
23826	Turtlehead, Snakehead ( <i>Chelone</i> sp.) <i>C. lyonii</i> 'Hot Lips'	Field Container	Neal	NC	2004	Over the top	Slight injury at 2.5 lb ai per acre rate, severe injury at 5 lb ai per acre and 10 lb ai per acre rates
23826	Turtlehead, Snakehead ( <i>Chelone</i> sp.) <i>C. lyonii</i> 'Hot Lips'	Field Container	Senesac	NY	2004	Over the top	All rates caused moderate injury (2.5. 5.0. 10.0 lb ai per acre).
23827	Golden Star ( <i>Chrysogonum virginianum</i> ) <i>C. virginianum</i> 'Alan Bush'	Field Container	Derr	VA	2004	Over the top	Moderate injury increasing with rate Excellent efficacy for large crabgrass and spotted spurge
23827	Golden Star ( <i>Chrysogonum virginianum</i> ) <i>C. virginianum</i> 'Alan Bush'	Field Container	Neal	NC	2004	Over the top	Slight injury at 2.5 lb ai per acre rate, moderate at 5 lb ai per acre and severe at 10 lb ai per acre
23827	Golden Star ( <i>Chrysogonum virginianum</i> ) <i>C. virginianum</i> 'Alan Bush'	Field Container	Senesac	NY	2005	Over the top	No significant injury at 2.5 lb ai per acre rate; moderate to severe injury at 5 lb ai per acre and 10 lb ai per acre.
24830	Leather Flower ( <i>Clematis</i> sp.) 'Midnight Showers'	Field Container	Mathers (OSU)	OH	2005	Over the top	All rates (2.5, 5.0. 10.0 lb ai per acre) caused slight injury but plants recovered
13129	Cosmos ( <i>Cosmos</i> sp.)	Field In-Ground	Miller	WA	2018	Over the top	Minor injury with 2.6, 5.2 and 10.4 pt per acre applied twice; moderate growth reduction with 4X.
13129	Cosmos ( <i>Cosmos</i> sp.)	Field In-Ground	Sellmer	PA	2001	Over the top	
25296	False Heather, Elfin Herb, Mexican Heather ( <i>Cuphea hyssopifolia</i> )	Field In-Ground	Chen	LA	2005	Over the top	Moderate injury with 2.5, 5.0 and 10 lb ai per acre; plants recovered in a short time
24832	Hardy Ice Plant, Yellow Ice Plant ( <i>Delosperma nubigenum</i> )	Field Container	Boydston	WA	2005	Over the top	Slight injury with 2.5 and 5.0 lb ai per acre; severe injury with 10 lb ai per acre. All rates caused slight stunting and delayed flowering.
24832	Hardy Ice Plant, Yellow Ice Plant ( <i>Delosperma nubigenum</i> ) D. Hardy Ice Plant yellow	Field Container	Boydston	WA	2006	Over the top	Significant injury and growth reduction at all rates (2.5, 5 and 10 lb ai per acre); marketability questionable
24832	Hardy Ice Plant, Yellow Ice Plant ( <i>Delosperma nubigenum</i> )	Field Container	Linderman	OR	2006	Over the top	No injury at 2.5, 5 and 10 lb ai per acre
24741	Foxglove ( <i>Digitalis</i> sp.) <i>D. purpurea</i> 'Foxy'	Field Container	Boydston	WA	2006	Over the top	No injury or growth reduction at 1X rate (2.5, lb ai per acre) only; 2X and 4X plants not saleable
26024	Foxglove ( <i>Digitalis</i> sp.) 'Excelsior Mix'	Greenhouse	Freiberger	NJ	2006	Over the top	No injury after 1st application at all rates (2.5, 5 and 10 lb ai per acre), moderate and high injury at 2X and 4X after 2nd application

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
24741	Foxglove ( <i>Digitalis</i> sp.) <i>D. thaspi</i> 'Spanish Peaks'	Field Container	Klett	CO	2005	Over the top	Fort Collins: Moderate to severe injury increasing with rate (2.5, 5.0, 10 lb ai per acre).
24741	Foxglove ( <i>Digitalis</i> sp.) <i>D. thaspi</i> 'Spanish Peaks'	Field Container	Klett	CO	2005	Over the top	Grand Junction: No to severe injury increasing with rate (2.5, 5.0, 10 lb ai per acre).
24741	Foxglove ( <i>Digitalis</i> sp.) <i>D. purpurea</i>	Field Container	Linderman	OR	2006	Over the top	No injury at 2.5, 5 and 10 lb ai per acre
26024	Foxglove ( <i>Digitalis</i> sp.) 'Pam's Choice'	Greenhouse	Mickelbart	IN	2006	Over the top	No significant injury or growth reduction (2.5, 5 and 10 lb ai per acre)
24741	Foxglove ( <i>Digitalis</i> sp.) <i>D. ambigua</i>	Field Container	Treat	WA	2006	Over the top	Severe injury (leaf burn) and stunting at all rates (2.5, 5 and 10 lb ai per acre)
33361	Eastern Purple Coneflower ( <i>Echinacea purpurea</i> ) 'Magnus'	Field In-Ground	Aulakh	CT	2018	Over the top	No injury or growth reduction with 2.6, 5.2 and 10.4 pt per acre after 1st applic, moderate after the 2nd application with 2X and 4X.
33361	Eastern Purple Coneflower ( <i>Echinacea purpurea</i> ) 'Cheyenne Spirit'	Field In-Ground	Derr	VA	2021	Over the top	No injury with 2 pint per acre.
24742	Purple Coneflower ( <i>Echinacea</i> sp.) <i>E. purpurea</i> 'White Swan'	Field Container	Boydston	WA	2006	Over the top	Significant injury at all rates (2.5, 5 and 10 lb ai per acre); 4X plants smaller
24742	Purple Coneflower ( <i>Echinacea</i> sp.) <i>E. purpurea</i> 'Magnus'	Field Container	Derr	VA	2005	Over the top	Very slight injury at 2.5 lb ai per acre and 5 lb ai per acre rates, slight injury at 10 lb ai per acre rate.
24742	Purple Coneflower ( <i>Echinacea</i> sp.) <i>E. purpurea</i>	Field Container	Fraelich	GA	2006	Over the top	Slight injury; no differences in growth or marketability
26027	Purple Coneflower ( <i>Echinacea</i> sp.) 'Magnus'	Greenhouse	Freiberger	NJ	2006	Over the top	No injury at all rates (2.5, 5 and 10 lb ai per acre)
24742	Purple Coneflower ( <i>Echinacea</i> sp.) <i>E. purpurea</i> 'Magnus'	Field Container	Gilliam	AL	2005	Over the top	All rates caused moderate to severe injury (leaf curl and leaf distortion) and stunting (2.5, 5, and 10 lb ai per acre).
24742	Purple Coneflower ( <i>Echinacea</i> sp.) <i>E. purpurea</i> 'Magnus'	Field Container	Linderman	OR	2006	Over the top	No injury at 2.5, 5 and 10 lb ai per acre
26027	Purple Coneflower ( <i>Echinacea</i> sp.) 'White Swan'	Greenhouse	Mickelbart	IN	2006	Over the top	Slight injury and growth reduction (2.5, 5 and 10 lb ai per acre)
23829	Epimedium ( <i>Epimedium</i> sp.) <i>E. X rubrum</i>	Field Container	Ahrens/Mervosh	CT	2005	Over the top	Slight injury at 2.5 lb ai per acre and 5 lb ai per acre rates; moderate to severe injury at 10 lb ai per acre
23829	Epimedium ( <i>Epimedium</i> sp.) <i>E. X rubrum</i>	Field Container	Senesac	NY	2005	Over the top	Moderate injury at all rates (2.5, 5.0, 10.0 lb ai per acre).
23830	Joe Pye Weed ( <i>Eupatorium maculatum</i> ) 'Gateway'	Field Container	Ahrens/Mervosh	CT	2005	Over the top	No injury at 2.5 lb ai per acre and 5 lb ai per acre rates; slight injury at 10 lb ai per acre
23831	Common Boneset ( <i>Eupatorium perfoliatum</i> ) <i>E. rugosum</i> 'Chocolate'	Field Container	Boydston	WA	2006	Over the top	No to moderate injury increasing with rate; slight growth reduction at all rates (2.5, 5 and 10 lb ai per acre); all plants saleable
23832	Joe Pye Weed ( <i>Eupatorium purpureum</i> ) <i>E. dubium</i> 'Little Joe'	Field Container	Boydston	WA	2006	Over the top	No injury or growth reduction at 1Xrate (2.5 lb ai per acre) only; 1X and 2X plants saleable

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
23832	Joe Pye Weed ( <i>Eupatorium purpureum</i> ) 'Gateway'	Field Container	Neal	NC	2005	Over the top	No injury at 2.5 lb ai per acre rate, moderate at 5 lb ai per acre and severe at 10 lb ai per acre
23832	Joe Pye Weed ( <i>Eupatorium purpureum</i> )	Field Container	Senesac	NY	2004	Over the top	Slight to moderate injury increasing with rate (2.5, 5.0, 10.0 lb ai per acre).
23832	Joe Pye Weed ( <i>Eupatorium purpureum</i> ) E. 'Sweet Scented'	Field Container	Treat	WA	2006	Over the top	Slight to severe injury increasing with rate (2.5, 5.0, 10.0 lb ai per acre).
24743	Thoroughwort ( <i>Eupatorium</i> sp.) E. rugosum 'Chocolate'	Field Container	Boydston	WA	2005	Over the top	No injury with 2.5, 5.0 and 10 lb ai per acre.
31434	Thoroughwort ( <i>Eupatorium</i> sp.)	Field Container	Larson	WA	2022	Over the top	No injury with 2.5, 5, and 10 lb ai per acre.
31434	Thoroughwort ( <i>Eupatorium</i> sp.) E. dubium 'Baby Joe'	Field Container	Senesac	NY	2022	Over the top	Minor to moderate injury increasing with rate (2.5, 5.0 10.0 lb ai per acre).
24743	Thoroughwort ( <i>Eupatorium</i> sp.) E. colestinum	Field Container	Treat	WA	2006	Over the top	Minor to severe injury increasing with rate (2.5, 5.0, 10.0 lb ai per acre) with recovery by end of the experiment at 2.5 lb ai per acre, but some mortality did occur with the 10.0 lb ai per acre rate.
24744	Treasureflower ( <i>Gazania linearis</i> ) 'Colorado Gold'	Field Container	Neal	NC	2005	Over the top	No injury at 2.5 lb ai per acre rate, moderate injury at 5 lb ai per acre severe at 10 lb ai per acre
24744	Treasureflower ( <i>Gazania linearis</i> ) 'Colorado Gold'	Field Container	Treat	WA	2006	Over the top	Significant injury with complete recovery by trial's end at 1X rate (2.5 lb ai per acre); unacceptable at higher rates
26831	<i>Gazania</i> ( <i>Gazania</i> sp.) 'Daybreak Mix'	Greenhouse	Freiberger	NJ	2006	Over the top	No injury at 2.5 lb ai per acre, slight and moderate at 5 and 10 lb ai
24746	Transvaal Daisy ( <i>Gerbera</i> sp.) <i>G. jamesonii</i>	Field Container	Chandran	WV	2005	Over the top	No injury at 2.5, 5 and 10 lb ai per acre
24746	Transvaal Daisy ( <i>Gerbera</i> sp.) <i>Gerbera jamesonii</i>	Field Container	Fraulich	GA	2006	Over the top	Moderate injury at higher rates but plants recovered; no differences in growth or marketability
24746	Transvaal Daisy ( <i>Gerbera</i> sp.) <i>G. jamesonii</i> 'Lambada'	Field Container	Lieth	CA	2005	Over the top	Significant injury and severe growth suppression
24746	Transvaal Daisy ( <i>Gerbera</i> sp.) <i>G. jamesonii</i>	Field Container	Linderman	OR	2006	Over the top	No injury at 2.5, 5 and 10 lb ai per acre
24746	Transvaal Daisy ( <i>Gerbera</i> sp.) 'Dark Eye Orange'	Field Container	Treat	WA	2006	Over the top	Severe injury (leaf burn and stunting) increasing with rates (2.5, 5 and 10 lb ai per acre)
24747	Globe Amaranth ( <i>Gomphrena</i> sp.)	Field Container	Chandran	WV	2005	Over the top	No significant injury at 2.5, 5 and 10 lb ai per acre
24747	Globe Amaranth ( <i>Gomphrena</i> sp.) <i>G. uvaria</i>	Field Container	Gilliam	AL	2005	Directed	Slight injury (leaf burn and discoloration) at 2.5 lb ai per acre and 5 lb ai per acre rates, moderate at 10 lb ai per acre
24747	Globe Amaranth ( <i>Gomphrena</i> sp.) <i>G. globosa</i> 'Buddy Series Purple'	Field Container	Treat	WA	2006	Over the top	Slight to moderate injury with complete recovery by trial's end at 1X rate (2.5 lb ai per acre); unacceptable at higher rates

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
23833	Common Sneezeweed ( <i>Helenium autumnale</i> )	Field Container	Lieth	CA	2004	Over the top	No injury at 2.5 lb ai per acre and 5 lb ai per acre rates slight injury at 10 lb ai per acre
23833	Common Sneezeweed ( <i>Helenium autumnale</i> ) 'Summer Sun'	Field Container	Neal	NC	2004	Over the top	No injury at 2.5 and 5 lb ai per acre; slight at 10 lb ai per acre, but plants recovered
23833	Common Sneezeweed ( <i>Helenium autumnale</i> )	Field Container	Reding	OH	2005	Over the top	No significant injury at 2.5 lb ai per acre and 5 lb ai per acre rates; plants outgrew slight to moderate injury at 10 lb ai per acre
23833	Common Sneezeweed ( <i>Helenium autumnale</i> )	Field Container	Senesac	NY	2005	Over the top	Minor injury after second application but plants recovered by end of experiment
26847	Sneezeweed ( <i>Helenium</i> sp.) 'Hoopes II Yellow'	Greenhouse	Freiberger	NJ	2006	Over the top	No injury at 2.5 lb ai per acre, slight at 5 and 10 lb ai
24748	Frostweed ( <i>Helianthemum</i> sp.) <i>H. nummularium</i> 'Belgravia Rose'	Field Container	Lieth	CA	2005	Over the top	No significant injury at 2.5 lb ai per acre and 5 lb ai per acre rates, all rates reduced growth.
23834	Sunflower, Willowleaf ( <i>Helianthus salicifolius</i> ) 'First Light'	Field Container	Mathers (OSU)	OH	2005	Over the top	No injury with 2.5, 5.0, 10.0 lb ai per acre.
23834	Sunflower, Willowleaf ( <i>Helianthus salicifolius</i> ) 'First Light'	Field Container	Senesac	NY	2005	Over the top	No significant injury at 2.5 lb ai per acre and 5 lb ai per acre rates, minor injury at 10 lb ai per acre
24749	Sunflower ( <i>Helianthus</i> sp.) <i>H. salicifolius</i> 'First light'	Field Container	Treat	WA	2006	Over the top	Minor to severe injury (leaf burn and stunting) increasing with rates (2.5, 5 and 10 lb ai per acre)
23836	Smooth Oxeye ( <i>Heliopsis helianthoides</i> ) 'Summer Sun'	Field Container	Boydston	WA	2005	Over the top	Slight injury with 2.5 and 5.0 lb ai per acre; moderate injury with 10 lb ai per acre. No significant height differences with 2.5 lb ai per acre, but significant stunting with 5.0 and 10.0 lb ai per acre.
26030	Smooth Oxeye ( <i>Heliopsis helianthoides</i> ) 'Summer Nights'	Greenhouse	Mickelbart	IN	2006	Over the top	Slight to high injury with increasing rates (2.5, 5 and 10 lb ai per acre); no growth reduction
23837	Hellebore, Black ( <i>Helleborus niger</i> ) <i>H. X orientalis</i> 'White Spotted Lady'	Field Container	Senesac	NY	2005	Over the top	Slight to moderate injury increasing with rate (2.5, 5.0, 10.0 lb ai per acre), but dissipating over time.
24750	Coral Bells ( <i>Heuchera sanguinea</i> )	Field Container	Boydston	WA	2005	Over the top	Moderate to severe injury increasing with rate (2.5, 5.0 and 10 lb ai per acre). Single application caused reddening and necrosis; second application caused additional injury.
24750	Coral Bells ( <i>Heuchera sanguinea</i> )	Field Container	Klett	CO	2005	Over the top	Fort Collins: Moderate to severe injury increasing with rate (2.5, 5.0, 10 lb ai per acre).
24750	Coral Bells ( <i>Heuchera sanguinea</i> )	Field Container	Klett	CO	2005	Over the top	Grand Junction: Severe injury at all rates (2.5, 5.0, 10 lb ai per acre).
24750	Coral Bells ( <i>Heuchera sanguinea</i> ) 'Firefly'	Field Container	Lieth	CA	2005	Over the top	Significant injury and growth suppression with increasing suppression at increased rates
25788	Rose-Of-Sharon ( <i>Hibiscus syriacus</i> ) <i>H. moscheutos</i> 'Splash Pinot Noir'	Field Container	Senesac	NY	2005	Over the top	Slight injury at 2.5 and 5, moderate at 10 lb ai per acre.
24833	Candytuft ( <i>Iberis</i> sp.) <i>I. sempervirens</i> 'Alexander's White'	Field Container	Boydston	WA	2006	Over the top	No injury or growth reduction at all rates (2.5, 5 and 10 lb ai per acre); all plants saleable
26840	Candytuft ( <i>Iberis</i> sp.) 'White'	Greenhouse	Freiberger	NJ	2006	Over the top	No injury at 2, 4 and 8 lb ai per acre.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
24833	Candytuft (Iberis sp.) I. sempervivens 'Snowflake'	Field Container	Lieth	CA	2005	Over the top	No significant injury; plant growth suppression
24833	Candytuft (Iberis sp.) I. sempervivens 'Purite'	Field Container	Treat	WA	2006	Over the top	No injury at 2.5 and 5 lb ai per acre; 10 lb ai per acre unacceptable
23838	Poker Plant (Kniphofia sp.) K. uvaria 'Flamenco'	Field Container	Boydston	WA	2005	Over the top	No injury with 2.5, 5.0 and 10.0 lb ai per acre.
23838	Poker Plant (Kniphofia sp.) K. uvaria 'Flamenco mix'	Field Container	Derr	VA	2005	Over the top	No injury at all rates (2.5, 5.0, 10.0 lb ai per acre).
23838	Poker Plant (Kniphofia sp.) K. uvaria	Field Container	Gilliam	AL	2005	Directed	No injury with 2.5, 5, and 10 lb ai per acre.
23838	Poker Plant (Kniphofia sp.) K. uvaria 'Border Ballet'	Field Container	Lieth	CA	2004	Over the top	No injury
23838	Poker Plant (Kniphofia sp.) K. uvaria 'Pfitzers Hybrid'	Field Container	Neal	NC	2005	Over the top	No injury at all rates (2.5, 5.0, 10.0 lb ai per acre).
25298	Lantana (Lantana sp.) L. hybrida 'New Gold'	Field In-Ground	Chen	LA	2005	Over the top	Slight injury with 2.5, 5.0 and 10 lb ai per acre.
24688	Lantana (Lantana sp.) L. montevidensis	Field Container	Lieth	CA	2005	Over the top	No injury or plant growth suppression
23839	Lavender, English (Lavandula angustifolia)	Field Container	Gilliam	AL	2004	Over the top	No injury with 2.1, 4.2, and 8.4 lb ai per acre.
23839	Lavender, English (Lavandula angustifolia) 'Vera'	Field Container	Lieth	CA	2004	Over the top	No injury with 2.1, 4.2, and 8.4 lb ai per acre.
23839	Lavender, English (Lavandula angustifolia) 'Munstead'	Field Container	Neal	NC	2004	Over the top	No injury at 2.5 lb ai per acre, slight injury at 5 lb ai per acre rate and unacceptable at 10 lb ai per acre
23839	Lavender, English (Lavandula angustifolia) 'Munstead'	Field Container	Senesac	NY	2004	Over the top	No injury with 2.5, 5.0, and 10.0 lb ai per acre.
34874	Daisy, Shasta (Leucanthemum x superbum)	Field Container	Larson	WA	2022	Over the top	No injury with 2.5, 5, and 10 lb ai per acre.
34874	Daisy, Shasta (Leucanthemum x superbum) 'Madonna'	Field Container	Senesac	NY	2022	Over the top	Moderate to severe injury increasing with rate (2.5, 5.0 10.0 lb ai per acre).
23840	Golden Rockets (Ligularia stenocephala) 'Dark Leaf'	Field Container	Boydston	WA	2005	Over the top	No injury with 2.5, 5.0 and 10.0 lb ai per acre.
23840	Golden Rockets (Ligularia stenocephala)	Field Container	Fraelich	GA	2005	Over the top	No injury or growth differences at all rates
23840	Golden Rockets (Ligularia stenocephala) 'The Rocket'	Field Container	Senesac	NY	2005	Over the top	No significant injury at all rates (2.5, 5.0, 10.0 lb ai per acre).
13195	Sealavender (Limonium sp.)	Field In-Ground	Sellmer	PA	2001	Over the top	
23841	Perennial Flax (Linum perenne) 'Sapphire'	Field Container	Lieth	CA	2004	Over the top	No injury at 2.1, 2.4, or 8.4 lb ai per acre.
23841	Perennial Flax (Linum perenne) 'Sapphyr"	Field Container	Neal	NC	2004	Over the top	No injury at 2.5 lb ai per acre and 5 lb ai per acre rates, slight injury at 10 lb ai per acre.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
23841	Perennial Flax ( <i>Linum perenne</i> )	Field Container	Senesac	NY	2004	Over the top	No injury at 2.5 lb ai per acre and 5 lb ai per acre rates, slight injury at 10 lb ai per acre.
24985	Lilyturf, Big Blue ( <i>Liriope muscari</i> ) 'Big Blue'	Field In-Ground	Chen	LA	2005	Over the top	No significant injury with 2.5, 5.0 and 10 lb ai per acre
23843	Nepeta ( <i>Nepeta x faassenii</i> ) 'Walker's Low'	Field Container	Derr	VA	2004	Over the top	All rates caused slight injury (2.5. 5.0. 10.0 lb ai per acre).
23843	Nepeta ( <i>Nepeta x faassenii</i> ) 'Dropmore'	Field Container	Gilliam	AL	2004	Over the top	No injury with 2.1, 2.4, and 8.4 lb ai per acre.
23843	Nepeta ( <i>Nepeta x faassenii</i> ) 'Dropmore'	Field Container	Lieth	CA	2004	Over the top	No injury with 2.1, 2.4, and 8.4 lb ai per acre.
23843	Nepeta ( <i>Nepeta x faassenii</i> ) N. nervosa 'Blue Carpet'	Field Container	Neal	NC	2004	Over the top	Severe injury at all rates (2..5, 5.0, 10.0 lb ai per acre)
23843	Nepeta ( <i>Nepeta x faassenii</i> ) 'Walker's Low'	Field Container	Senesac	NY	2004	Over the top	No injury with 2.5, 5.0 and 10.0 lb ai per acre.
23844	Devil's-tongue; Eastern Prickly Pear ( <i>Opuntia humifusa</i> ) 'Lemon Form'	Field Container	Senesac	NY	2005	Over the top	No injury with 2.5, 5.0, 10.0 lb ai per acre.
24754	Hopflower Oregano; Majoram ( <i>Origanum libanoticum</i> )	Field Container	Klett	CO	2005	Over the top	Fort Collins: Moderate to severe injury increasing with rate (2.5, 5.0, 10 lb ai per acre).
24754	Hopflower Oregano; Majoram ( <i>Origanum libanoticum</i> )	Field Container	Klett	CO	2005	Over the top	Grand Junction: Severe injury at all rates (2.5, 5.0, 10 lb ai per acre).
24834	Switch Grass ( <i>Panicum virgatum</i> ) P. virgatum 'Trailblazer'	Field Container	Boydston	WA	2006	Over the top	No injury or growth reduction at 1X and 2X rates (2.5 and 5 lb ai per acre), 4x rate plants slightly smaller; all plants saleable
26833	Switch Grass ( <i>Panicum virgatum</i> )	Greenhouse	Freiberger	NJ	2006	Over the top	High injury at all rates ( 2.5, 5 and 10 lb ai per acre)
24834	Switch Grass ( <i>Panicum virgatum</i> )	Field Container	Lieth	CA	2006	Over the top	No injury but unacceptable growth reduction at 2.5, 5 and 10 lb ai per acre
24834	Switch Grass ( <i>Panicum virgatum</i> ) 'Dallas Blues'	Field Container	Mathers (OSU)	OH	2005	Over the top	No injury
24755	Beard-Tongue ( <i>Penstemon</i> sp.) P. digitalis 'Scarlet Queen'	Field Container	Boydston	WA	2005	Over the top	No injury with 2.5, 5.0 and 10 lb ai per acre.
24755	Beard-Tongue ( <i>Penstemon</i> sp.) P. digitalis 'Husker Red'	Field Container	Derr	VA	2005	Over the top	Slight injury at 5 lb ai per acre but plants outgrew injury
24755	Beard-Tongue ( <i>Penstemon</i> sp.) P. X mexicali 'Red Rocks'	Field Container	Lieth	CA	2005	Over the top	Very slight, no effect on plant growth
24755	Beard-Tongue ( <i>Penstemon</i> sp.) P. digitalis 'Husker Red'	Field Container	Neal	NC	2005	Over the top	No injury at 2.5 lb ai per acre, moderate at 5 lb ai per acre, severe at 10 lb ai per acre
26836	Pentas ( <i>Pentas</i> sp.) 'New Look Pink'	Greenhouse	Freiberger	NJ	2006	Over the top	High injury at all rates (2.5, 5 and 10 lb ai per acre)
25300	Phlox, Carolina ( <i>Phlox carolinia</i> ) P. subulata 'Candy Strip'	Field In-Ground	Chen	LA	2005	Over the top	Slight injury with 2.5, 5.0 and 10 lb ai per acre.
33364	Phlox, Fall ( <i>Phlox paniculata</i> )	Field Container	Larson	WA	2022	Over the top	No injury with 2.5, 5, and 10 lb ai per acre.

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
24758	Flax, New Zealand ( <i>Phormium tenax</i> ) <i>P. colinsoi</i>	Field Container	Lieth	CA	2005	Over the top	No injury, significant increase in plant growth
23845	Jacob's Ladder ( <i>Polemonium</i> sp.) 'Heavenly Habit'	Field Container	Boydston	WA	2005	Over the top	No injury to minor injury increasing with rate (2.5, 5.0 and 10 lb ai per acre). However, significant stunting occurred with 10 lb ai per acre.
23845	Jacob's Ladder ( <i>Polemonium</i> sp.) <i>P. reptans</i>	Field Container	Fraelich	GA	2006	Over the top	Slight injury and stunting; no reduction in marketability
23845	Jacob's Ladder ( <i>Polemonium</i> sp.) <i>P. reptans</i> 'Stairway to Heaven'	Field Container	Linderman	OR	2006	Over the top	No injury at 2.5, 5 and 10 lb ai per acre
23845	Jacob's Ladder ( <i>Polemonium</i> sp.) <i>P. 'Caeruleum Blue'</i>	Field Container	Treat	WA	2006	Over the top	Severe injury at 2.5, 5 and 10 lb ai per acre.
24760	Primrose, Fairy ( <i>Primula malacoides</i> )	Field Container	Lieth	CA	2005	Over the top	Significant injury and plant growth suppression
24760	Primrose, Fairy ( <i>Primula malacoides</i> ) 'Prima Carmine Rose'	Field Container	Treat	WA	2006	Over the top	Severe injury (leaf burn and stunting) at all rates (2.5, 5 and 10 lb ai per acre)
24759	Lungwort ( <i>Pulmonaria</i> sp.) <i>P. x cevenensis</i> 'Silver Bouquet'	Field Container	Senesac	NY	2022	Over the top	Minor to moderate injury increasing with rate (2.5, 5.0 10.0 lb ai per acre).
33362	Black-Eyed Susan ( <i>Rudbeckia hirta</i> ) 'Marmalade'	Field In-Ground	Aulakh	CT	2018	Over the top	No injury or growth reduction with 2.6 pt per acre applied twice, moderate with 5.2, and severe with 10.4 pt per acre.
25305	Carolina Wild Petunia ( <i>Ruellia caroliniensis</i> ) <i>R. brittoniana</i> 'Katie'	Field In-Ground	Chen	LA	2005	Over the top	No significant injury with 2.5, 5.0 and 10 lb ai per acre.
24762	Lavender Cotton ( <i>Santolina chamaecyparissus</i> ) <i>S. incana</i>	Field Container	Boydston	WA	2006	Over the top	No injury or growth reduction at all rates (2.5, 5 and 10 lb ai per acre); all plants saleable
24762	Lavender Cotton ( <i>Santolina chamaecyparissus</i> ) 'Compacta'	Field Container	Lieth	CA	2005	Over the top	No injury at 2.5 lb ai per acre, slight injury at 5 lb ai per acre and moderate at 10 lb ai per acre rate; no effect on plant growth
26033	Pincushions ( <i>Scabiosa</i> sp.) 'Blue Diamonds'	Greenhouse	Mickelbart	IN	2006	Over the top	No significant injury or growth reduction at 2.5 lb ai per acre; significant at higher rates
23848	Houseleek, Cobweb ( <i>Sempervivum arachnoideum</i> )	Field Container	Senesac	NY	2004	Over the top	No injury with 2.5, 5.0, 10.0 lb ai per acre.
26842	Houseleek ( <i>Sempervivum</i> sp.)	Greenhouse	Freiberger	NJ	2006	Over the top	No injury after 1st application at all rates (2.5, 5 and 10 lb ai per acre), high at 2X and 4X after 2nd application
23847	Houseleek, Common ( <i>Sempervivum tectorum</i> ) 'Cobweb'	Field Container	Ahrens/Mervosh	CT	2005	Over the top	Slight injury at 2.5 lb ai per acre rate, moderate at 5 lb ai per acre and 10 lb ai per acre.
23847	Houseleek, Common ( <i>Sempervivum tectorum</i> ) 'Sunset'	Field Container	Lieth	CA	2004	Over the top	No injury
23847	Houseleek, Common ( <i>Sempervivum tectorum</i> )	Field Container	Senesac	NY	2004	Over the top	No injury
23851	Goldenrod, Wrinkleleaf ( <i>Solidago rugosa</i> ) 'Goldenrod'	Field Container	Fraelich	GA	2005	Over the top	No injury or growth differences at all rates



PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
23851	Goldenrod, Wrinkleleaf ( <i>Solidago rugosa</i> ) 'Fireworks'	Field Container	Lieth	CA	2004	Over the top	No injury at 2.5 lb ai per acre rate, slight injury at 5 lb ai per acre and 10 lb ai per acre rates
23851	Goldenrod, Wrinkleleaf ( <i>Solidago rugosa</i> )	Field Container	Neal	NC	2004	Over the top	No injury at 2.5 lb ai per acre rate, slight injury at 5 lb ai per acre and 10 lb ai per acre rates
23851	Goldenrod, Wrinkleleaf ( <i>Solidago rugosa</i> )	Field Container	Reding	OH	2005	Over the top	Slight, transient injury at 2.5 lb ai per acre and 5 lb ai per acre rates
23852	Goldenrod, Seaside ( <i>Solidago sempervirens</i> )	Field Container	Fraelich	GA	2005	Over the top	No injury or growth differences at all rates
23852	Goldenrod, Seaside ( <i>Solidago sempervirens</i> )	Field Container	Fraelich	GA	2006	Over the top	No injury or growth differences at all rates
23852	Goldenrod, Seaside ( <i>Solidago sempervirens</i> )	Field Container	Senesac	NY	2004	Over the top	No injury with 2.5, 5.0, 10.0 lb ai per acre.
33378	Goldenrod ( <i>Solidago</i> sp.) <i>S. rugosa</i> 'Fireworks'	Field In-Ground	Aulakh	CT	2018	Over the top	No injury or growth reduction with 2.6, 5.2 and 10.4 pt per acre after 1st applic, moderate after the 2nd application with 2X and 4X.
24763	Goldenrod ( <i>Solidago</i> sp.) <i>S. speciosa</i>	Field Container	Fraelich	GA	2006	Over the top	No injury or growth differences at all rates
24763	Goldenrod ( <i>Solidago</i> sp.) <i>S. sempervirens</i>	Field Container	Linderman	OR	2006	Over the top	No injury at 2.5, 5 and 10 lb ai per acre
24763	Goldenrod ( <i>Solidago</i> sp.) <i>S. cabadensis</i>	Field Container	Treat	WA	2006	Over the top	No injury at 2.5, 5 and 10 lb ai per acre.
23853	Goldenrod, Showy ( <i>Solidago speciosa</i> )	Field Container	Larson	WA	2022	Over the top	No injury with 2.5, 5, and 10 lb ai per acre.
27704	Aster, White Heath ( <i>Symphyotrichum ericoides</i> ) 'Snow Flurry'	Field Container	Chandran	WV	2005	Over the top	No significant injury at 2.5, 5, and 10 lb ai per acre.
23854	Foamflower, Heartleaf ( <i>Tiarella cordifolia</i> )	Field Container	Neal	NC	2004	Over the top	All rates caused significant injury (2.5, 5.0, 10.0 lb ai per acre).
23854	Foamflower, Heartleaf ( <i>Tiarella cordifolia</i> )	Field Container	Senesac	NY	2005	Over the top	Moderate to severe injury increasing with rate (2.5, 5.0, 10.0 lb ai per acre).
24766	Ohio Spiderwort; Bluejacket ( <i>Tradescantia ohioensis</i> ) 'Osprey'	Field Container	Boydston	WA	2005	Over the top	All rates (2.5, 5.0 and 10 lb ai per acre) caused moderate to severe injury, stunting and delayed flowering; treated plants not saleable.
26843	Spiderwort ( <i>Tradescantia</i> sp.)	Greenhouse	Freiberger	NJ	2006	Over the top	Slight injury after 1st, high injury after 2nd application at all rates (2.5, 5 and 10 lb ai per acre)
24765	Spiderwort ( <i>Tradescantia x andersoniana</i> ) 'Concord Grape'	Field Container	Derr	VA	2005	Over the top	No injury at 2.5 lb ai per acre slight injury at 5 lb ai per acre, moderate at 10 lb ai per acre
24765	Spiderwort ( <i>Tradescantia x andersoniana</i> ) 'Baby Bunny Bellies'	Field Container	Lieth	CA	2006	Over the top	Unacceptable injury and growth reduction at 2.5, 5 and 10 lb ai per acre
24765	Spiderwort ( <i>Tradescantia x andersoniana</i> ) 'Sweet Kate'	Field Container	Mathers (OSU)	OH	2005	Over the top	All rates (2.5, 5.0, 10.0 lb ai per acre) caused severe injury (distorted and dead plants)
25306	Vervain ( <i>Verbena</i> sp.) <i>V. canadensis</i> 'Homestead Purple'	Field In-Ground	Chen	LA	2005	Over the top	Moderate injury with 2.5, 5.0 and 10 lb ai per acre; plants recovered in a short time

PR#	Crop	Production Site	Researcher	State	Year	Application Type	Results
24689	Vervain (Verbena sp.) V. canadensis 'Homestead Purple'	Field Container	Neal	NC	2005	Over the top	No injury at 2.5 lb ai per acre rate, moderate at 5 lb ai per acre rate and severe at 10 lb ai per acre
23855	Ironweed, New York (Vernonia noveboracensis)	Field Container	Neal	NC	2004	Over the top	No injury at 2.5 lb ai per acre rate, slight injury at 5 lb ai per acre and 10 lb ai per acre rates
23855	Ironweed, New York (Vernonia noveboracensis)	Field Container	Senesac	NY	2004	Over the top	No injury with 2.5, 5.0, 10.0 lb ai per acre.
13131	Strawflower, Bracted (Xerochrysum bracteatum)	Field In-Ground	Sellmer	PA	2001	Over the top	
33358	Zinnia (Zinnia sp.) 'Purple Prince'	Field In-Ground	Aulakh	CT	2018	Over the top	Minor injury with good recovery, and no growth reduction, with 2.6, 5.2 and 10.4 pt per acre applied twice.

## Label Suggestions

We recommend that the 5 crops exhibiting no injury in these experiments be added to the current Pennant Magnum label for over-the-top applications.

*Ligularia stenocephala*  
*Linum perenne*  
*Nepeta faassenii*  
*Penstemon digitalis*  
*Solidago rugosa*

We also suggest that the 22 plant species / genera exhibiting significant injury with over-the-top applications be added to a sensitive plant list.

*Agastache aurantiaca*  
*Agastache rugosa x foeniculum*  
*Alchemilla mollis*  
*Amsonia hubrichtii*  
*Amsonia tabernaemontana*  
*Armeria maritima*  
*Bergenia cordifolia*  
*Chelone lyonii*  
*Chrysogonum virginianum*  
*Delosperma nubigenum*  
*Digitalis thapsi*  
*Echinacea purpurea*  
*Epimedium x rubrum*  
*Gerbera jamesonii*  
*Heuchera sanguinea*  
*Nepeta nervosa*  
*Origanum libanoticum*  
*Pentas sp.*  
*Primula malacoides*  
*Tiarella cordifolia*  
*Tradescantia andersoniana*  
*Tradescantia ohiensis*

## Appendix 1: Contributing Researchers

Dr. John Ahrens <i>(retired)</i>	Connecticut Agricultural Experiment Station Valley Laboratory 153 Cook Hill Road, P.O. Box 248 Windsor, CT
Dr. Jatindee Aulakh	Connecticut Agricultural Experiment Station 153 Cook Hill Road Windsor, CT 06095
Dr. Rick Boydston	USDA-ARS IAREC 24106 N Bunn Road Prosser, WA 99350
Dr. Rakesh S. Chandran	West Virginia University 1076 Agricultural Science Building P. O. Box 6108 Morgantown, WV 26506-6108
Dr. Yan Chen	Louisiana State University AgCenter Hammond Research Station 21549 Old Covington Hwy. Hammond, LA 70403
Dr. Jeffrey Derr	Hampton Roads Ag. Exp. Station 1444 Diamond Springs Road Virginia Beach, VA 23455
Mr. Ben Fraelich	USDA-ARS CPES P.O. Box 748 Tifton, GA 31793
Mr. Tom Freiberger <i>(retired)</i>	Rutgers University Cream Ridge Experiment Station 283 Rt. 539 Cream Ridge, NJ 08514
Dr. Charles Gilliam <i>(retired)</i>	Auburn University Department of Horticulture 101 Funchess Hall Auburn, AL 36849

Dr. Jim Klett  
Colorado State University  
Department of Horticulture and Landscape Architecture  
Fort Collins, CO 80523

Mr. Duane Larson  
USDA-ARS  
4230 Konnawac Pass Road  
Wapato, WA, 98941

Dr. Heiner Lieth  
Department of Plant Sciences  
University of California  
One Shield Avenue  
Davis, CA 95616

Dr. Bob Linderman  
*(retired)*  
Horticultural Crops Research Lab  
USDA-ARS  
3420 NW Orchard Ave.  
Corvallis, OR 97330

Dr. Hannah Mathers  
*(past affiliate)*  
The Ohio State University  
Dept. Hort. and Crop Science  
2001 Fyffe Ct.  
Columbus, OH 43210

Dr. Michael Mickelbart  
Purdue University  
Dept. Hort. and Landscape Architecture  
625 Agriculture Mall Dr.  
West Lafayette, IN 47907

Dr. Tim Miller  
Washington State University  
16650 State Route 536  
Mount Vernon, WA 98273-9761

Dr. Joe Neal  
North Carolina State University  
Department of Horticultural Science  
262 Kilgore Hall  
Box 7609, NCSU  
Raleigh, NC 27695-7609

Dr. Michael Reding  
USDA-ARS  
Hort Insects Lab  
1680 Madison Ave.  
Wooster, OH, 44691

Dr. Andy Senesac

Long Island Horticultural Research Laboratory  
39 Sound Avenue  
Riverhead, NY 11901

Mr. Thomas L. Treat  
*(retired)*

USDA-ARS  
Yakima Agricultural Research Laboratory  
5230 Konnowac Pass Road  
Wapato, WA 98951